

Characters, Character sheaves and Beyond

Toshiaki Shoji

Nagoya University

March 14, 2012

Kostka polynomials $K_{\lambda,\mu}(t)$

$\lambda = (\lambda_1, \dots, \lambda_k) : \text{partition of } n$

$$\lambda_i \in \mathbb{Z}_{\geq 0}, \quad \lambda_1 \geq \dots \geq \lambda_k \geq 0, \quad \sum_i \lambda_i = n$$

$\mathcal{P}_n = \{\text{partitions of } n\}$

$s_\lambda(x) = s_\lambda(x_1, \dots, x_k) \in \mathbb{Z}[x_1, \dots, x_k] : \text{Schur function}$

$$s_\lambda(x) = \det(x_i^{\lambda_j+k-j})_{1 \leq i, j \leq k} / \det(x_i^{k-j})_{1 \leq i, j \leq k}$$

$P_\lambda(x; t) = P_\lambda(x_1, \dots, x_k; t) \in \mathbb{Z}[x_1, \dots, x_k; t] : \text{Hall-Littlewood function}$

$$P_\lambda(x_1, \dots, x_k; t) = \sum_{w \in S_k / S_k^\lambda} w \left(x_1^{\lambda_1} \dots x_k^{\lambda_k} \prod_{\lambda_i > \lambda_j} \frac{x_i - tx_j}{x_i - x_j} \right)$$

Take $k \gg 0$

$\{s_\lambda(x) \mid \lambda \in \mathcal{P}_n\}, \{P_\lambda(x; t) \mid \lambda \in \mathcal{P}_n\}$ are basis of the space of homog. symmetric polynomials of degree n in $\mathbb{Z}[x_1, \dots, x_k : t]$

For $\lambda, \mu \in \mathcal{P}_n$, $K_{\lambda, \mu}(t)$: **Kostka polynomial** defined by

$$s_\lambda(x) = \sum_{\mu \in \mathcal{P}_n} K_{\lambda, \mu}(t) P_\mu(x; t)$$

$$K_{\lambda, \mu}(t) \in \mathbb{Z}[t]$$

$(K_{\lambda, \mu}(t))_{\lambda, \mu \in \mathcal{P}_n}$: transition matrix of two basis $\{s_\lambda(x)\}, \{P_\mu(x; t)\}$

Geometric realization of Kostka polynomials

In 1981, Lusztig gave a geometric realization of Kostka polynomials in connection with the closure of nilpotent orbits.

$$V = \mathbb{C}^n, \quad G = GL(V)$$

$$\mathcal{N} = \{x \in \text{End}(V) \mid x : \text{nilpotent}\} : \text{nilpotent cone}$$

$$\mathcal{P}_n \simeq \mathcal{N}/G$$

$$\lambda \leftrightarrow G\text{-orbit } \mathcal{O}_\lambda \ni x : \text{Jordan type } \lambda$$

• Closure relations :

$$\overline{\mathcal{O}}_\lambda = \coprod_{\mu \leq \lambda} \mathcal{O}_\mu \quad (\overline{\mathcal{O}}_\lambda : \text{Zariski closure of } \mathcal{O}_\lambda)$$

dominance order on \mathcal{P}_n

$$\text{For } \lambda = (\lambda_1, \lambda_2, \dots, \lambda_k), \quad \mu = (\mu_1, \mu_2, \dots, \mu_k),$$

$$\mu \leq \lambda \Leftrightarrow \sum_{i=1}^j \mu_i \leq \sum_{i=1}^j \lambda_i \quad \text{for each } j.$$

Notation: $n(\lambda) = \sum_{i \geq 1} (i-1)\lambda_i$

Define $\tilde{K}_{\lambda,\mu}(t) = t^{n(\mu)} K_{\lambda,\mu}(t^{-1})$: **modified Kostka polynomial**

$K = \text{IC}(\overline{\mathcal{O}}_\lambda, \mathbb{C})$: Intersection cohomology complex

$$K : \cdots \longrightarrow K_{i-1} \xrightarrow{d_{i-1}} K_i \xrightarrow{d_i} K_{i+1} \xrightarrow{d_{i+1}} \cdots$$

$K = (K_i)$: bounded complex of \mathbb{C} -sheaves on $\overline{\mathcal{O}}_\lambda$

$\mathcal{H}^i K = \text{Ker } d_i / \text{Im } d_{i-1}$: i -th cohomology sheaf

$\mathcal{H}_x^i K$: the stalk at $x \in \overline{\mathcal{O}}_\lambda$ of $\mathcal{H}^i K$ (finite dim. vector space over \mathbb{C})

Known fact : $\mathcal{H}^i K = 0$ for odd i .

Theorem (Lusztig)

For $x \in \mathcal{O}_\mu$,

$$\tilde{K}_{\lambda,\mu}(t) = t^{n(\lambda)} \sum_{i \geq 0} (\dim_{\mathbb{C}} \mathcal{H}_x^{2i} K) t^i$$

In particular, $K_{\lambda,\mu}[t] \in \mathbb{Z}_{\geq 0}[t]$. (theorem of Lascoux-Schützenberger)

Representation theory of $GL_n(\mathbb{F}_q)$

\mathbb{F}_q : finite field of q elements with $\text{ch } \mathbb{F}_q = p$

$\overline{\mathbb{F}}_q$: algebraic closure of \mathbb{F}_q

$$G = GL_n(\overline{\mathbb{F}}_q) \supset B = \left\{ \begin{pmatrix} * & \cdots & * \\ 0 & \ddots & \vdots \\ 0 & 0 & * \end{pmatrix} \right\} \supset U = \left\{ \begin{pmatrix} 1 & \cdots & * \\ 0 & \ddots & \vdots \\ 0 & 0 & 1 \end{pmatrix} \right\}$$

B : Borel subgroup, U : maximal unipotent subgroup

$F : G \rightarrow G, (g_{ij}) \mapsto (g_{ij}^q)$: Frobenius map

$G^F = \{g \in G \mid F(g) = g\} = G(\mathbb{F}_q)$: finite subgroup

$\text{Ind}_{B^F}^{G^F} 1$: the character of G^F obtained by inducing up 1_{B^F}

$$\text{Ind}_{B^F}^{G^F} 1 = \sum_{\lambda \in \mathcal{P}_n} (\deg \chi^\lambda) \rho^\lambda,$$

ρ^λ : irreducible character of G^F corresp. to $\chi^\lambda \in S_n^\wedge \simeq \mathcal{P}_n$.

$G_{\text{uni}} = \{g \in G \mid u: \text{unipotent}\} \subset G, \quad G_{\text{uni}} \simeq \mathcal{N}, u \leftrightarrow u - 1$

• $G_{\text{uni}}/G \simeq \mathcal{P}_n, \quad \mathcal{O}_\lambda \leftrightarrow \lambda$

$\mathcal{O}_\lambda : F\text{-stable} \implies \mathcal{O}_\lambda^F : \text{single } G^F\text{-orbit}, u_\lambda \in \mathcal{O}_\lambda^F$

Theorem (Green)

$$\rho^\lambda(u_\mu) = \tilde{K}_{\lambda,\mu}(q)$$

Remark : Lusztig's result \implies the character values of ρ^λ at **unipotent elements** are described in terms of intersection cohomology complex.

Theory of character sheaves \implies describes **all the character values** of ρ^λ in terms of certain simple perverse sheaves.

Character sheaves on GL_n

X : alg. variety over \mathbb{F}_q with Frobenius map $F : X \rightarrow X$

K : perverse sheaf on X , K : F -stable $\Leftrightarrow F^*K \simeq K$.

For F -stable perverse sheaf K , fix $\varphi : F^*K \rightarrow K$ isomorphism

Define $\chi_{K,\varphi} : X^F \rightarrow \bar{\mathbb{Q}}_l$ by

$$\chi_{K,\varphi}(x) = \sum_i (-1)^i \text{Tr}(\varphi, \mathcal{H}_x^i K) \quad (x \in X^F)$$

$\chi_{K,\varphi}$: **Characteristic function** of K with respect to φ .

• If K : G -equiv. perverse sheaf $\Rightarrow \chi_{K,\varphi} : G^F$ -invariant function on X^F .

Lusztig : All the irreducible characters of $G^F = GL_n(\mathbb{F}_q)$ are obtained as characteristic functions of certain G -equivariant F -stable simple perverse sheaves (i.e., **character sheaves**) of G

Representation theory of finite reductive groups

Green (1955) : classified all irreducible representations of G^F on \mathbb{C} (or $\bar{\mathbb{Q}}_l$), and determined irreducible characters

basic tool : $R_T^G(\theta)$ Green's basic function
(Deligne-Lusztig's virtual character)

$$Q_T^G := R_T^G(\theta)|_{G_{\text{uni}}^F} : G_{\text{uni}}^F \rightarrow \bar{\mathbb{Q}}_l \quad \text{Green function}$$

- $\pm R_T^G(\theta)$: irreducible character for generic pair (T, θ)
- Any irreducible character is a linear combination of $R_T^G(\theta)$
- Computation of $R_T^G(\theta) \iff$ computation of Green functions

Green functions are described by Kostka polynomials !!

Deligne-Lusztig (1976) : Deligne-Lusztig's virtual rep. $R_T^G(\theta)$ for connected reductive groups by using ℓ -adic cohomology theory

Lusztig (1980's) : Classification of irreducible representations for connected reductive groups

Lusztig (1985) : Theory of character sheaves (geometric theory of characters of reductive groups)

Lusztig's conjecture : Uniform algorithm of computing irreducible characters (in principle)

S (1995) : Solved Lusztig's conjecture in the case where the center of G is connected

Bonnafe, S, Waldspurger (2004 ~) : Lusztig's conjecture for disconnected center case for SL_n, Sp_{2n}, SO_{2n} (open in general)

Enhanced nilpotent cone

$\lambda = (\lambda^{(1)}, \dots, \lambda^{(r)}), \quad \sum_{i=1}^r |\lambda^{(i)}| = n : r\text{-partition of } n$

$\mathcal{P}_{n,r}$: the set of r -partitions of n

For $\lambda, \mu \in \mathcal{P}_{n,r}$, one can define $K_{\lambda, \mu}(t) \in \mathbb{Q}(t)$: **Kostka functions associated to complex reflection groups.**

Achar-Henderson (2008) : geometric realization
of Kostka functions for $r = 2$

$V = \mathbb{C}^n, \quad \mathcal{N}$: nilpotent cone

$\mathcal{N} \times V$: **enhanced nilpotent cone**, diagonal action of $G = GL(V)$

Achar-Henderson, Travkin :

$$(\mathcal{N} \times V)/G \simeq \mathcal{P}_{n,2}, \quad \mathcal{O}_\lambda \leftrightarrow \lambda$$

• **Closure relations**

$$\overline{\mathcal{O}_\lambda} = \coprod_{\mu \leq \lambda} \mathcal{O}_\mu$$

$K = \text{IC}(\overline{\mathcal{O}}_\lambda, \mathbb{C})$: Intersection cohomology complex

Theorem (Achar-Henderson)

$\mathcal{H}^i K = 0$ for odd i . For $\lambda, \mu \in \mathcal{P}_{n,2}$, and $(x, v) \in \mathcal{O}_\mu \subseteq \overline{\mathcal{O}}_\lambda$,

$$t^{a(\lambda)} \sum_{i \geq 0} (\dim_{\mathbb{C}} \mathcal{H}_{(x,v)}^{2i} K) t^{2i} = \tilde{K}_{\lambda, \mu}(t),$$

where $a(\lambda) = 2n(\lambda^{(1)}) + 2n(\lambda^{(2)}) + |\lambda^{(2)}|$ for $\lambda = (\lambda^{(1)}, \lambda^{(2)})$.

$\mathcal{N} \times V \rightsquigarrow G_{\text{uni}} \times V \hookrightarrow G \times V$ (over $\overline{\mathbb{F}}_q$) : diagonal action of $G = GL(V)$

Finkelberg-Ginzburg-Travkin (2008) : Theory of character sheaves on $G \times V$ (certain G -equiv. simple perverse sheaves)

\implies **“character table”** of $(G \times V)^F$

S (2010) : Generalization to $(G \times V^{r-1})^F$, in connection with Kostka functions assoc. to $\mathcal{P}_{n,r}$ (in progress)

Finite symmetric space $GL_{2n}(\mathbb{F}_q)/Sp_{2n}(\mathbb{F}_q)$

$$G = GL(V) \simeq GL_{2n}(\overline{\mathbb{F}}_q), \quad V = (\overline{\mathbb{F}}_q)^{2n}, \quad \text{ch } \mathbb{F}_q \neq 2$$

$$\theta : G \rightarrow G, \theta(g) = J^{-1}({}^t g^{-1})J : \text{involution, } J = \begin{pmatrix} 0 & I_n \\ -I_n & 0 \end{pmatrix}$$

$$K := \{g \in G \mid \theta(g) = g\} \simeq Sp_{2n}(\overline{\mathbb{F}}_q) \quad G/K : \text{symmetric space over } \overline{\mathbb{F}}_q$$

$$G^F \simeq GL_{2n}(\mathbb{F}_q) \supset Sp_{2n}(\mathbb{F}_q) \simeq K^F$$

G^F acts on $G^F/K^F \rightsquigarrow 1_{K^F}^{G^F}$: induced representation

$H(G^F, K^F) := \text{End}_{G^F}(1_{K^F}^{G^F})$: **Hecke algebra** asoc. to (G^F, K^F)

- $H(G^F, K^F)$: commutative algebra
- $H(G^F, K^F)^\wedge$: natural labeling by $(GL_n^F)^\wedge$
- $K^F \backslash G^F / K^F$: natural labeling by $\{ \text{conj. classes of } GL_n^F \}$

Theorem (Bannai-Kawanka-Song, 1990)

The character table of $H(G^F, K^F)$ can be obtained from the character table of GL_n^F by replacing $q \mapsto q^2$.

More precisely, there exist **basic functions, Green functions** assoc. to $H(G^F, K^F)$, which have the same role as those for $GL_n(\mathbb{F}_q)$.

Geometric setting for G/K

$$\begin{aligned} G^{\iota\theta} &= \{g \in G \mid \theta(g) = g^{-1}\} \\ &= \{g\theta(g)^{-1} \mid g \in G\}, \end{aligned}$$

where $\iota : G \rightarrow G, g \mapsto g^{-1}$.

The map $G \rightarrow G, g \mapsto g\theta(g)^{-1}$ gives isom. $G/K \simeq G^{\iota\theta}$.

K acts by left mult $\curvearrowright G/K \simeq G^{\iota\theta} \curvearrowleft K$ acts by conjugation.

$$K \backslash G/K \simeq \{K\text{-conjugates of } G^{\iota\theta}\}$$

- Irred. character of $H(G^F, K^F) \iff K^F$ -inv. function on $(G^{\iota\theta})^F$
- character sheaves $\iff K$ -equiv. simple perverse sheaves on $G^{\iota\theta}$

Henderson : Geometric reconstruction of BKS (not complete)

Lie algebra analogue

$\mathfrak{g} = \mathfrak{gl}_{2n}$, $\theta : \mathfrak{g} \rightarrow \mathfrak{g} : \text{involution}$, $\mathfrak{g} = \mathfrak{g}^{\theta} \oplus \mathfrak{g}^{-\theta}$,

$\mathfrak{g}^{\pm\theta} = \{x \in \mathfrak{g} \mid \theta(x) = \pm x\}$, K -stable subspace of \mathfrak{g}

$\mathfrak{g}_{\text{nil}}^{-\theta} = \mathfrak{g}^{-\theta} \cap \mathcal{N}_{\mathfrak{g}}$: analogue of nilpotent cone \mathcal{N} , K -stable subset of $\mathfrak{g}^{-\theta}$

$$\mathfrak{g}_{\text{nil}}^{-\theta}/K \simeq \mathcal{P}_n, \quad \mathcal{O}_{\lambda} \leftrightarrow \lambda$$


Theorem (Henderson + BKS, 2008)

Let $K = \text{IC}(\overline{\mathcal{O}}_{\lambda}, \overline{\mathcal{Q}}_l)$, $x \in \mathcal{O}_{\mu} \subset \overline{\mathcal{O}}_{\lambda}$. Then $\mathcal{H}^i K = 0$ unless $i \equiv 0 \pmod{4}$, and

$$t^{2n(\lambda)} \sum_{i \geq 0} (\dim \mathcal{H}_x^{4i} K) t^{2i} = \tilde{K}_{\lambda, \mu}(t^2)$$

Exotic symmetric space $GL_{2n}/Sp_{2n} \times V$

(Joint work with K. Sorlin)



$G = GL(V) \simeq GL_{2n}(\overline{\mathbb{F}}_q)$, $\dim V = 2n$, $K = G^\theta$.
 $G^{\iota\theta} \times V : K$ acts diagonally

Problem

- Find a good class of K -equivariant simple perverse sheaves on $G^{\iota\theta} \times V$, i.e., “**character sheaves**” on $G^{\iota\theta} \times V$
- Find a good basis of K^F -equivariant functions on $(G^{\iota\theta} \times V)^F$, i.e., “**irreducible characters**” of $(G^{\iota\theta} \times V)^F$, and compute their values, i.e., computation of the “**character table**”

Remark : $\mathcal{X}_{\text{uni}} := G_{\text{uni}}^{\iota\theta} \times V \simeq \mathfrak{g}_{\text{nil}}^{-\theta} \times V$: **Kato's exotic nilcone**

Kato $(\mathfrak{g}_{\text{nil}}^{-\theta} \times V)/K \simeq \mathcal{P}_{n,2}$, $\mathcal{O}_\mu \leftrightarrow \mu \in \mathcal{P}_{n,2}$

Natural bijection with GL_n -orbits of enhanced nilcone, compatible with closure relations (**Achar-Henderson**)

Springer correspondence

$B = TU$: θ -stable Borel subgroup, maximal torus, unipotent radical

$M_0 \subset M_1 \subset \cdots \subset M_n$: isotropic flag stable by B

$W_n = N_K(T^\theta)/T^\theta$: Weyl group of type C_n

$$\tilde{\mathcal{X}}_{\text{uni}} = \{(x, v, gB^\theta) \in G_{\text{uni}}^{\iota\theta} \times V \times K/B^\theta \mid (g^{-1}xg, g^{-1}v) \in U^{\iota\theta} \times M_n\}$$

$$\pi_1 : \tilde{\mathcal{X}}_{\text{uni}} \rightarrow \mathcal{X}_{\text{uni}}, \quad (x, v, gB^\theta) \mapsto (x, v)$$

Theorem 1 (Springer correspondence)

$(\pi_1)_* \bar{\mathbb{Q}}_l$ is a semisimple perverse sheaf on \mathcal{X}_{uni} with W_n -action, is decomposed as

$$(\pi_1)_* \bar{\mathbb{Q}}_l[\dim \mathcal{X}_{\text{uni}}] \simeq \bigoplus_{\mu \in \mathcal{P}_{n,2}} V_\mu \otimes \text{IC}(\bar{\mathcal{O}}_{\mu^\bullet}, \bar{\mathbb{Q}}_l)[\dim \mathcal{O}_{\mu^\bullet}],$$

V_μ : standard irred. W_n -module, $\mathcal{O}_{\mu^\bullet} \mapsto V_\mu$ gives bijection $\mathcal{X}_{\text{uni}}/K \simeq W_n^\wedge$

Remark : Theorem 1 was first proved by Kato for the exotic nilcone by using Ginzburg theory on affine Hecke algebras. We give an alternate proof based on the theory of character sheaves

$$T_{\text{reg}}^{\iota\theta} = \{t = \text{Diag}(t_1, \dots, t_n, t_1, \dots, t_n) \mid t_i \neq t_j\}$$

$$\tilde{G}_{\text{reg}}^{\iota\theta} = \{(x, gB^\theta) \in G^{\iota\theta} \times K/B^\theta \mid g^{-1}xg \in T_{\text{reg}}^{\iota\theta}\}$$

$$\psi_0 : \tilde{G}_{\text{reg}}^{\iota\theta} \rightarrow G_{\text{reg}}^{\iota\theta} = \bigcup_{g \in K} g(T_{\text{reg}}^{\iota\theta})g^{-1}, \quad (x, gB^\theta) \mapsto x$$

$$\tilde{G}_{\text{reg}}^{\iota\theta} \simeq K \times_{(Z_K(T^{\iota\theta}) \cap B^\theta)} T_{\text{reg}}^{\iota\theta} \xrightarrow{\xi} K \times_{Z_K(T^{\iota\theta})} T_{\text{reg}}^{\iota\theta} \xrightarrow{\eta} G_{\text{reg}}^{\iota\theta}$$

- η is a finite Galois covering with group $S_n \simeq N_K(T^{\iota\theta})/Z_K(T^{\iota\theta})$
- ξ is a \mathbb{P}_1^n -bundle, with $\mathbb{P}_1^n \simeq (SL_2/B_2)^n$

$$(\psi_0)_* \bar{Q}_I \simeq H^\bullet(\mathbb{P}_1^n, \bar{Q}_I) \otimes \eta_* \bar{Q}_I$$

- η : finite Galois covering $\implies \eta_* \bar{Q}_I$ has a natural action of S_n
- $(\mathbb{Z}/2\mathbb{Z})^n$: Weyl group of $(SL_2)^n$ acts on $H^\bullet(\mathbb{P}_1^n, \bar{Q}_I)$
- $(\psi_0)_* \bar{Q}_I$ has a natural action of $W_n = S_n \ltimes (\mathbb{Z}/2\mathbb{Z})^n$

For $0 \leq m \leq n$,

$$\tilde{\mathcal{Y}}_m = \{(x, v, gB^\theta) \in G_{\text{reg}}^{\iota\theta} \times V \times K/B^\theta \mid (g^{-1}xg, g^{-1}v) \in B_{\text{reg}}^{\iota\theta} \times M_m\}$$

$$\mathcal{Y}_m = \bigcup_{g \in K} g(B_{\text{reg}}^{\iota\theta} \times M_m)$$

$$\psi : \tilde{\mathcal{Y}}_n \rightarrow \mathcal{Y}_n = G_{\text{reg}}^{\iota\theta} \times V, \quad (x, v, gB^\theta) \mapsto (x, v)$$

Proposition 1

$$\psi_* \bar{Q}_I[\dim \mathcal{Y}_n] \simeq \bigoplus_{\mu \in \mathcal{P}_{n,2}} V_\mu \otimes \text{IC}(\mathcal{Y}_{m(\mu)}, \mathcal{L}_\mu)[\dim \mathcal{Y}_{m(\mu)}],$$

$m(\mu) = |\mu^{(1)}|$ for $\mu = (\mu^{(1)}, \mu^{(2)})$, \mathcal{L}_μ : simple local system on $\mathcal{Y}_{m(\mu)}^0$.

For $0 \leq m \leq n$

$$\tilde{\mathcal{X}}_m = \{(x, v, gB^\theta) \in G^{l\theta} \times V \times K/B^\theta \mid (g^{-1}xg, g^{-1}v) \in B^{l\theta} \times M_m\}$$

$$\mathcal{X}_m = \bigcup_{g \in K} g(B^{l\theta} \times M_m)$$

$$\pi: \tilde{\mathcal{X}}_n \rightarrow \mathcal{X}_n = G^{l\theta} \times V, \quad (x, v, gB^{l\theta}) \mapsto (x, v)$$

\mathcal{Y}_m is open dense in \mathcal{X}_m , $\pi_* \bar{\mathbb{Q}}_l|_{\mathcal{X}_{\text{uni}}} \simeq (\pi_1)_* \bar{\mathbb{Q}}_l$

Proposition 2

① $\pi_* \bar{\mathbb{Q}}_l$ is equipped with W_n action, is decomposed as

$$\pi_* \bar{\mathbb{Q}}_l[\dim \mathcal{X}_n] \simeq \bigoplus_{\mu \in \mathcal{P}_{n,2}} V_\mu \otimes \text{IC}(\mathcal{X}_{m(\mu)}, \mathcal{L}_\mu)[\dim \mathcal{X}_{m(\mu)}]$$

② $\text{IC}(\mathcal{X}_{m(\mu)}, \mathcal{L}_\mu)|_{\mathcal{X}_{\text{uni}}} \simeq \text{IC}(\bar{\mathcal{O}}_{\mu^\bullet}, \bar{\mathbb{Q}}_l)[a]$ for some $\mu^\bullet \in \mathcal{P}_{n,2}$,
where $a = \dim \mathcal{O}_{\mu^\bullet} - \dim \mathcal{X}_{\text{uni}} - \dim \mathcal{X}_{m(\mu)} + \dim \mathcal{X}_n$.

Theorem 2 (explicit correspondence)

Under the notation of Theorem 1, we have $\mu^\bullet = \mu$. Hence the Springer correspondence is given by $\mathcal{O}_\mu \leftrightarrow V_\mu$.

Theorem 2 was proved by Kato. Our proof uses “restriction theorem”.

$P = LU_P$: θ -stable parabolic subgroup of G s.t. $L^\theta \simeq GL_1 \times Sp_{2n-2}$.
 $V_1 \oplus V' \subset V$, $GL_1 = GL(V_1)$, $Sp_{2n-2} = Sp(V')$.

For $z = (x, v) \in G_{\text{uni}}^{\iota\theta} \times V$, $z' = (x', v') \in L_{\text{uni}}^{\iota\theta} \times V'$,

$$Y_{z,z'} = \{g \in K \mid g^{-1}xg \in x'U_P^{\iota\theta}, g^{-1}v \in v' + V_1\}$$

Put $d_{z,z'} = (\dim Z_K(z) - \dim Z_{L^\theta}(z'))/2 + \dim U_P^\theta$

Restriction Theorem

Let ρ_z^G irred. rep. of W_n corresp. to $z \in \mathcal{O}$, $\rho_{z'}^L$, irred. rep. of W_{n-1} corresp. to $z' \in \mathcal{O}'$. Then $\langle \rho_z^G, \rho_{z'}^L \rangle_{W_{n-1}}$ coincides with the number of irreducible components of $Y_{z,z'}$ with dimension $d_{z,z'}$

Kostka polynomials and exotic nilcone

For $\mathcal{O}_\lambda \subset \mathfrak{g}_{\text{uni}}^{-\theta} \times V$, let $K = \text{IC}(\overline{\mathcal{O}}_\lambda, \overline{\mathcal{Q}}_l)$.

Conjecture (Achar-Henderson)

- $\mathcal{H}^i K = 0$ unless $i \equiv 0 \pmod{4}$.
- For $(x, v) \in \mathcal{O}_\mu \subset \overline{\mathcal{O}}_\lambda$, $t^{a(\lambda)} \sum_i \dim(\mathcal{H}_{(x,v)}^{4i} K) t^{2i} = \tilde{K}_{\lambda, \mu}(t)$

W_n acts on $\mathbb{C}[x] = \mathbb{C}[x_1, \dots, x_n] \supset V_\lambda$: Specht module

Define $R^\lambda = \mathbb{C}[x]/I_\lambda$, where $I_\lambda = \{P \in \mathbb{C}[x] \mid P(\partial)f = 0 \forall f \in V_\lambda\}$

$R^\lambda = \bigoplus_{i \geq 0} R_i^\lambda$: Graded W_n -module

Conjectue (S) $\sum_{i \geq 0} \langle R_i^\lambda, \chi^\mu \rangle t^i = \tilde{K}_{\lambda, \mu}(t)$

Remark Recently Kato proved Conjecture (S)

Conjecture (S) + his another result \implies Conjecture (AH)

Future Problem

- Discuss the case for $G^{\iota\theta} \times V$ with $\text{ch } \mathbb{F}_q = 2$. Known by Kato that there exists an interesting relationship with Springer correspondence for symplectic groups with even characteristic.
- Extension to the case $G^{\iota\theta} \times V^{r-1}$ for $r \geq 2$, and discuss the relationship with Kostka functions associated to complex reflection groups.
- Extension to the general symmetric space.
 $\theta : G \rightarrow G$ involution, $K = G^\theta$. Consider the variety V with K action such that the number of K -orbits on $G^{\iota\theta} \times V$ is finite. Develop the theory of character sheaves on $G^{\iota\theta} \times V$.