

Addendum to "Termination of 4-fold canonical flips"

By

Osamu FUJINO *

Abstract

The definition of the weighted version of difficulty in "Termination of 4-fold canonical flips" is nonsense. In this paper, we explain why it is meaningless and how to correct it. Anyway, any sequence of 4-fold canonical flips terminates: Theorem 5.2.

§1. Introduction

Professor Alexeev pointed out that Lemma 2.1 in [F], which is a copy of $[K^+, (4.12.2.1)]$, is wrong. Therefore, the weighted version of difficulty $d_{S,b}(X, B)$ in Definition 2.3 in [F] is infinite if $b < \max_j \{b_j\}$. So, the proof in [F] is nonsense. In this paper, we change the definition of $d_{S,b}(X, B)$ to make it finite when (X, B) is canonical and B has no reduced components, that is, the round down $\lfloor B \rfloor = 0$. Roughly speaking, in [F, Definition 2.3] we exclude valuations obtained by one blow-up along generic points of codimension two subvarieties when we count valuations with small discrepancies. In this paper, we exclude valuations whose centers are codimension two subvarieties with good properties. By this change, the new version of $d_{S,b}(X, B)$ defined in Definition 4.4 becomes finite and the arguments in [F] work without any changes. Proposition 3.1 is a key result in this paper. Note that the problems in [F] are not in the arguments but in the definitions. As mentioned above,

Communicated by ?????.

2000 Mathematics Subject Classification(s): Primary 14E05; Secondary 14J35.

*Graduate School of Mathematics, Nagoya University, Chikusa-ku Nagoya 464-8602, Japan.
Current address: Institute for Advanced Study, Einstein Drive, Princeton NJ 08540, USA.

we have to assume $\perp B \perp = 0$ to make $d_{S,b}(X, B)$ finite. Thus the main theorem: Theorem 1.1 in [F] becomes slightly weaker. However, this assumption is harmless for applications if we use the special termination theorem (see [Fj2]). For the precise statements of the termination theorems, see Theorems 5.1 and 5.2 below. Anyway, *any sequence of 4-fold canonical flips terminates*.

Acknowledgments. I would like to thank Professor Valery Alexeev, who pointed out an error in [F], gave me comments, and obtained the same correction independently. This paper was written in the Institute for Advanced Study. I am grateful to it for its hospitality. I was partially supported by a grant from the National Science Foundation: DMS-0111298.

We summarize the contents of this paper. In Section 2, we describe the errors in discrepancy lemmas in [K⁺, 4.12]. In Section 3, we formulate a new discrepancy lemma. Proposition 3.1 is the main result in this paper. In Section 4, we explain how to modify the definition of the weighted version of difficulty. Section 5 is devoted to the statements of the termination of 4-fold canonical flips. We will use the same notation as in [F] throughout this paper.

§2. Errors in discrepancy lemmas

The following example contradicts [F, Lemma 2.1], which is a copy of [K⁺, (4.12.2.1)].

Example 2.1. Let $X = \mathbb{P}^2$, $B = \frac{2}{3}L$, where L is a line on X . Let P be any point on L . First, blow up X at P . Then we obtain an exceptional divisor E_P such that $a(E_P, X, B) = \frac{1}{3}$. Let L' be the strict transform of L . Next, take a blow-up at $L' \cap E_P$. Then we obtain an exceptional divisor F_P whose discrepancy $a(F_P, X, B) = \frac{2}{3}$. Note that this F_P is *essential* in the notation in [F, Definition 2.1]. On the other hand, it is easy to see that $\text{discrep}(X, B) = \frac{1}{3}$. Thus, $\min\{1, 1 + \text{discrep}(X, B)\} = 1$.

Remark 2.2. Note that [K, Corollary 3.2 (iii)] is wrong by Example 2.1. So, [M, Lemma 2.2] and [K⁺, (4.12.1.3)], which are copies of [K, Corollary 3.2 (iii)], are incorrect. We think that a right formulation is [KM, Proposition 2.36 (2)]. We note that [K, Lemma 3.3] and [M, Lemma 2.3] are correct since ϵ is sufficiently small (see Lemmas 2.3 and 2.5 below). Anyway, the termination theorems proved in [K], [M], and [K⁺, Chapter 4] are true.

It is not difficult to see that [K⁺, (4.12.1.3)] is true if we assume that $b_j \leq \frac{1}{2}$ for all j . We write the precise statement for the reader's convenience. The proof is an exercise.

Lemma 2.3. *Let Y be a smooth variety with a (not necessarily effective) \mathbb{Q} -divisor $B = \sum_i b_i B_i$ such that $\sum_i B_i$ has simple normal crossings, B_i is a prime divisor for every i , $B_k \neq B_l$ for $k \neq l$, and that $b_i \leq \frac{1}{2}$ for all i . Assume that $b_k + b_l \leq 0$ whenever B_k and B_l intersect. If ν is an algebraic valuation with small center on Y such that $a(\nu, Y, B) < 1$ then ν is obtained by blowing up the generic point of a subvariety $W \subset Y$ such that $\text{codim}_Y W = 2$, only one of the B_k (say B_{k_0}) contains W and $b_{k_0} > 0$.*

Remark 2.4. In Example 2.1, we put $D = cL$. Then $a(E_P, X, D) = 1 - c$. Thus, the coefficient of E_P (resp. D' , the strict transform of D) is $c - 1$ (resp. c). Thus, $(c - 1) + c \leq 0$ if and only if $c \leq \frac{1}{2}$. This computation shows that we have to assume $b_j \leq \frac{1}{2}$ for all j in Lemma 2.3.

Thus we obtain the following lemma, which is a correction of [K⁺, (4.12.2.1)]. Note that [K⁺, (4.12.2.2)] is contained in [KM, Proposition 2.36 (2)].

Lemma 2.5. *Let X be a normal variety and $D = \sum_j d_j D_j$ an effective \mathbb{Q} -divisor on X such that $K_X + D$ is \mathbb{Q} -Cartier, where D_j is a prime divisor for every j and $D_k \neq D_l$ for $k \neq l$. Assume that $\text{discrep}(X, D) \geq -\frac{1}{2}$ and $d_j \leq \frac{1}{2}$ for all j . Let ν be an algebraic valuation with small center on X . Then there is a finite set of valuations $\{\nu_i\}$ such that if*

$$a(\nu, X, D) < \min\{1, 1 + \text{discrep}(X, D)\} \text{ and } \nu \notin \{\nu_i\}$$

then ν is obtained from blowing up the generic point of a subvariety $W \subset D \subset X$ such that D and X are generically smooth along W (and thus only one of the D_j contains W) and $\dim W = \dim X - 2$.

Unfortunately, Lemmas 2.3 and 2.5 are useless for our purpose. The assumption that $b_j \leq \frac{1}{2}$ for all j is too strong. Proposition 3.1 below seems to be a better formulation.

§3. New discrepancy lemma

The following proposition is a key result in this paper. The proof is essentially the same as one of [K⁺, (4.12.2.1)]. We give a proof for the reader's convenience.

Proposition 3.1. *Let X be a normal variety and $B = \sum_i b_i B_i$ a \mathbb{Q} -divisor on X with $\lfloor B \rfloor \leq 0$, where B_i is a prime divisor for every i and $B_k \neq B_l$ for $k \neq l$. Assume that $K_X + B$ is \mathbb{Q} -Cartier and $\text{discrep}(X, B) > -1$. Note*

that (X, B) is called a sub klt pair in some literatures. Let ν be an algebraic valuation with small center on X . Then there is a finite set of valuations $\{\nu_i\}$ such that if

$$a(\nu, X, D) < \min\{1, 1 + \text{discrep}(X, D)\} \text{ and } \nu \notin \{\nu_i\}$$

then $V := \text{Center}_X \nu \subset B \subset X$, B and X are generically smooth along V , $\dim V = \dim X - 2$, and only one of the B_k (say B_{k_0}) contains V and $b_{k_0} > 0$.

Proof. First, we take a log resolution $f : Y \rightarrow X$ as in [KM, Proposition 2.36]. Thus, we have $f^*(K_X + B) = K_Y + A - C$, where A and C are both effective divisors with the following properties:

- (i) $A = \sum_{a_i > 0} a_i A_i$ and $C = \sum_{c_j \geq 0} c_j C_j$ have no common irreducible components,
- (ii) $\text{Exc}(f) \cup \text{Supp} f_*^{-1} B = \sum_i A_i \cup \sum_j C_j$, and
- (iii) $\sum_i A_i \cup \sum_j C_j$ is a simple normal crossing divisor and $\sum_i A_i$ is smooth.

Note that c_j may be zero and that $A = f_*^{-1} B + D$, where D is an effective \mathbb{Q} -divisor such that $\text{Supp} D \cap \text{Supp} f_*^{-1} B = \emptyset$. Next, if E is an exceptional divisor over Y such that $a(E, Y, A - C) < 1$, then $V := \text{Center}_Y E \subset A \subset Y$ and $\dim V = \dim Y - 2$ by the following lemma. We note that in general $a(E, Y, F') \leq a(E, Y, F)$ if $F' \geq F$ for any divisor E . If V is contained in D , then $a(E, Y, A - C) \geq 1 + \text{discrep}(X, B)$. Finally, the number of the exceptional divisors over Y whose centers are in $f_*^{-1} B \cap C$ with $a(\cdot, Y, A - C) < 1$ is finite (see Lemma 3.2 below), and it is obvious that the number of f -exceptional divisors is finite. Thus, we obtain the required finite set of valuations $\{\nu_i\}$. \square

Lemma 3.2. *Let Y be a smooth variety and $H = dP$, where P is a smooth prime divisor on Y and $0 < d < 1$. Then $\text{discrep}(Y, H) = 1 - d$. If $a(E, Y, H) < 1$ for an exceptional divisor E over Y , then $\text{Center}_Y E$ is a codimension two subvariety of Y such that $\text{Center}_Y E \subset P \subset Y$.*

Let W be a codimension two subvariety of Y such that $W \subset P \subset Y$. Then there are only finitely many algebraic valuations ν 's with the following properties:

- (1) $a(\nu, Y, H) < 1$,
- (2) $\text{Center}_Y \nu = W$.

Furthermore, ν attains the minimum, that is, $a(\nu, Y, H) = 1 - d$, if and only if ν is obtained by blowing up Y along W .

Proof. This follows from easy computations. See [KM, Lemmas 2.45 and 2.29]. \square

§4. How to define a weighted difficulty

We introduce the notion of *significant* divisors. Proposition 3.1 and Lemma 2.3 imply that the notion of significant divisors are much better than one of *essential* divisors in [F, Definition 2.3] for our purpose.

Definition 4.1. Let (X, B) be a canonical pair. We say that an exceptional divisor E (over X) is *significant* unless $W = \text{Center}_X E$ is a subvariety $W \subset B \subset X$ such that B and X are generically smooth along W (and thus only one of the irreducible components of $\text{Supp} B$ contains W) and $\dim W = \dim X - 2$.

The following corollary is obvious by Proposition 3.1. We will use this to define a weighted version of difficulty.

Corollary 4.2. Let (X, B) be a canonical pair with $\lfloor B \rfloor = 0$. Then we have

$$\#\{E \mid E \text{ is significant and } a(E, X, B) < 1\} < \infty.$$

Remark 4.3. Let (X, B) be a canonical pair. Assume that $\lfloor B \rfloor \neq 0$. Let $f : Y \rightarrow X$ be a log resolution of (X, B) with $f^*(K_X + B) = K_Y + \sum_i a_i E_i$ such that $\sum_i E_i = \text{Exc}(f) \cup \text{Supp} f_*^{-1} B$. We can assume that $a_0 = 1$. If E_0 intersects E_1 such that $0 \leq a(E_1, X, B) = -a_1 < 1$ and $\text{codim}_X f(E_0 \cap E_1) \geq 3$, then we have infinitely many significant divisors whose centers are $f(E_0 \cap E_1)$ with $a(\cdot, X, B) = -a_1$ by blowing up $E_0 \cap E_1$.

We define a weighted version of difficulty. To define this, we have to assume that the boundary divisor has no reduced components.

Definition 4.4 (A weighted version of difficulty). Let (X, B) be a pair with only canonical singularities, where $B = \sum_{j=1}^l b_j B^j$ with $0 < b_1 < \dots < b_l < 1$ and B^j is a reduced divisor for every j . We note that B^j is not necessarily irreducible and that we assume $b_l < 1$. If (X, B) has only terminal singularities,

then $\lfloor B \rfloor = 0$. Thus the assumption $b_l < 1$ always holds for terminal pairs. We put $b_0 = 0$, and $S := \sum_{j \geq 0} b_j \mathbb{Z}_{\geq 0} \subset \mathbb{Q}$. Note that $S = 0$ if $B = 0$. We set

$$d_{S,b}(X, B) := \sum_{\xi \in S, \xi \geq b} \#\{E \mid E \text{ is significant and } a(E, X, B) < 1 - \xi\}.$$

Then $d_{S,b_j}(X, B)$ is finite by Corollary 4.2.

§5. Statements of the termination theorems

Now the proof in [F, §3] works without any changes only if we replace the word "essential" with "significant". Thus we obtain the following theorem, which is slightly weaker than the original theorem: Theorem 1.1 in [F].

Theorem 5.1. *Let X be a normal projective 4-fold and B an effective \mathbb{Q} -divisor such that (X, B) is canonical and $\lfloor B \rfloor = 0$. Consider a sequence of log flips starting from $(X, B) = (X_0, B_0)$:*

$$\begin{array}{ccccccc} (X_0, B_0) & \dashrightarrow & (X_1, B_1) & \dashrightarrow & (X_2, B_2) & \dashrightarrow & \cdots \\ & \searrow & \swarrow & \searrow & \swarrow & & \\ & & Z_0 & & Z_1 & & \end{array},$$

where $\phi_i : X_i \rightarrow Z_i$ is a contraction and $\phi_i^+ : X_i^+ = X_{i+1} \rightarrow Z_i$ is the log flip. Then this sequence terminates after finitely many steps.

As we pointed out before, $\lfloor B \rfloor = 0$ if (X, B) has only terminal singularities. Under the assumption that the varieties are \mathbb{Q} -factorial and all the flipping contractions have the relative Picard number one, we obtain the following theorem by using the special termination theorem. These assumptions are harmless for applications.

Theorem 5.2. *Let X be a normal projective 4-fold and B an effective \mathbb{Q} -divisor such that (X, B) is canonical. Assume that X is \mathbb{Q} -factorial. Consider a sequence of log flips starting from $(X, B) = (X_0, B_0)$:*

$$\begin{array}{ccccccc} (X_0, B_0) & \dashrightarrow & (X_1, B_1) & \dashrightarrow & (X_2, B_2) & \dashrightarrow & \cdots \\ & \searrow & \swarrow & \searrow & \swarrow & & \\ & & Z_0 & & Z_1 & & \end{array},$$

where $\phi_i : X_i \rightarrow Z_i$ is a contraction and $\phi_i^+ : X_i^+ = X_{i+1} \rightarrow Z_i$ is the log flip. We further assume that the relative Picard number $\rho(X_i/Z_i) = 1$ for every i . Then this sequence terminates after finitely many steps.

Proof. By applying the special termination theorem (see [Fj2]) and shifting the index, we can assume that the flipping and flipped loci are disjoint from $\lfloor B_i \rfloor$ for every i . So, we can replace B_i with its fractional part. Thus this sequence terminates by Theorem 5.1. \square

Remark 5.3. The final remark in [F] should be removed. In [Fj], we only need the termination of 4-fold semi-stable terminal flips. See Definition 2.3 in [Fj]. Therefore, Theorems 5.1 and 5.2 are sufficient for [Fj].

References

- [F] O. Fujino, Termination of 4-fold canonical flips, *Publ. Res. Inst. Math. Sci.* **40** (2004), no.1, 231–237.
- [Fj] O. Fujino, On termination of 4-fold semi-stable log flips, to appear in *Publ. Res. Inst. Math. Sci.*
- [Fj2] O. Fujino, Special termination and reduction theorem, to be contained in the book prepared by A. Corti et al.
- [K] J. Kollár, Flops, *Nagoya Math. J.* **113** (1989), 15–36.
- [KM] J. Kollár, and S. Mori, *Birational geometry of algebraic varieties*, Cambridge Tracts in Mathematics, **134**. Cambridge University Press, Cambridge, 1998.
- [K⁺] J. Kollár et al., *Flips and Abundance for algebraic threefolds*, *Astérisque* **211**, (1992).
- [M] K. Matsuki, Termination of flops for 4-folds, *Amer. J. Math.* **113** (1991), no. 5, 835–859.