



CIPMA - UNESCO CHAIRE

GROUP-ALGEBRAIC CHARACTERIZATION OF SPIN PARTICLES: SEMI-SIMPLICITY, $SO(2N)$ STRUCTURE AND IWASAWA DECOMPOSITION

Francis Atta Howard

(Joint work with Mahouton Norbert Hounkonnou and Kinvi Kangni)

International Chair of Mathematical Physics and Applications
(ICMPA-UNESCO Chair)
University of Abomey-Calavi
072 B.P. 50 Cotonou, Republic of Benin

Outline

- 1 Motivation
- 2 Preliminaries
- 3 Spin Semi-simplicity
- 4 Real Lie algebra of Spin particle
- 5 Iwasawa Decomposition of Spin ($\frac{1}{2}$) particle
- 6 Spin Particle Decomposition
- 7 Is spin in Mathematics the same as the spin in particle Physics?
- 8 Concluding remarks

Motivation

3

- [Doran in 1993](#) showed that every linear transformation can be represented as a monomial of vectors in geometric algebra, every Lie algebra as a bivector algebra, and every Lie group as a spin group.
- Schwinger's realization of $su(1,1)$ Lie algebra with creation and annihilation operators was defined with spatial reference in the Pauli matrix representation [[W. Pauli, 1927](#); [J. Schwinger, 1945](#)].
- Several relations as well as connections were observed in spin particles such as fermionic, bosonic, parastatistic Lie algebras, and in geometric algebras such as the Clifford algebra, Grassmannian algebra and so on [[G. Sobezyk, 2015](#)].
- Sobczyk proved that the spin half particles can be represented by geometric algebras. [[G. Sobezyk, 2015](#)]
- [T.D. Palev in 1976](#) highlighted that a semi-simple Lie algebra can be generated by the creation and annihilation operators. In all the above mentioned works, the classical groups such as B_n and D_n play a crucial role in

Motivation

Motivated by all the above mentioned works, a natural questions arise:

- ① Is spin in Mathematics the same as the spin in particle Physics?
- ② If spin (n) is a double cover of $SO(n)$ group, what is the cover of spin (2) and what is also the cover for spin (1/2) are they both related?
- ③ Is it possible to construct the Iwasawa decomposition at both the Lie algebra and Lie group levels of the spin particles ?

But before dealing with the main results, and as a matter of clarity in the development, let us briefly recall the main definitions, the known results, and the appropriate notations useful in the sequel.

Para-fermionic algebra

5

Definition 1 [Green, 1953]

Let a_1^\pm, \dots, a_n^\pm be the creation and annihilation operators for a system consisting of n -fermions with commutator relations :

$$[a_i^-, a_j^+] = \delta_{ij} \quad (3.1)$$

$$[a_i^-, a_j^-] = [a_i^+, a_j^+] = 0, \quad (3.2)$$

or, of n -parafermions with

$$[[a_i^-, a_i^+], a_j^\pm] = \pm 2\delta_{ij} a_j^\pm, \quad (3.3)$$

where

$$[X, Y] := XY - YX. \quad (3.4)$$

Para-fermionic algebra

6

Let T be the associative free algebra of $a_i, a_j; i, j \in N = \{1, 2, \dots, n\}$, and I be the two sided ideal in T generated by the relation (3.3). The Quotient (factor algebra)

$$Q = \frac{T}{I} \quad (3.5)$$

is called para-Fermi algebra, for all $X, Y \in Q$. This is an infinite dimensional Lie algebra with respect to the bracket defined by the equation (3.4).

Semi-simple Lie algebra generated by creation and annihilation operators

7

Definition 2 [Paley, 1976]

Let g be a semi-simple Lie algebra generated by n pairs a_1^\pm, \dots, a_n^\pm of creation and annihilation operators. The elements

$$h_i = \frac{1}{2}[a_i^-, a_i^+], i = 1, \dots, n \quad (3.6)$$

are contained in a Cartan subalgebra H of g . The rank of $g \geq n$. If the semi-simple Lie algebra g of rank n is generated by n pairs of creation and annihilation operators, then, with respect to the basis of the Cartan subalgebra, the creation (resp. annihilation) operators are negative (resp. positive) root vectors. The correspondence with their roots is:

$$a_i^\pm \longleftrightarrow \pm h^* i. \quad (3.7)$$

Lie algebra of spin group

Definitions 3

Let now m be an n -dimensional oriented real vector space with an inner product \langle, \rangle .

- We define the Clifford algebra [La Harpe, 1972] $Cl(m)$ over m by the quotient $T(m)/I$, where $T(m)$ is a tensor algebra over m and I is the ideal generated by all elements $v \otimes v + \langle v, v \rangle 1$, $v \in m$.
- The multiplication of $Cl(m)$ will be denoted by $x \cdot y$.
- Let $p : T(m) \rightarrow Cl(m)$ be the canonical projection. Then, $Cl(m)$ is decomposed into the direct sum $Cl^+(m) \oplus Cl^-(m)$ of the p -images of the elements of even and odd degrees of $T(m)$, and m is identified with the subspace of $Cl(m)$ through the projection p .
- Let e_1, e_2, \dots, e_n be an oriented orthonormal basis of m . The map: $e_{i_2} \cdot e_{i_2} \cdot \dots \cdot e_{i_p} \mapsto (-1)^p e_{i_p} \cdot \dots \cdot e_{i_2} \cdot e_{i_1}$ defines a linear map of $Cl(m)$ and the image of $x \in Cl(m)$ by this linear map is denoted by \bar{x} .

- The spin group is defined by:

$$\text{Spin}(V) = \{x \in \text{Cl}^+(V) : xVx^{-1} \subset V \text{ and } x\bar{x} = 1\}.$$

- Let $\text{Cl}(V) = \text{Cl}_{p,q} = \text{Cl}(\mathbb{R}_{p,q}) = \mathbb{R}_{p,q}$ be the Clifford algebra over \mathbb{R} . Consider $\mathbb{R}_{p,q}^* = \text{Cl}_{p,q}^*$ the group of invertible elements of $\mathbb{R}_{p,q}$. The exponential of $y \in \text{Cl}_{p,q}$ is defined by:

$$\exp(y) = \sum_{n=0}^{\infty} \frac{1}{n!} y^n.$$

- Let $\pi : \text{Spin}(V) \rightarrow \text{SO}(V)$ be defined by $\pi(x)v = xv x^{-1}$. Then, the differential $\dot{\pi}$ of π is given by: $\dot{\pi}(x)v = xv - vx$, for $x \in \mathfrak{spin}(V)$ and $v \in V$.

Spin Lie group and its Lie algebra

10

Suppose now M is an oriented Riemannian manifold.

Definition 4[Milnor; Rodrigues]

A spin structure on M consists of a principal fiber bundle $\pi : P_{\text{Spin}_{p,q}^e}(M) \longrightarrow M$ with a group $P_{\text{Spin}_{p,q}^e}$ and the fundamental map (two fold cover)

$$s : P_{\text{Spin}_{p,q}^e}(M) \longrightarrow P_{\text{SO}_{p,q}^e}(M),$$

satisfying the following conditions:

- (i) $\pi(s(p)) = \pi_s(p)$ for every $p \in P_{\text{Spin}_{p,q}^e}(M)$; π is the projection map of the bundle $P_{\text{SO}_{p,q}^e}(M)$.
- (ii) $s(pu) = s(p)Ad_u$ for every $p \in P_{\text{Spin}_{p,q}^e}(M)$ and

$$Ad : \text{Spin}_{p,q}^e \rightarrow \text{Aut}(\mathcal{Cl}_{p,q}), \quad Ad_u : \mathbb{R}_{p,q} \mapsto uxu^{-1} \in \mathcal{Cl}_{p,q}.$$

the following diagram must commute

$$\begin{array}{ccc}
 P_{\text{Spin}_{p,q}^e}(M) & \xrightarrow{s} & P_{\text{SO}_{p,q}^e}(M) \\
 & \searrow \pi_s & \swarrow \pi \\
 & M &
 \end{array}$$

Definitions 5 [Milnor; Rodrigues]

- A spin manifold is an orientable manifold M together with a spin structure on the tangent bundle of M .
- A spin group is a compact dimensional Lie group.

Spin Lie Group

12

Spin Lie group

The Lie algebra $\mathfrak{spin}(j)$ of spin particles can be represented by classical matrices, which makes it easier to see their algebraic nature:

$$\mathfrak{spin}(j) = \begin{cases} \text{higgs} & j = 0; \\ \text{fermions} & j = \frac{\mathbb{Z}}{2} \text{ when odd integer spins are considered;} \\ \text{bosons} & j = \mathbb{Z} \text{ when positive integer spins are considered.} \end{cases}$$

The Lie algebra $\mathfrak{sl}(2n, \mathbb{C})$ can represent the fermion spin Lie algebra of elementary particles in quantum physics. As indicated in the mapping below. We define $\frac{\mathbb{Z}}{2}$ as fraction of the form $(\frac{2k+1}{2}, k = 0, 1, 2, 3, \dots)$:


$$\mathfrak{sl}(2n, \mathbb{C}) \longrightarrow \mathfrak{spin}\left(\frac{\mathbb{Z}}{2}\right) \longrightarrow \text{fermions.}$$

- The Lie group $SL(2n, \mathbb{C})$ structure can represent the fermion Spin Lie group analogue:

$$SL(2n, \mathbb{C}) \longrightarrow \text{Spin}\left(\frac{\mathbb{Z}}{2}\right) \longrightarrow \text{fermions},$$

while the Lie group $SL(2n + 1, \mathbb{C})$ represents the boson Spin Lie group analogue:

$$SL(2n + 1, \mathbb{C}) \longrightarrow \text{Spin}(\mathbb{Z}) \longrightarrow \text{bosons}.$$



Spin Semi-simplicity

14

Lemma 1

Any spin particle Lie algebra admits a Clifford algebra and a spin group structure.

Proof.

Consider any $\mathfrak{spin}(j)$ with $j = 0, \frac{1}{2}, 1, \dots$, satisfying the spin particle commutator and anticommutator relations (3.1), (3.2), (3.3) as well as the spin Lie algebra commutation bracket rule. It is obvious that the Lie algebra $\mathfrak{spin}(j)$ is a Clifford algebra. Thus, the $\mathfrak{spin}(j)$ exponential is just

$$\exp : \mathfrak{spin}(j) \rightarrow \text{Spin}(J)$$

where $\text{Spin}(J)$ is the spin group. Hence, any spin particle admits a spin group. \square

Lemma 11

Any spin group of a spin particle admits an almost complex spin manifold (Riemannian manifold) and a spin Lie group structure.

Proof.

From Lemma 14, any spin particle admits a spin group. Also, from Definition 10, the spin group, say $\text{Spin}(J)$, has a group structure with an almost complex manifold. Thus, from Definition 3.3, the spin particle, say $\text{Spin}(J)$ with $J = 0, \frac{1}{2}, \dots$, admits a spin manifold. Next, we see that any spin particle has a spin group, say $\text{Spin}(J)$. Since any spin particle has a spin manifold, we observe that $\text{Spin}(J)$ is a spin group and, hence, a spin Lie group. \square

Proposition 1

Any spin half odd integer (resp. integer spin) Lie group is a fourfold cover of the compact Lie group $SO(2n)$ (resp. a double cover of $SO(2n + 1)$).




Theorem 1

Any spin Lie group $\text{Spin}(J)$ of a spin particle is:

- (i) connected;
- (ii) semi-simple if and only if its simple roots are one of the Dynkin's root systems $\Pi(B_n)$ or $\Pi(D_n)$ associated with the classical groups $\text{SO}(2n+1)$ and $\text{SO}(2n)$, respectively.

Proof

We let $\text{Spin}(J)$ be a spin Lie group with $J = 0, \frac{1}{2}, 1, \dots$. For $\text{Spin}(0)$, $\text{Spin}(\frac{1}{2})$, and $\text{Spin}(1)$, we have, respectively, the diagram:



The $sl(2, \mathbb{C})$ Lie algebra can be decomposed into the compact real $su(2)$ and imaginary $isu(2)$ forms, or $sl(2, \mathbb{R})$ and $isl(2, \mathbb{R})$. It is only natural to seek the real form of the spin half particle Lie algebra in terms of Pauli matrices [[Pauli Jr, 1927](#)], which are $sl(2, \mathbb{C})$ matrix basis elements.

Real Lie algebra of Spin particle

20

Proposition II

The real Lie algebra $\mathfrak{spin}(\frac{1}{2})$ of spin half particles ($\text{Spin}(\frac{1}{2})$) is given by $\mathfrak{spin}(\frac{1}{2}) = \{S \in M_2(\mathbb{R}) \mid \text{Tr}S = 0\}$.

- 1 The elements $S_k = \frac{\hbar}{2} \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$, $S_z = \frac{\hbar}{2} \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$, $S_+ = \hbar \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}$ form a basis of the $\mathfrak{spin}(\frac{1}{2})$.
- 2 The commutation relations are given by :
 $[S_k, S_z] = -\hbar S_x$, $[S_k, S_+] = \hbar S_z$, $[S_z, S_+] = \hbar S_+$.

Take an arbitrary angular momentum $\mathfrak{spin}(\frac{1}{2})$ with spinors

$$\chi = \begin{pmatrix} a \\ b \end{pmatrix} = a\chi_{\frac{1}{2}} + b\chi_{-\frac{1}{2}}, \quad \chi_{\frac{1}{2}} = \begin{pmatrix} 1 \\ 0 \end{pmatrix} \quad \text{and} \quad \chi_{-\frac{1}{2}} = \begin{pmatrix} 0 \\ 1 \end{pmatrix}.$$

Let

$$S_x = \frac{S_+ + S_-}{2} \quad \text{and} \quad S_y = \frac{S_+ - S_-}{2i}.$$

From the above equations (??) and (??), we can write S^2 and S_z in terms of spinors. Indeed,

$$S^2 \chi_{\frac{1}{2}} = \hbar^2 \frac{1}{2} \left(\frac{1}{2} + 1 \right) \left| \chi_{\frac{1}{2}} \right\rangle = \frac{3}{4} \hbar^2 \chi_{\frac{1}{2}}, \quad S^2 \chi_{-\frac{1}{2}} = \hbar^2 \frac{3}{4} \chi_{-\frac{1}{2}}. \quad (5.1)$$

From equations (5.1), we can deduce

$$S^2 = \frac{3}{4} \hbar^2 \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} = \frac{3}{4} \hbar^2 I,$$

where $I = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$ is the identity matrix. Similarly,

$$S_z \chi_{\frac{1}{2}} = \frac{\hbar}{2} \chi_{\frac{1}{2}} \quad \text{and} \quad S_z \chi_{-\frac{1}{2}} = -\frac{\hbar}{2} \chi_{-\frac{1}{2}}.$$

Therefore,

$$S_z = \frac{\hbar}{2} \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} = \frac{\hbar}{2} \sigma_z.$$

By analogous computations, we get:

$$S_+ = \hbar \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} = \hbar \sigma_+ \quad (5.2)$$

and

$$S_- = \hbar \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix} = \hbar \sigma_- \quad (5.3)$$

Lemma III

For a $\mathfrak{spin}(\frac{1}{2})$ there exists an orthogonal (skew symmetric) matrix element S_k (with $\hbar = 1$), which can be transformed into an $SO(2)$ compact Lie group. For $g = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in SL(2, \mathbb{R}) = G$, the stabilizer of $i \in \mathbb{C}$ under the action of g is the subgroup $K = SO(2)$.

Remark

The $\mathfrak{spin}(\frac{1}{2}) \in \mathfrak{sl}(2, \mathbb{C})$. Thus, it is complex, and for good notation, we write $\mathfrak{spin}(\frac{1}{2}, \mathbb{C}) \subset \mathfrak{sl}(2, \mathbb{C})$. For the real form, we write $\mathfrak{spin}_{\mathbb{R}}(\frac{1}{2}) \subset \mathfrak{sl}(2, \mathbb{R})$. Finally,

$$\mathfrak{spin}\left(\frac{1}{2}, \mathbb{C}\right) = \mathfrak{spin}_{\mathbb{R}}\left(\frac{1}{2}\right) \oplus i\mathfrak{spin}_{\mathbb{R}}\left(\frac{1}{2}\right).$$

Iwasawa Decomposition of Spin ($\frac{1}{2}$) particle

24

Theorem II

- (i) Let θ, t, ξ be arbitrary real numbers, and put $\hbar k_\theta = \exp(\theta S_k)$, $\hbar d_t^{\frac{1}{2}} = \exp(t S_z)$, and $\hbar n_\xi = \exp(\xi S_+)$. Then, the subgroups $\hbar^3 KDN$ of Spin ($\frac{1}{2}$) are defined by: $\hbar K_\theta = \{\hbar k_\theta | \theta \in R\}$, $\hbar D = \{\hbar d_t^{\frac{1}{2}} | t \in R\}$ and $\hbar N = \{\hbar n_\xi | \xi \in R\}$. We have:

$$\hbar k_\theta = \begin{pmatrix} \hbar \cos \frac{\theta}{2} & \hbar \sin \frac{\theta}{2} \\ -\hbar \sin \frac{\theta}{2} & \hbar \cos \frac{\theta}{2} \end{pmatrix}, \quad \hbar d_t^{\frac{1}{2}} = \begin{pmatrix} \hbar e^{\frac{t}{2}} & 0 \\ 0 & \hbar e^{-\frac{t}{2}} \end{pmatrix},$$

$$\hbar n_\xi = \begin{pmatrix} \hbar & \hbar \xi \\ 0 & \hbar \end{pmatrix},$$

$$\hbar K \cong \frac{\mathbb{R}}{4\pi\mathbb{Z}} \cong T, \quad \hbar D \cong \mathbb{R}, \quad \hbar N \cong \mathbb{R}.$$

- (ii) Any spin ($\frac{1}{2}$) particle is uniquely decomposable in the form:

$$\text{spin} \left(\frac{1}{2} \right) = \hbar^3 k_\theta d_t^{\frac{1}{2}} n_\xi = \exp(\theta \langle {}^s_m | S_k | {}^s_m \rangle) \cdot \exp(t \langle {}^s_m | S_z | {}^s_m \rangle) \cdot \exp(\xi \langle {}^s_m | S_+ | {}^s_m \rangle).$$

Iwasawa decomposition of Lie algebra and Lie group Levels 25

Theorem II: Iwasawa Decomposition of Spin ($\frac{1}{2}$) particle

If $spin(\frac{1}{2}) = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in SL(2, \mathbb{R})$, then, θ, t, ξ in Theorem II (i) are given by the relations:

$$\exp\left(i\frac{\theta}{2}\right) = \frac{a - ic}{\hbar^3 \sqrt{a^2 + c^2}}, \quad (6.2)$$

$$\exp(t) = \frac{a^2 + c^2}{\hbar^6}, \quad (6.3)$$

and

$$\xi = \frac{\hbar^6(ab + cd)}{a^2 + c^2}. \quad (6.4)$$

Now we know that Spin ($\frac{1}{2}$) is spanned by two states: $\{|\frac{1}{2} \frac{1}{2}\rangle, |\frac{1}{2}, -\frac{1}{2}\rangle\}$. From equations (??), (??) and (??), we can calculate the angular momentum for spin half integers such as $\frac{1}{2}$, $\frac{3}{2}$, $\frac{5}{2}$ and so on... [Thankappan, 1972].

A question arises: What can the general term (last term) of a spin half integer be? From a theoretical point of view, this can be useful in the study of particle rotational forms. We have the following results:

Theorem III

For any $\text{spin}(\frac{2n-1}{2})$ (fermions) quantum state, where $n = 1, 2, 3, \dots$, we have:

- (i) $S^2 |S, M\rangle_n = \left(\frac{4n^2-1}{4}\right) \hbar^2 |S, M\rangle.$
- (ii) $S_z |S, M\rangle_n = \pm \left(\frac{2n-k}{2}\right) \hbar |S, M\rangle$ where $k \leq 2n$, and $n = 1, 2, \dots$, with $k = 1, 3, 5, \dots$
- (iii) The n^{th} possible state of a spin half particle is given by:

$$M_{s_n} = 2S_n + 1 = 2n,$$

where $n = 1, 2, 3, \dots$. The quantum state of the fermion is spanned by $2n$ states:

$$\left| \left(\frac{2n-1}{2} \right), \pm \left(\frac{2n-1}{2} \right) \right\rangle, \dots, \left| \left(\frac{2n-1}{2} \right), \pm \left(\frac{2n-k}{2} \right) \right\rangle,$$

(iv) The ladder operators act as follows:

$$S_{+n} \left| \left(\frac{2n-1}{2}, \frac{2n-k}{2} \right) \right\rangle = \hbar \sqrt{(k-1)n - \left(\frac{(k-1)(k-1)}{4} \right)} \left| S, M+1 \right\rangle, \quad (6.5)$$

$$S_{+n} \left| \left(\frac{2n-1}{2}, -\frac{2n-k}{2} \right) \right\rangle = \hbar \sqrt{(k+1)n - \left(\frac{(k+1)(k+1)}{4} \right)} \left| S, M+1 \right\rangle, \quad (6.6)$$

Note

$$S_{y_n} = \frac{S_{+n} - S_{-n}}{2i} = \frac{\hbar \sigma_{k_n}}{2i} = -i S_{k_n}. \quad (6.7)$$

Theorem IV

For any $\mathfrak{spin}(\frac{2n-1}{2})$, the quantum state of the particle is spanned by $2n$ states and there exists orthogonal matrix element S_{k_n} in the S_{y_n} matrix which can be transformed into the classical group $SO(2n)$ with natural numbers $n = 1, 2, 3, \dots$. This compact Lie group $SO(2n)$ corresponds to the Dynkin's root $\Pi(D_n)$.

Proof.

From Lemma III, From Lemma 23 we observe that the theorem is true for $n = 1$. For $\mathfrak{spin}(\frac{2n-1}{2})$ particle quantum state spanned by $2n$ states, we consider similar arguments for Theorem II, replacing the S_k matrix by the n^{th} matrix S_{k_n} and deducing in the same manner as in Lemma III to obtain the above Theorem III. Specifically, from Theorem II, there exists S_{k_n} matrix in the S_{y_n} matrix from equation 6.10. One can check that these matrices are orthogonal and generate $SO(2n)$ with $n = 1, 2, 3, \dots$. For $n = 1$ we have the compact Lie group $SO(2)$ as in Lemma III. Finally, the correspondence to the Dynkin's roots $\Pi(D_n)$.

Proposition III [Sugiura, 1977]

For an element $g \in \text{Spin}_{\mathbb{R}}\left(\frac{1}{2}\right)$ and $\theta \in \mathbb{R}$, let

$$gk_{\theta} = k_{g \cdot \theta} d_{t(g, \theta)}^{\frac{1}{2}} n_{\xi(g, \theta)}$$

be the Iwasawa decomposition of gk_{θ} . If $\hbar = 1$, then, the following cocycle conditions hold, for g, g' which are projections in $\text{SL}(2, \mathbb{R}) \supset \text{Spin}_{\mathbb{R}}\left(\frac{1}{2}\right)$:

- ❶ (i) $(gg') \cdot \theta \equiv g \cdot (g' \cdot \theta) \pmod{4\pi}$;
- (ii) $t(gg', \theta) = t(g, g' \cdot \theta) + t(g', \theta)$;
- (iii) $g \cdot (\theta + 2\pi) = g \cdot \theta + 2\pi \pmod{4\pi}$, $t(g, \theta + 2\pi) = t(g, \theta)$.

Proposition III [Sugiura, 1977]

1 If $g = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$, then, we have:

- (i) $\exp\left(\frac{i(g \cdot \theta)}{2}\right) = \frac{(a - ic) \cos \frac{\theta}{2} + (-b + id) \sin \frac{\theta}{2}}{|(a - ic) \cos \frac{\theta}{2} + (-b + id) \sin \frac{\theta}{2}|}$;
- (ii) $\exp(t(g, \theta)) = |(a - ic) \cos \frac{\theta}{2} + (-b + id) \sin \frac{\theta}{2}|^2$;
- (iii) $\xi(g, \theta) = \exp(-t(g, \theta)) \times [(ab + cd) \cos \theta + \frac{1}{2}(a^2 - b^2 + c^2 - d^2) \sin \theta]$;
- (iv) $d \frac{d(g \cdot \theta)}{d\theta} = \exp(-t(g, \theta))$.

Theorem V[Main Result: Particle Decomposition]

Any spin Lie group G can be uniquely decomposed in the form:

$$G = \mathbb{K}KD^sN$$

where K is compact, D^s is a rotational function (d -function), and N is nilpotent (Ladder operators). We denote by $\mathbb{K}(\alpha^{-1})$ the fine structure constant and all other translational energy of elementary spin particles.

Fine structure constant

$$\alpha^2 = \frac{(k_c)(q)^2}{(h/\pi)(c)}, \quad \alpha^{-1} = 137.036\dots$$

$$\alpha = \frac{1}{137} = \mathbb{K}.$$

Is Spin(n) in Mathematics the same as the Spin(J) in particle Physics?


33


In mathematics the spin group $spin(n)$ is a double cover of the special orthogonal group $SO(n)$.

$$Spin(3)$$



$$SO(3)$$

however, in particle Physics





while in Physics;



Concluding remarks

35

Concluding remarks

In this work,

- we provided an extension of semi-simplicity of spin particle Lie algebra to the Lie group level,
- we showed that a spin particle Lie algebra admits a Clifford algebra, an almost complex manifold and a spin Lie group structure,
- we demonstrated that any spin half particle, (resp. integer spin), spin Lie group is a fourfold, (resp. double), cover of the $SO(2n)$, (resp. $SO(2n + 1)$), we also proved that any spin Lie group of a spin particle is connected and semi-simple,
- we constructed the real Lie algebra of the Spin ($\frac{1}{2}$) particle,
- we also performed the Iwasawa decomposition of the spin half into KDN ,
- finally, we applied the angular momentum coupling to the Spin ($\frac{2n-1}{2}$) particle and demonstrated that the orthogonal basis transforms into the

Motivation

Preliminaries

Spin Semi-simplicity

Real Lie algebra of Spin particle

Iwasawa Decomposition of Spin ($\frac{1}{2}$) particle

Spin Particle Decomposition

Is spin in Mathematics the same as the spin in particle Physics?

Concluding remarks

QUESTIONS?

THANKS FOR YOUR ATTENTION!