

Mapping class group, Donaldson-Thomas theory and S-duality

Kentaro Nagao

November 9, 2011

Abstract

We study Donaldson-Thomas theory associated to a triangulated surface. We show that the generating function of the Donaldson-Thomas invariants is “invariant” under an action of the mapping class group, which is identified with the mapping class group action in the (decorated) Teichmüller theory. This gives an example of constraints of the generating function induced by the derived auto-equivalences. From the view point of string theory, this is nothing but S-duality of the BPS spectrum of the 4d gauge theory given by Gaiotto-type construction.

Introduction

The DT invariant for a Calabi-Yau 3-fold Y is a counting invariant of coherent sheaves on Y , which is introduced in [Tho00] as a holomorphic analogue of the Casson invariant on a real 3-manifold. Although the category of coherent sheaves on Y is an Abelian category, it has been known that we take it as a counting invariant of objects in the derived category.

An ideal application of this formulation might be the following: The derived category sometimes have a non-trivial auto-equivalence group. In such a case, the generating function might have a good transformation formula with respect to this action, which would help us to determine the generating function.

In this notes, we will show a new example¹ of such a phenomenon. We study Donaldson-Thomas theory associated to a triangulated surface. The mapping class group acts on the derived category and the generating function of the Donaldson-Thomas invariants is “invariant” under this action.

¹As far as the author understand, it is the first example.

Plan

In §1, we briefly review the construction in [FST08, LF09] of quivers with potential associated to triangulated surfaces. In §2, we study the mapping class group actions on the derived category and the associated Poisson torus. The later is identified with the mapping class group action on the decorated Teichmüller space as is shown in §5. The main result of this paper appears in §3. Finally, we explain an interpretation of the main result in terms of S -duality ([Gai]) in §4.

Acknowledgement

The author would like to thank Daniel Labardini-Fragoso for answering his questions on [LF09], Takahiro Nishinaka and Masahito Yamazaki for exciting discussions. The author is deeply grateful to Yuji Tachikawa for his elegant and mathematician-friendly talks on Gaiotto functor in New developments in Infinite analysis 2011. The main result in this paper was first announced in the same conference in 2010. The author would like to thank Akihiro Tsuchiya and Yuji Yamada for organizing these conferences.

The main result of this paper was obtained more than a year ago ([Nagb]), but the author did not submit it since it is just a direct corollary of the formulae in [KS] and [Naga]. Recently, the author found that the result may have a good chance to be developed and so decided to enforce the manuscript and submit it.

1 QP for a triangulated surface

In this section, we briefly explain how to associate a quiver with a potential for a triangulated surface [FST08, LF09].

1.1 Ideal triangulations of a surface

Let Σ be a compact connected oriented surface with (possibly non-empty) boundary and M be a finite set of points on Σ , called marked points. We assume that M is non-empty and has at least one point on each connected component of the boundary of Σ . The marked points that lie in the interior of Σ will be called **punctures**, and the set of punctures of (Σ, M) will be denoted P .²

²We will always assume that (Σ, M) is none of the following:

- a sphere with less than five punctures;

We decompose Σ into “triangles” (in the topological sense) so that each edge is either

- a curve (which is called an *arc*) whose endpoints are in M or
- a connected component of $\partial\Sigma \setminus M$.

A triangle may contain exactly two arcs (see Figure 1). Such a triangle (and its doubled arc) is said to be *self-folded*.




Figure 1: A self-folded triangle

Given a triangulation τ and a (non self-folded) arc i , we can *flip* i to get a new triangulation $f_i(\tau)$ (see Figure 2).




Figure 2: A flip of a triangulation

Theorem 1.1 ([FST08]). *Any two triangulations are related by a sequence of flips.*

1.2 Quiver for a triangulation

Let τ be a triangulation. We will define a quiver $Q(\tau)$ without loops and 2-cycles whose vertex set I is the set of arcs in τ .

-
- an unpunctured monogon, digon or triangle;
 - a once-punctured monogon.

Here, by a monogon (resp. digon, triangle) we mean a disk with exactly one (resp. two, three) marked point(s) on the boundary.

For a (non self-folded³) triangle Δ and arcs i and j , we define a skew-symmetric integer matrix B^Δ by

$$B_{i,j}^\Delta := \begin{cases} 1 & \Delta \text{ has sides } i \text{ and } j, \text{ with } i \text{ following } j \text{ in the clockwise order,} \\ -1 & \text{the same holds, but in the counter-clockwise order,} \\ 0 & \text{otherwise.} \end{cases}$$

We put

$$B(\tau) := \sum_{\Delta} B^\Delta$$

where the sum is taken over all triangles in τ . Let $Q(\tau)$ denote the quiver without loops and 2-cycles associated to the matrix $B(\tau)$.

Theorem 1.2 ([FST08]). *Given a triangulation τ and its (non self-folded) arc i , we have*

$$Q(f_i(r)) = \mu_i(Q(\tau))$$

where μ_i denote the mutation of the quiver at the vertex i .

1.3 Potential for a triangulation

For a triangle Δ in τ , we define a potential ω_Δ as in Figure 3. For a puncture




Figure 3: ω_Δ

P in τ , we define a potential ω_P as in Figure 4.

Finally, we put

$$\omega(\tau) := \sum_{\Sigma} \omega_\Sigma + \sum_P \omega_P.$$

Theorem 1.3 ([LF09]). *Given a triangulation τ and its (non self-folded) arc i , we have*

$$\omega(f_i(r)) = \mu_i(\omega(\tau))$$

where μ_i denote the mutation of the potential at the vertex i in the sense of [DWZ08].

³We omit the definition of B^Δ for a self-folded triangle Δ .




Figure 4: ω_P

Remark 1.4. *More strongly, the QP $(Q(\tau), \omega(\tau))$ is f-mutable (see [Naga, Definition 1.2.(2)]). This is important for us since otherwise we can not apply the results in [Naga] for a technical reason. See the comments at the end of the introduction of [Naga].*

2 Mapping class group action

2.1 Mapping class group

We define

$$\text{Diffeo}(\Sigma, M) := \{\phi: \Sigma \rightarrow \Sigma \mid \phi: \text{diffeomorphism}, \phi(M) = M\}.$$

Let $\text{Diffeo}(\Sigma, M)_0$ denote the connected component of $\text{Diffeo}(\Sigma, M)$ which contains id_Σ . The quotient

$$\text{MCG}(\Sigma, M) := \text{Diffeo}(\Sigma, M) / \text{Diffeo}(\Sigma, M)_0$$

is called the *mapping class group*.

2.2 Derived category for a triangulation

Let $\Gamma(\tau)$ be Ginzburg's dg algebra associated to the quiver with the potential $(Q(\tau), \omega(\tau))$ and $\mathcal{D}(\tau) = \mathcal{D}\Gamma(\tau)$ be the derived category of right dg-modules over Γ . By the result of Keller ([Kel]), $\Gamma(\tau)$ and $\Gamma(f_i(\tau))$ are equivalent⁴.

For a triangulation τ and an element $\phi \in \text{MCG}(\Sigma, M)$, we get another triangulation $\phi(\tau)$. Note that $(Q(\tau), \omega(\tau))$ and $(Q(\phi(\tau)), \omega(\phi(\tau)))$ (and hence $\mathcal{D}(\tau)$ and $\mathcal{D}(\phi(\tau))$) are canonically identified.

By Theorem 1.1, τ and $\phi(\tau)$ are related by a sequence of flips. Each flip gives a derived equivalence. By composing the derived equivalences, we get

⁴Since we have two derived equivalences, we have to choose one of them. Given a sequence of flips, we have a canonical choice. See [Naga, §2.2].

a derived equivalence

$$\Psi_\phi: \mathcal{D}(\tau) \xrightarrow{\sim} \mathcal{D}(\phi(\tau)) = \mathcal{D}(\tau).$$

Thanks to the result [FST08, Theorem 3.10] and the pentagonal identity for the derived equivalences, Ψ_ϕ is independent of choices of a sequence of flips and well-defined. Finally we get an action of the mapping class group on the derived category:

$$\Psi: \text{MCG}(\Sigma, M) \rightarrow \text{Aut}(\mathcal{D}(\tau)).$$

2.3 Cluster transformation

We put $T = T(\tau) := \mathbb{C}[x_i, x_i^{-1}]_{i \in I}$. We define $\text{CT}_k: T(f_k(\tau)) \xrightarrow{\sim} T(\tau)$ by

$$\text{CT}_k(x'_i) = \begin{cases} (x_k)^{-1} \left(\prod (x_j)^{Q(j,k)} + \prod (x_j)^{Q(k,j)} \right) & i = k, \\ x_i & i \neq k \end{cases}$$

where $Q(i, k)$ is the number of arrows from i to k and x'_i is the generator of $T(f_k(\tau))$.

In the same way as the previous section, we get

$$\text{CT}_\phi: T(\phi(\tau)) \xrightarrow{\sim} T(\tau).$$

Under the identification $T(\phi(\tau)) = T(\tau)$ induced by Ψ_ϕ , we get

$$\text{CT}: \text{MCG}(\Sigma, M) \rightarrow \text{Aut}(T(\tau)).$$

Remark 2.1. *As we will explain in §5, this is compatible with the action of mapping class group on the decorated Teicummuller space.*

3 Donaldson-Thomas theory

Let J_τ be the Jacobi algebra associated to the quiver with the potential $(Q(\tau), W(\tau))$.

Let P_τ^i be the indecomposable projective J_τ -module associated to $i \in I$. For $\mathbf{v} \in \mathbb{Z}_{\geq 0}^I$, we define

$$\text{Hilb}_\tau^i(\mathbf{v}) := \{P_\tau^i \twoheadrightarrow V \mid \underline{\dim} V = \mathbf{v}\}.$$

This is called the *Hilbert scheme*⁵.

⁵The name comes from the Hilbert scheme in algebraic geometry which parameterizes quotient sheaves of the structure sheaf.

Definition 3.1. We define $\text{DT}_\tau: T \xrightarrow{\sim} T$ by

$$\text{DT}_\tau(x_i) := (x_i)^{-1} \cdot \sum_{\mathbf{v}} \text{Eu}(\text{Hilb}_\tau^i(\mathbf{v})) \cdot y^{-\mathbf{v}}$$

where

$$y^{-\mathbf{v}} := \prod_i (y_i)^{-v_i}, \quad y_i := \prod_j (x_i)^{Q(i,j)}.$$

As a direct application of the main theorem in [Naga], we get the following:

Theorem 3.2. For any element $\phi \in \text{MCG}(\Sigma, M)$, we have

$$\text{DT}_\tau \circ \text{CT}_\phi = \text{CT}_\phi \circ \text{DT}_\tau.$$

4 S-duality interpretation

4.1 Gaiotto functor

Let \mathcal{F} is an n -dimensional quantum field theory. Then for any fixed k -dimensional manifold K , the correspondence

$$M \longmapsto \mathcal{F}(K \times M)$$

provides an $(n - k)$ -dimensional quantum field theory.

We take a 6d $\mathcal{N} = (2, 0)$ quantum field theory S_G , where G is compact Lie group of type ADE . Fixing a Riemann surface C , we get a 4d $\mathcal{N} = 2$ theory by the construction above. Let $S_{G,C}$ denote this theory ([Gai]).

In summary, 6d $\mathcal{N} = (2, 0)$ theory provides the following correspondence :

$$\begin{array}{ccc} \{\text{Riemann surfaces}\} & \longrightarrow & \{4\text{d } \mathcal{N} = 2 \text{ QFT}\} \\ C & \longmapsto & S_{G,C}. \end{array}$$

Following Y. Tachikawa, we call this ‘‘Gaiotto functor’’⁶ (see [MT]).

4.2 4d BPS spectrum

In this paper, we have studied the DT theory associated to a triangulation of a surface C . It is expected that the generating function provides the BPS spectrum of the 4d QFT $S_{\text{SU}(2),C}$ ([GMN, ACC⁺b, ACC⁺a]).⁷

⁶This is not a functor of categories in mathematical sense.

⁷For $S_{\text{SU}(N),C}$, we need to take the triangulations which appear in the higher Teichmüller space ([FG06]).

Remark 4.1. *For BPS spectrum of the 4d QFT, there should be a wall-crossing theory which is compatible with those of DT theory under the expectation above [GMN, Moo].*

4.3 S-duality

Fixing a topological type of a 2-dimensional manifold, the Teichmüller space is the space of complex structures on it. The mapping class group acts on the Teichmüller space so that the quotient space gives the moduli space of complex structures.

Under Gaiotto functor, Teichmüller space should give the space of parameters⁸ of 4-dimensional quantum field theories. Since two points on a mapping class group orbit in the Teichmüller space give a common complex structure, they provide a common 4d theory. This is the S-duality in the sense of Gaiotto ([Gai]).

Remark 4.2. *The original S-duality is the duality between strongly/weakly coupled regions in the space of parameters. Strongly or weakly coupled regions appear as neighborhoods of cusps in the fundamental region.*

Combining the observations in §4.2, we get an interpretation of Theorem 3.2 as the S-duality on the BPS spectrum.

Remark 4.3. *In this paper, we understand the mapping class group action in DT theory via wall-crossing. We can understand the S-duality as a consequence of wall-crossing of 4d QFT, without passing through DT theory (Figure 5).*




Figure 5: Summary

⁸parameter space of exactly marginal gauge couplings.

5 Appendix : Teichmüller theory

Let $\mathcal{T}(\Sigma)$ denote the Teichmüller space and $\tilde{\mathcal{T}}(\Sigma)$ denote the decorated Teichmüller space, which is a $(\mathbb{R}_{>0})^s$ -bundle over $\mathcal{T}(\Sigma)$ whose fiber is the set of s -tuples of horocycles around each of the marked points ([Pen87, Pen04]).

We assume that a triangulation τ does not contain self-folded arcs. Let τ_1 be the set of edges of a triangulation τ . Each edge e in τ_1 , we take the (unique) geodesic represents e . The coordinate $l_e(P)$ is defined as the hyperbolic length of the segment of the geodesic that lies between the two horocycles surrounding the punctures connected by e , taken with positive sign if the two horocycles are disjoint, with negative sign otherwise.

Theorem 5.1. [Pen87, Pen04]

(1) For an ideal triangulation τ without self-folded arcs, the function

$$\vec{l} : \tilde{\mathcal{T}}(\Sigma) \rightarrow \mathbb{R}^{\tau_1}, \quad P \mapsto (l_e(P))_{e \in \tau_1}$$

is a homeomorphism. (This is called the Penner coordinate of the decorated Teichmüller space.)

(2) We put

$$\lambda_e := \sqrt{2} \exp(l_e/2)$$

which is called the Lambda length of e . Let τ' be the triangulation obtained by flipping the edge e . The coordinates associated to τ and τ' agree for each edge which the two triangulations have in common, and

$$\lambda_{e'} = \frac{\lambda_a \lambda_c + \lambda_b \lambda_d}{\lambda_e}.$$

We define the inclusion

$$\begin{array}{ccccc} \tilde{\mathcal{T}}(\Sigma) & \simeq & \mathbb{R}^{\tau_1} & \hookrightarrow & (\mathbb{C}^*)^{\tau_1} & \simeq & \text{Spec}(T(\tau)) \\ (l_e) & \mapsto & (x_e) & = & (\lambda_e) & & \end{array}.$$

We call $T(\tau)$ as the complexified decorated Teichmüller space. The mapping class group action on $\text{Spec}(T(\tau))$ given in §2.3 preserves $\tilde{\mathcal{T}}(\Sigma)$. If we can realize all the mapping classes by a sequence of flips without self-folded arcs then restricted action coincides with the geometric one.

Remark 5.2. In [NTY], we study hyperbolic structures on the mapping torus of a pseudo-Anosov mapping class g of a surface. We show that a fixed point on $T(\tau)$ with respect to the action of g gives a hyperbolic structures on the mapping torus, while the fixed point set on $\tilde{\mathcal{T}}(\Sigma)$ is empty due to the Nielsen-Thurston classification.

References

- [ACC⁺a] M. Alim, S. Cecotti, C. Cordova, S. Espahbodi, A. Rastogi, and C. Vafa, *BPS Quivers and Spectra of Complete $\mathcal{N} = 2$* , arXiv:1109.4941.
- [ACC⁺b] ———, *BPS Quivers Theories*, to appear.
- [DWZ08] H. Derksen, J. Weyman, and A. Zelevinsky, *Quivers with potentials and their representations I: Mutations*, Selecta Math. **14** (2008).
- [FG06] V. Fock and A. Goncharov, *Moduli spaces of local systems and higher Teichmüller theory*, Publ. Math. IHES **103** (2006), no. 1.
- [FST08] S. Fomin, M. Shapiro, and D. Thurston, *Cluster algebras and triangulated surfaces. part i: Cluster complexes*, Acta Math. **201** (2008).
- [Gai] D. Gaiotto, *$\mathcal{N} = 2$ dualities*, arXiv:0904.2715.
- [GMN] D. Gaiotto, G. W. Moore, and A. Neitzke, *Wall-crossing, Hitchin systems, and the WKB approximation*, arXiv:0907.3987.
- [Kel] B. Keller, *Deformed Calabi-Yau Completions*, arXiv:0908.3499v5.
- [KS] M. Kontsevich and Y. Soibelman, *Stability structures, motivic Donaldson-Thomas invariants and cluster transformations*, arXiv:0811.2435v1.
- [LF09] D. Labardini-Fragoso, *Quivers with potentials associated to triangulated surfaces*, Proc. London Math. Soc. **98** (2009).
- [Moo] G. Moore, *PiTP lectures on BPS states and wall-crossing in $d = 4$, $\mathcal{N} = 2$ theories*, http://www.physics.rutgers.edu/gmoore/PiTP_July26_2010.pdf.
- [MT] G. Moore and Y. Tachikawa, *On 2d TQFTs whose values are holomorphic symplectic varieties*, arXiv:1106.5698, to appear in the proceedings of String-Math 2011 conference.
- [Naga] K. Nagao, *Donaldson-Thomas theory and cluster algebras*, arXiv:1002.4884.
- [Nagb] K. Nagao, *Triangulated surface, mapping class group and Donaldson-Thomas theory*, Proceedings of Kinosaki algebraic geometry symposium 2010.

- [NTY] K. Nagao, Y. Terashima, and M. Yamazaki, *Pseudo-asonov mapping tori and cluster algebras*, to appear.
- [Pen87] R. C. Penner, *The decorated Teichmüller space of punctured surfaces*, Comm. Math. Phys. **113** (1987).
- [Pen04] ———, *Decorated teichmauller theory of bordered surfaces*, Comm. Anal. Geom. **12** (2004).
- [Tho00] R. P. Thomas, *A holomorphic Casson invariant for Calabi-Yau 3-folds, and bundles on K3 fibrations*, J. Differential Geom. **54** (2000), no. 2, 367–438.

Kentaro Nagao
 Graduate School of Mathematics, Nagoya University
 kentaron@math.nagoya-u.ac.jp