

A variant of the induction theorem for Springer representations

Toshiaki Shoji*

Graduate School of Mathematics
Nagoya University
Chikusa-ku, Nagoya 464-8602, Japan

ABSTRACT. Let G be a simple algebraic group over \mathbf{C} with the Weyl group W . For a unipotent element $u \in G$, let \mathcal{B}_u be the variety of Borel subgroups of G containing u . Let L be a Levi subgroup of a parabolic subgroup of G with the Weyl subgroup W_L of W . Assume that $u \in L$ and let \mathcal{B}_u^L be a similar variety as \mathcal{B}_u for L . For a certain choice of L , $u \in L$ and $e \geq 1$, we describe the W -modules $\bigoplus_{n \equiv k \pmod{e}} H^{2n}(\mathcal{B}_u)$ for $k = 0, \dots, e-1$, in terms of the W_L -module $H^*(\mathcal{B}_u^L)$ with some additional data, which is a refinement of the induction theorem due to Lusztig. As an application, we give an explicit formula for the values of Green functions at root of unity, in the case where u is a regular unipotent element in L .

0. INTRODUCTION

Let G be a connected reductive group over an algebraically closed field k , and W the Weyl group of G . For a unipotent element $u \in G$, let \mathcal{B}_u be the variety of Borel subgroups containing u . According to Springer [Sp2], Lusztig [L1], W acts naturally on the l -adic cohomology group $H^n(\mathcal{B}_u) = H^n(\mathcal{B}_u, \bar{\mathbf{Q}}_l)$, the so-called Springer representations of W . Assume that $k = \mathbf{C}$, or the characteristic p of k is good. Then it is known that $H^{\text{odd}}(\mathcal{B}_u) = 0$. We consider the graded W -module $H^*(\mathcal{B}_u) = \bigoplus_{n \geq 0} H^{2n}(\mathcal{B}_u)$. Let L be a Levi subgroup of a parabolic subgroup of G . Let W_L be the Weyl group of L , which is naturally a subgroup of W . If $u \in L$, the variety \mathcal{B}_u^L is defined by replacing G by L , and we have a graded W_L -module $H^*(\mathcal{B}_u^L)$.

Lusztig proved in [L3] an induction theorem for Springer representations, which describes the W -module structure of $H^*(\mathcal{B}_u)$ in terms of the W_L -module structure of $H^*(\mathcal{B}_u^L)$, in the case where $u \in L$. However in this theorem, the information on the graded W -module structure is eliminated. In this paper, we try to recover partly the graded W -module structure, i.e., for a fixed positive integer e , we consider the W -modules $V_{e,k} = \bigoplus_{n \equiv k \pmod{e}} H^{2n}(\mathcal{B}_u)$ for $k = 0, \dots, e-1$. Let G be a simple group modulo center defined over \mathbf{C} . We show, under a certain choice of L , u and e , that the W -module $V_{e,k}$ can be described in terms of the graded W_L -module $H^*(\mathcal{B}_u^L)$.

*A part of this work was done during the author's stay in CIB in Lausanne in June 2005. The author is grateful for their hospitality.

with some additional data. In particular, we see that $\dim V_{e,k}$ is independent of the choice of k .

In the case where $u = 1$, $H^*(\mathcal{B}_u)$ is isomorphic, as a graded W -module, to the coinvariant algebra of W . In this case $V_{e,k}$ has been studied by many authors, by Stembridge [St] for e corresponding to the regular elements in W , by Morita and Nakajima [MN1] for $W = \mathfrak{S}_n$ with e such that $1 \leq e \leq n$, and by Bonnafé, Lehrer and Michel [BLM] for complex reflection groups W in the most general framework. Our result partly covers the result of [BLM]. For general $u \neq 1$, Morita and Nakajima [MN2] considered certain types of unipotent elements for $G = GL_n$, which is a special case of ours.

The proof of the induction theorem in [L3] is done by passing to the finite field \mathbf{F}_q , and using a certain specialization argument $q \mapsto 1$ together with the properties of Deligne-Lusztig's virtual character $R_T(1)$. Our argument is a variant of that in [L3]. We use a specialization $q \mapsto \zeta$, where ζ is a primitive e -th root of unity. Thus our argument is closely related to the values of Green functions at root of unity. In the case where u is a regular unipotent element in L , we obtain an explicit formula for such values, which is regarded as a generalization of the result by Lascoux, Leclerc and Thibon [LLT] for the case of Green polynomials of GL_n .

1. THE STATEMENT OF THE MAIN RESULT

1.1. Let k be an algebraic closure of a finite field with $\text{ch}(k) = p > 0$ or the complex number field \mathbf{C} . Let G be a connected reductive group G over k . Let \mathcal{B} be the variety of Borel subgroups of G , and W the Weyl group of G . For any $g \in G$, put $\mathcal{B}_g = \{B' \in \mathcal{B} \mid g \in B'\}$. We consider the Springer representations of W on $H^n(\mathcal{B}_g, \bar{\mathbf{Q}}_l)$ (or on $H^n(\mathcal{B}_g, \mathbf{C})$ in the case where $k = \mathbf{C}$).

Let L be a Levi subgroup of a parabolic subgroup P of G . The Weyl group W_L of L is naturally identified with a subgroup of W . Let \mathcal{B}^L be the variety of Borel subgroups of L . For a unipotent element $u \in L$, we consider $\mathcal{B}_u^L = \{B' \in \mathcal{B}^L \mid u \in B'\}$. Thus we have a W_L -module $H^n(\mathcal{B}_u^L, \bar{\mathbf{Q}}_l)$, and a W -module $H^n(\mathcal{B}_u, \bar{\mathbf{Q}}_l)$. The induction theorem for Springer representations asserts that

$$(1.1.1) \quad \sum_{n \geq 0} (-1)^n H^n(\mathcal{B}_u, \bar{\mathbf{Q}}_l) = \text{Ind}_{W_L}^W \left(\sum_{n \geq 0} (-1)^n H^n(\mathcal{B}_u^L, \bar{\mathbf{Q}}_l) \right)$$

as virtual W -modules.

Remark 1.2. The induction theorem was stated in [AL], with a brief indication of the proof, in the case where $k = \mathbf{C}$, and was proved in [L3] for any k . Note that if p is good, the unipotent classes in G are parametrized in the same way as the case of $k = \mathbf{C}$, independent of p . Moreover in that case, it is known that $H^n(\mathcal{B}_u, \bar{\mathbf{Q}}_l) = 0$ for odd n . Then the algorithm of computing Green functions implies that the W -module structure of $H^n(\mathcal{B}_u, \bar{\mathbf{Q}}_l)$ is independent of p . Thus by a general principle $H^n(\mathcal{B}_u, \bar{\mathbf{Q}}_l)$ is isomorphic to the W -module $H^n(\mathcal{B}_{u'}, \mathbf{C})$, where $u', \mathcal{B}_{u'}$ are the corresponding objects in the algebraic group $G_{\mathbf{C}}$ over \mathbf{C} . In what follows, we express $H^n(\mathcal{B}_u, \bar{\mathbf{Q}}_l)$ or $H^n(\mathcal{B}_{u'}, \mathbf{C})$ by $H^n(\mathcal{B}_u)$ by abbreviation.

1.3. Assume that $k = \mathbf{C}$. We consider the following variant of the induction theorem. Let Γ be a cyclic group of order e generated by a . Let ζ be a primitive e -th root of unity in \mathbf{C} . Let $V = \bigoplus_{n \geq 0} V_n$ be a graded W -module. Then V turns out to be a $\Gamma \times W$ -module by defining the action of Γ on V by $ax = \zeta^n x$ for $x \in V_n$. We denote by $V^{(\zeta)}$ the thus obtained $\Gamma \times W$ -module V .

For $u \in L$, we consider the graded W_L -module $H^*(\mathcal{B}_u^L) = \bigoplus_{n \geq 0} H^{2n}(\mathcal{B}_u^L)$, where the degree n part is given by $H^{2n}(\mathcal{B}_u^L)$, and similarly we consider the graded W -module $H^*(\mathcal{B}_u) = \bigoplus_{n \geq 0} H^{2n}(\mathcal{B}_u)$. Let Γ be as before. We choose Γ such that $\Gamma \subset N_W(W_L)$, and consider the semidirect product $\widetilde{W}_L = \Gamma \ltimes W_L$. We assume that the W_L -module $H^n(\mathcal{B}_u^L)$ can be extended to a \widetilde{W}_L -module for each n . (In the case where $a \in Z_W(W_L)$, we have $\widetilde{W}_L = \Gamma \times W_L$. In this case, one can choose a trivial extension to \widetilde{W}_L , i.e., we may assume that $\Gamma \subset \widetilde{W}_L$ acts trivially on $H^*(\mathcal{B}_u^L)$.) Then one can define a $\Gamma \times \widetilde{W}_L$ -module $H^*(\mathcal{B}_u^L)$ as above, replacing W_L by \widetilde{W}_L , which we denote by $H^*(\mathcal{B}_u^L)^{(\zeta)}$. (When we need to distinguish the group Γ as the first factor of $\Gamma \times \widetilde{W}_L$ from the subgroup of \widetilde{W}_L , we write the latter as Γ_0 .) $\Gamma \times W$ -module $H^*(\mathcal{B}_u)^{(\zeta)}$ is defined as before. Put $V^{(\zeta)} = H^*(\mathcal{B}_u^L)^{(\zeta)}$, and let $V_n^{(\zeta)}$ be the degree n -part of $V^{(\zeta)}$. Let us consider the induced W -module

$$\text{Ind}_{W_L}^W V^{(\zeta)} = \bigoplus_{w \in W/W_L} w \otimes V^{(\zeta)}.$$

Then $\text{Ind}_{W_L}^W V^{(\zeta)}$ turns out to be a $\Gamma \times W$ -module by defining the action of Γ by $b(w \otimes x) = \zeta^n (wb^{-1} \otimes bx)$ for $b \in \Gamma_0, x \in V_n^{(\zeta)}$, which we denote by $\Gamma\text{-Ind}_{W_L}^W V^{(\zeta)}$.

1.4. In the remainder of this paper, we assume that G is simple modulo center. Let $T \subset B$ be a pair of maximal torus and a Borel subgroup of G . Put $W = N_G(T)/T$. Let L be a Levi subgroup of a parabolic subgroup P of G containing B such that $L \supset T$. We have $W_L = N_L(T)/T$. Let $\Phi \subset X(T)$ be a root system for G with respect to T , with a simple root system Π (with respect to B), where $X(T)$ is the character group of T . We denote by Φ_L the sub system of Φ corresponding to L with the simple root system $\Pi_L \subset \Pi$. Let Π' be the set of simple roots which are orthogonal to Π_L with respect to the standard inner product on $V = \mathbf{R} \otimes_{\mathbf{Z}} X(T)$. We denote by L' the Levi subgroup containing T corresponding to Π' . Let $W_{L'} = N_{L'}(T)/T$ be the Weyl group of L' . Then we have $W \supset W_L \times W_{L'}$, and so $W_{L'} \subset N_W(W_L)$.

We recall here the notion of regular elements of reflection groups due to Springer [Sp1]. Let W be a reflection group in $GL(V)$. A vector $v \in V$ is called regular if v is not contained in any reflecting hyperplane in V . An element $a \in W$ is called regular if a has an eigenvector v which is a regular element in V . If $av = \zeta v$, with ζ a primitive e -th root of unity, then the order of a is equal to e ([Sp1, 4.2]). In particular, if a is regular of order e , there exists an eigenvalue ζ which is a primitive e -th root of unity.

The regular elements $a \in W$ in the case of classical groups are given as follows (cf. [Sp1]).

Type A_{n-1} . In this case $W = \mathfrak{S}_n$ and there are two types of regular elements.

- (a) e is a divisor of n , and a is an n/e -product of (disjoint) e -cycles in \mathfrak{S}_n .
- (b) e is a divisor of $n - 1$, and a is an $(n - 1)/e$ -product of e -cycles in \mathfrak{S}_n

Type B_n . There are two types of regular elements.

- (a) e is an odd divisor of n , and a is an n/e -product of positive cycles of length e .
- (b) e is an even divisor of $2n$, and a is a $2n/e$ -product of negative cycles of length $e/2$.

Type D_n . In this case there are 4 types of regular elements.

- (a) e is an odd divisor of n , and a is a product of positive cycles of length e .
- (b) e is an odd divisor of $n - 1$, and a is a product of positive cycle of length 1 and $(n - 1)/e$ positive cycles of length e .
- (c) n is even, and e is an even divisor of n . a is a product of negative cycles of length $e/2$.
- (d) e is an even divisor of $2n - 2$, and a is a product of $(n - 1)/e$ negative cycles of length $e/2$ and one cycle of length 1, which is positive or negative according as $(2n - 2)/e$ is even or odd.


Regular elements in the exceptional Weyl groups are listed in [Sp1].

Returning to the original setting, we consider the subgroups $W_L, W_{L'}$ of W . Let V' be the subspace of V generated by $\Pi_{L'}$. $W_{L'}$ is realized as a reflection group on V' . Assume that a is a regular element of $W_{L'}$ of order e . Let ζ be a primitive e -th root of unity, and $V(a, \zeta)$ the eigensubspace of a in V with eigenvalue ζ . Since a is regular, $V(a, \zeta)$ is not contained in any reflecting hyperplane H_α for $\alpha \in \Phi_{L'}$. We say that a is L -regular if $V(a, \zeta)$ is not contained in any H_α for $\alpha \in \Phi - \Phi_{L'}$. If L is the torus T , all the regular elements are L -regular. But if $L \neq T$, regular elements are not necessarily L -regular. For example, if L is not simple modulo center, regular elements in $W_{L'}$ are not L -regular in many cases. In the case where L is simple modulo center, L -regular elements are classified as follows.

Lemma 1.5. *Assume that L is simple modulo center.*

- (i) *If W is of type A_n, B_n, D_n , take L such that W_L is of the same type as W of rank m , and $W_{L'}$ is of type A_{n-m-1} . Then a regular element of $W_{L'}$ of type (a) in 1.4 is L -regular.*
- (ii) *If W is of type G_2, F_4 or E_8 , there does not exist L -regular elements for any $L \neq T$.*
- (iii) *Assume that W is of type E_6 or E_7 . Let $\Pi = \{\alpha_1, \dots, \alpha_7\}$ (resp. $\{\alpha_1, \dots, \alpha_6\}$) be the set of simple roots in E_7 (resp. in E_6) as in the figure. Take $\Pi_L = \{\alpha_k, \alpha_{k+1}, \dots, \alpha_7\}$ (resp. $\{\alpha_k, \alpha_{k+1}, \dots, \alpha_6\}$) for $k \geq 3$. Then $W_{L'}$ is of type A_j or of type $A_j + A_1$ for some j except the case where W is of type E_7 and $\Pi_L = \{\alpha_7\}$, in which case $\Pi_{L'}$ is of type D_5 . In the former case, we choose a a regular element of type (a) for type A , and in the latter case, we choose a a regular element of type (a) for type D in 1.4, respectively. Then a is L -regular.*

Proof. If there exists $\beta \in \Phi - \Phi_L$ such that β is orthogonal to $V_{L'}$, then any regular element in $W_{L'}$ cannot be L -regular. By direct inspections, one can find such β unless L is the type given in (i), (iii) of the lemma. Assume that L is as in the



lemma, and let a be a regular element in $W_{L'}$. If $W_{L'}$ is of type A_j or type $A_j + A_1$, then a regular vector $v \in V'$ can be written explicitly, and one can check the L -regularity by direct inspections. If $W_{L'}$ is of type D_5 (in the case where W is of type E_7), a must be of type (a) (otherwise it is easy to see that a is not L -regular). But this element is nothing but the regular element in A_4 , and the checking is reduced to the previous case. The details are omitted. \square

1.6. In what follows we consider a specific cyclic group $\Gamma \in N_W(W_L)$, and $u \in L$ according to the following two cases.

Case (a): $W_{L'} \neq \{1\}$.

In this case, we assume that L is simple modulo center. We choose an L -regular element $a \in W_{L'}$, and put $\Gamma = \langle a \rangle$. Let e be the order of Γ . Thus $\Gamma \subset W_{L'}$ and we have $\Gamma \times W_L \subset W$. We take any unipotent element $u \in L$.

Case (b): $W_{L'} = \{1\}$.

In this case, we assume that L is of type $X_0 + e(A_{n_1-1} + \cdots + A_{n_r-1})$ with X_0 irreducible. We further assume that any $\beta \in \Phi - \Phi_L$ is not orthogonal to the root system $e(A_{n_1-1} + \cdots + A_{n_r-1})$. (Note: since $W_{L'} = \{1\}$, any irreducible component of the Dynkin diagram corresponding to $\Pi - \Pi_L$ consists of 1 or 2 nodes. The latter condition is satisfied for type B_n if all the irreducible components consist of one node, and for type A_n, D_n if the number of irreducible components having two nodes is at most 1.)

We choose $a \in W$ so that a permutes each component A_{n_i-1} in a cyclic way, and acts trivially on X_0 . Thus $a \in \mathfrak{S}_{en_1} \times \cdots \times \mathfrak{S}_{en_r}$, and a is a product of disjoint cycles of length e . In particular, $\Gamma = \langle a \rangle \subset N_W(W_L)$, and the subgroup of W generated by Γ and W_L coincides with the semidirect product $\Gamma \ltimes W_L$. Now L is isogenic to $G_0 \times G_1 \times \cdots \times G_r$ modulo center, where G_0 is of type X_0 , and $G_i \simeq GL_{n_i} \times \cdots \times GL_{n_i}$ (e -factors). We choose a unipotent element $u \in L$ so that u corresponds to (u_0, u_1, \dots, u_r) , where $u_0 \in G_0$ is arbitrary, and u_i is a diagonal element in G_i , i.e., $u_i = (v_i, \dots, v_i)$ with $v_i \in GL_{n_i}$ for $i = 1, \dots, r$.

We can state our main theorem, whose proof will be given in the next section.

Theorem 1.7. *Assume that G is defined over \mathbf{C} . Let L be a Levi subgroup in G . Assume that a cyclic subgroup Γ of order e in $N_W(W_L)$ and $u \in L$ are given as in 1.4. Put $\widetilde{W}_L = \Gamma \ltimes W_L$. Then the followings hold.*

- (i) W_L -module $H^*(\mathcal{B}_u^L)$ can be extended to a \widetilde{W}_L -module so that $\Gamma \times \widetilde{W}_L$ -module $H^*(\mathcal{B}_u^L)^{(\zeta')}$ is defined for any e -th root of unity ζ' .
- (ii) There exists a primitive e -th root of unity ζ such that

$$(1.7.1) \quad \Gamma\text{-Ind}_{W_L}^W (H^*(\mathcal{B}_u^L)^{(\zeta)}) \simeq H^*(\mathcal{B}_u)^{(\zeta)}$$

as $\Gamma \times W$ -modules.

Remarks 1.8. (i) The extension of W_L -module $H^*(\mathcal{B}_u^L)$ to \widetilde{W}_L -module is not unique. The theorem asserts that the statement (ii) holds for some choice of extension.

(ii) The theorem asserts that (1.7.1) holds for some choice of primitive e -th root of unity ζ , but then it holds for any choice of primitive root of unity ζ' . In fact, we can write $\zeta' = \zeta^j$ for some j prime to e , and we have an automorphism τ on Γ such that $\tau(a) = a^j$. It follows from (1.7.1) that we have an isomorphism of $\Gamma \times W$ modules, where the action of Γ is twisted by τ . It is easy to check that the twisted $\Gamma \times W$ -module $\Gamma\text{-Ind}_{W_L}^W(H^*(\mathcal{B}_u^L)^{(\zeta)})$ is isomorphic to $\Gamma\text{-Ind}_{W_L}^W(H^*(\mathcal{B}_u^L)^{(\zeta')})$, and similarly the twisted $H^*(\mathcal{B}_u)^{(\zeta)}$ is isomorphic to $H^*(\mathcal{B}_u)^{(\zeta')}$. Thus (1.7.1) holds also for ζ' .

2. PROOF OF THEOREM 1.7

2.1. In the case where $e = 1$, Theorem 1.7 is nothing but the original induction theorem. So we assume that $e \geq 2$ in what follows. Since the structure of the W -module $H^n(\mathcal{B}_u)$ is independent of p provided that p is a good prime, it is enough to show the corresponding formula for an appropriate p . So, we assume that G is defined over \mathbf{F}_p , of split type, with Frobenius map F . We assume that $T \subset B$ are both F -stable, and that $L \subset P$ are F -stable. Thus F acts trivially on W and on W_L . We first note that

Lemma 2.2. *Let $a \in N_W(W_L)$ and choose $\dot{a} \in N_G(T) \cap N_G(L)$. Assume that $\dot{a} \in Z_G(u)$. Then $\text{ad } \dot{a}$ stabilizes \mathcal{B}_u^L , and acts on $H^*(\mathcal{B}_u^L)$ in such a way that $\text{ad } \dot{a}(w) = awa^{-1}$ for $w \in W_L$.*

Proof. Since $\dot{a} \in N_G(L)$, \dot{a} acts on \mathcal{B}^L by the adjoint action $\text{ad } \dot{a}$, which stabilizes \mathcal{B}_u^L since $\dot{a} \in Z_G(u)$. Hence \dot{a} acts naturally on $H^*(\mathcal{B}_u^L)$. In order to compare this action with the action of W_L , we shall recall the construction of Springer representations of W_L . Let

$$\widetilde{L} = \{(x, gB) \in L \times \mathcal{B}^L \mid g^{-1}xg \in B\},$$

and $\pi : \widetilde{L} \rightarrow L$ be the first projection. Let L_r be the set of regular semisimple elements in L . Then $\pi^{-1}(L_r)$ is isomorphic to

$$\widetilde{L}_r = T_r \times L/T,$$

where $T_r = T \cap L_r$. Let $\pi_0 : \widetilde{L}_r \rightarrow L_r$ be the map defined by $\pi_0 : (t, gT) \mapsto g^{-1}tg$, which coincides with the restriction of π on \widetilde{L}_r under the identification $\pi^{-1}(L_r) \simeq \widetilde{L}_r$. Then π_0 is an unramified Galois covering with group W_L , and for a constant sheaf $\overline{\mathbf{Q}}_l$ on \widetilde{L}_r , $\mathcal{L} = \pi_* \overline{\mathbf{Q}}_l$ is a W_L -equivariant local system on L_r . Thus $K = \text{IC}(L, \mathcal{L})$ is a W_L -equivariant complex on L , and it is known by Lusztig that $K \simeq \pi_* \overline{\mathbf{Q}}_l$. Thus for each $u \in L$, the stalk $\mathcal{H}_u^i(K)$ at u of the i -th cohomology sheaf of K gives rise to a W_L -module $H^i(\mathcal{B}_u^L)$.

Now \dot{a} acts on \tilde{L}_r (resp. on L_r) by $\text{ad } \dot{a} : (t, gT) \mapsto (\dot{a}t\dot{a}^{-1}, \dot{a}g\dot{a}^{-1}T)$ (resp. $\text{ad } \dot{a} : x \mapsto \dot{a}x\dot{a}^{-1}$), and π_0 commutes with $\text{ad } \dot{a}$. Hence \mathcal{L} becomes an \dot{a} -equivariant local system. Since $\pi_0^{-1}(t) = \{(wtw^{-1}, wT) \mid w \in W_L\}$ for $t \in T_r$, the stalk \mathcal{L}_t has a natural structure of the regular W_L -module. Then the isomorphism $\mathcal{L}_{\dot{a}t\dot{a}^{-1}} \rightarrow \mathcal{L}_t$ is given by $\text{ad } \dot{a}^{-1}$ under the identification $\mathcal{L}_x \simeq \bar{\mathbf{Q}}_l[W_L]$ for $x \in L_r$. It follows that \mathcal{L} is $\langle \dot{a} \rangle \times W_L$ -equivariant, where $\langle \dot{a} \rangle$ is a cyclic group generated by \dot{a} , and \dot{a} acts on W_L by $\text{ad } \dot{a}(w) = awa^{-1}$. By the functoriality of IC functor, K turns out to be a \dot{a} -equivariant complex on L under the adjoint action of \dot{a} , which is regarded as a $\langle \dot{a} \rangle \times W_L$ -equivariant complex on L . Hence for $u \in L$ such that $\dot{a}u\dot{a}^{-1} = u$, $\mathcal{H}_u^i(K)$ has a structure of $\langle \dot{a} \rangle \times W_L$ -module.

On the other hand, \dot{a} acts naturally on \tilde{L} and on L by the adjoint action, which commute with π . Thus $\pi_*\bar{\mathbf{Q}}_l$ is \dot{a} -equivariant, which is isomorphic to K as the complex with \dot{a} -action. Hence the action of \dot{a} on $\mathcal{H}_u^i(K)$ coincides with the action on $H^i(\mathcal{B}_u^L)$ induced from the adjoint action of \dot{a} on \mathcal{B}_u^L . The lemma follows from this. \square

Next we show the following lemma.

Lemma 2.3. *There exists a representative $\dot{a} \in N_G(T) \cap N_G(L) \cap Z_G(u)$ such that \dot{a} acts trivially on $H^*(\mathcal{B}_u)$ and that \dot{a}^e acts trivially on $H^*(\mathcal{B}_u^L)$. In particular, $H^*(\mathcal{B}_u^L)$ has a structure of \tilde{W}_L -module.*

Proof. First consider the case (a) in 1.6. Let H be the subgroup of G generated by U_α with $\alpha \in \Phi_{L'}$, where U_α is the root subgroup corresponding to α . Then H is a connected reductive subgroup of L' whose Weyl group coincides with $W_{L'}$. Since $H \subset Z_G(u)$, we have $H \subset Z_G^0(u)$. One can choose a representative $\dot{a} \in N_H(T_1)$ of $a \in W_{L'}$, where T_1 is a maximal torus of H contained in T . Then $\dot{a} \in Z_G^0(u) \cap N_G(L)$ and $\dot{a}^e \in T_1$. Since $T_1 \subset Z_G(u)$, we see that $T_1 \subset Z_L^0(u)$. Thus, $\dot{a}^e \in Z_L^0(u)$. Hence \dot{a} satisfies the condition.

Next consider the case (b) in 1.6. Let L_1 be the Levi subgroup containing L of type $X_{n_0} + A_{en_1-1} + \cdots + A_{en_r-1}$. We have a natural projection $\pi : L_1 \rightarrow \bar{L}_1 = L_1/Z^0(L_1)$, and an isogeny map $\theta : \tilde{L}_1 = G_0 \times SL_{en_1} \times \cdots \times SL_{en_r} \rightarrow \bar{L}_1$, where G_0 is the simply connected semisimple group of type X_0 . Put $\bar{u} = \pi(u) \in \bar{L}_1$. Now $Z_{L_1}(u)$ acts on $H^*(\mathcal{B}_u)$. Since $Z^0(L_1)$ acts trivially on $H^*(\mathcal{B}_u)$, we have an action of $Z_{L_1}(u)/Z^0(L_1) = Z_{\bar{L}_1}(\bar{u})$ on $H^*(\mathcal{B}_u)$. Let \tilde{u} be an element in \tilde{L}_1 such that $\theta(\tilde{u}) = \bar{u}$. $\tilde{u} = (u_0, u_1, \dots, u_r)$ can be chosen as given in 1.4. We choose $\dot{a} \in \tilde{L}_1$ as follows; put $\dot{a} = (a_0, a_1, \dots, a_r)$ with $a_0 \in G_0$, and $a_i \in SL_{en_i}$ for $1 \leq i \leq r$. We put $a_0 = 1$ and choose a_1, \dots, a_r so that $a_i \in Z_{SL_{en_i}}^0(u_i)$ and that $a_i^e \in Z(SL_{en_i})$. Such a choice is always possible for u_i of type (n_i, \dots, n_i) . Thus $\dot{a} \in Z_{\tilde{L}_1}^0(\tilde{u})$. It follows that $\theta(\dot{a})$ is contained in a connected subgroup of $Z_{\bar{L}_1}(\bar{u}_1)$, and by the previous remark, $\theta(\dot{a})$ acts trivially on $H^*(\mathcal{B}_u)$. Now take $\dot{a} \in Z_{L_1}(u)$ such that $\pi(\dot{a}) = \theta(\dot{a})$. Then $\dot{a} \in N_G(T) \cap N_G(L)$, and acts trivially on $H^*(\mathcal{B}_u)$. On the other hand, similar to π, θ , we have a map $\pi' : L \rightarrow \bar{L} = L/Z^0(L)$ and $\theta' : \tilde{L} = G_0 \times (SL_{n_1})^e \times \cdots \times (SL_{n_r})^e \rightarrow \bar{L}$. Let $\bar{u} = \pi'(u) \in \bar{L}$, and $\tilde{u} \in \tilde{L}$ such that $\bar{u} = \theta'(\tilde{u})$. Then we have an isomorphism $H^*(\mathcal{B}_u^L) \simeq H^*(\mathcal{B}_{\bar{u}}^{\bar{L}}) \simeq H^*(\mathcal{B}_{\tilde{u}}^{\tilde{L}})$ compatible with the actions of $Z_L(u)$, $Z_{\bar{L}}(\bar{u})$ and $Z_{\tilde{L}}(\tilde{u})$ with respect to π', θ' . We have $\dot{a}^e \in Z(SL_{n_1})^e \times Z(SL_{n_2})^e \times \cdots$. Since the action of

$Z(SL_{n_1})^e \times Z(SL_{n_2})^e \times \cdots$ can be extended to an action of $Z(GL_{n_1})^e \times Z(GL_{n_2})^e \times \cdots$ on $H^*(\mathcal{B}_u^L)$, \tilde{a}^e acts trivially on $H^*(\mathcal{B}_u^L)$, and so \tilde{a}^e acts trivially on $H^*(\mathcal{B}_u^L)$. \square

2.4. Let $\mathcal{Z} = Z_L^0$ be the identity component of the center of L . Put $\mathcal{B}_{\mathcal{Z}} = \{B' \in \mathcal{B} \mid \mathcal{Z} \subset B'\}$. Then $\mathcal{B}_{\mathcal{Z}}$ is decomposed into connected components

$$\mathcal{B}_{\mathcal{Z}} = \coprod_{d \in W_L \backslash W} \mathcal{B}_{\mathcal{Z},d},$$

where $\mathcal{B}_{\mathcal{Z},d} = \{x^d B \mid x \in L\}$, which is isomorphic to \mathcal{B}^L under the map $B' \mapsto B' \cap L$.

Put

$$\mathcal{Z}_{\text{reg}} = \{z \in \mathcal{Z} \mid Z_G^0(z) = L\}.$$

Then for any $t \in \mathcal{Z}_{\text{reg}}$, we have $\mathcal{B}_t = \mathcal{B}_{\mathcal{Z}}$ by Lemma 2.2 (c) in [L3], and so $\mathcal{B}_{tu} = \mathcal{B}_u \cap \mathcal{B}_t = \mathcal{B}_u \cap \mathcal{B}_{\mathcal{Z}}$. It follows that

$$\mathcal{B}_{tu} = \coprod_{d \in W_L \backslash W} (\mathcal{B}_{\mathcal{Z},d} \cap \mathcal{B}_u),$$

where $\mathcal{B}_{\mathcal{Z},d} \cap \mathcal{B}_u$ is isomorphic to \mathcal{B}_u^L under the map $B' \mapsto B' \cap L$. This implies that

$$(2.4.1) \quad H^{2n}(\mathcal{B}_{tu}) \simeq \bigoplus_{d^{-1} \in W/W_L} H^{2n}(\mathcal{B}_{\mathcal{Z},d} \cap \mathcal{B}_u).$$

The right hand side of (2.4.1) has a natural structure of the induced W -module $\text{Ind}_{W_L}^W H^{2n}(\mathcal{B}_u^L)$. It is proved in [L3, Proposition 1.4] that (2.4.1) is actually an isomorphism of W -modules. Let $a \in W$ be as in the theorem. Since $\tilde{a} \in N_G(L)$, it stabilizes \mathcal{Z} , and so \tilde{a} acts on $\mathcal{B}_{\mathcal{Z}}$ via $\text{ad } \tilde{a}$. It is easy to see that \tilde{a} induces a permutation action on the components of $\mathcal{B}_{\mathcal{Z}}$; $\tilde{a} : \mathcal{B}_{\mathcal{Z},d} \mapsto \mathcal{B}_{\mathcal{Z},ad}$. It follows that \tilde{a} induces an automorphism on $H^{2n}(\mathcal{B}_{tu})$, which maps the factor corresponding to $d^{-1} \in W/W_L$ to $d^{-1}a^{-1} \in W/W_L$. Under the isomorphism $H^{2n}(\mathcal{B}_{\mathcal{Z},d} \cap \mathcal{B}_u) \simeq H^{2n}(\mathcal{B}_u^L)$, the factor corresponding to $d^{-1} \in W/W_L$ is written as $d^{-1} \otimes H^{2n}(\mathcal{B}_u^L)$, and \tilde{a} maps $d^{-1} \otimes H^{2n}(\mathcal{B}_u^L) \rightarrow d^{-1}a^{-1} \otimes H^{2n}(\mathcal{B}_u^L)$. On the other hand, by Lemma 2.3, \tilde{a}^e acts trivially on $H^{2n}(\mathcal{B}_u^L)$, and induces an action of \widetilde{W}_L on it. Hence \tilde{a} induces an action of Γ_0 on $H^{2n}(\mathcal{B}_{tu}) \simeq \text{Ind}_{W_L}^W H^{2n}(\mathcal{B}_u^L)$, which is given by $\tilde{a} : d^{-1} \otimes x \mapsto d^{-1}a^{-1} \otimes \tilde{a}x$ for each factor $d^{-1} \otimes H^{2n}(\mathcal{B}_u^L)$.

Now we define an action of Γ on $H^*(\mathcal{B}_{tu})$ by $a : x \mapsto \zeta^n \tilde{a}x$ for $x \in H^{2n}(\mathcal{B}_{tu})$, where $\tilde{a}x$ is the action of Γ_0 on $H^{2n}(\mathcal{B}_{tu})$ given as above. Since the action of $\tilde{a} \in G$ commutes with that of W , $H^*(\mathcal{B}_{tu})$ turns out to be a $\Gamma \times W$ -module, which we denote by $H^*(\mathcal{B}_{tu})^{[\zeta]}$. The following lemma is immediate from the above discussion.

Lemma 2.5. *There exists an isomorphism of $\Gamma \times W$ -modules*

$$H^*(\mathcal{B}_{tu})^{[\zeta]} \simeq \Gamma\text{-Ind}_{W_L}^W H^*(\mathcal{B}_u^L)^{(\zeta)}.$$

In view of Lemma 2.5, in order to prove the theorem it is enough to show the following proposition.

Proposition 2.6. *Under an appropriate choice of (a good prime) p , there exists an isomorphism of $\Gamma \times W$ -modules for any $t \in \mathcal{Z}_r$,*

$$H^*(\mathcal{B}_u)^{(\zeta)} \simeq H^*(\mathcal{B}_{tu})^{[\zeta]}.$$

2.7. The remainder of this section is devoted to the proof of the proposition. We shall prove it by modifying the arguments in [L3]. By [Sh1], [Sh2], [BS], the following fact is known; assume that G is simple modulo center. Then for each unipotent class C of G , there exists $u_1 \in C^F$, called a split unipotent element, such that F acts on $H^{2n}(\mathcal{B}_{u_1})$ as a scalar multiplication by p^n . (In the case where G is of type E_8 , we assume that $p \equiv 1 \pmod{4}$). Since the component group $A_G(u_1) = Z_G(u_1)/Z_G^0(u_1)$ is isomorphic to S_3, S_4, S_5 or $(\mathbf{Z}/2\mathbf{Z})^k$ for some k , there exists a positive integer s_0 (independent of p) such that F^{s_0} acts on $H^{2n}(\mathcal{B}_u)$ by a scalar multiplication by $p^{s_0 n}$ for any unipotent element u of G^F (e.g., one can take $s_0 = |S_5|$.) Similarly, F^{s_0} acts on $H^{2n}(\mathcal{B}_u^L)$ by a scalar multiplication by $p^{s_0 n}$ for any unipotent element $u \in L^F$. Note that the isomorphism in (2.4.1) is F -equivariant. Hence F^{s_0} acts also as a scalar multiplication by $p^{s_0 n}$ for $H^{2n}(\mathcal{B}_{tu})$.

Note that \dot{a} acts trivially on $H^{2n}(\mathcal{B}_u)$ by Lemma 2.3. It follows that one can write

$$(2.7.1) \quad \text{Tr}((F^s \dot{a})^i w, H^*(\mathcal{B}_u)) = \sum_{n \geq 0} a_n(w) p^{isn},$$

$$(2.7.2) \quad \text{Tr}((w, a^i), H^*(\mathcal{B}_u)^{(\zeta)}) = \sum_{n \geq 0} a_n(w) \zeta^{in},$$

for any $w \in W, 0 \leq i \leq e - 1$ and for any positive integer s divisible by s_0 , where $a_n(w) = \text{Tr}(w, H^n(\mathcal{B}_u))$ are integers for each $n \geq 0$.

On the other hand, by the description of the action of F and of \dot{a} on $H^n(\mathcal{B}_{tu})$ in 2.4, together with Lemma 2.5, one can write

$$(2.7.3) \quad \text{Tr}((F^s \dot{a})^i w, H^*(\mathcal{B}_{tu})) = \sum_{n \geq 0} b_{n,i}(w) p^{isn},$$

$$(2.7.4) \quad \text{Tr}((w, a^i), H^*(\mathcal{B}_{tu})^{[\zeta]}) = \sum_{n \geq 0} b_{n,i}(w) \zeta^{in},$$

for w, i, s as above, where $b_{n,i}(w)$ are certain integers.

For an integer x and a prime number l , we denote by $m_l(x)$ the multiplicative order of x in $\mathbf{Z}/l\mathbf{Z}$, i.e., the smallest positive integer m such that $x^m \equiv 1 \pmod{l}$. The following is a key for the proof of Proposition 2.6.

Lemma 2.8. *Assume that $p \equiv 1 \pmod{4}$. Let s_0, e be fixed positive integers coprime to p . Then there exist infinitely many prime numbers l satisfying the following properties.*

- (i) $m_l(p^s) = e$ for a certain integer s divisible by s_0 .
- (ii) $l - 1$ is divisible by e .

Proof. By our assumption, the image of s_0e on $\mathbf{F}_p = \mathbf{Z}/p\mathbf{Z}$ is non-zero. Hence the map $x \mapsto s_0ex + 1$ induces a bijective map on \mathbf{F}_p . Thus there exists $c \in \mathbf{Z}$ such that the image of $s_0ec + 1$ in \mathbf{F}_p is contained in $\mathbf{F}_p^* - (\mathbf{F}_p^*)^2$. Put $\alpha = s_0ec + 1$. Then α is prime to p , and so $(\alpha - 1)p$ and α are coprime each other. Then by Dirichlet's theorem on arithmetic progression, there exist infinitely many prime numbers l of the form $l = n(\alpha - 1)p + \alpha$ for some positive integer n . It is enough to show that these $l \geq 3$ satisfy the assertion of the lemma. For an integer a and a prime number p , let $\left(\frac{a}{p}\right)$ be the Legendre symbol, i.e.,

$$\left(\frac{a}{p}\right) = \begin{cases} 1 & \text{if } x^2 \equiv a \pmod{p} \text{ for some } x \in \mathbf{Z}, \\ -1 & \text{otherwise.} \end{cases}$$

We show that

$$(2.8.1) \quad \left(\frac{p}{l}\right) = -1.$$

In fact, by the quadratic reciprocity law (e.g., [Se]), we have

$$\left(\frac{p}{l}\right)\left(\frac{l}{p}\right) = (-1)^{\frac{p-1}{2} \cdot \frac{l-1}{2}} = 1.$$

The second equality follows from the assumption that $p \equiv 1 \pmod{4}$. Hence we have $\left(\frac{p}{l}\right) = \left(\frac{l}{p}\right)$. But $l \equiv \alpha \pmod{p}$, and so $\left(\frac{l}{p}\right) = \left(\frac{\alpha}{p}\right) = -1$ since the image of α is not contained in \mathbf{F}_p^2 by our choice of α . Hence (2.8.1) holds.

Now (2.8.1) is equivalent to $p^{(l-1)/2} \equiv -1 \pmod{l}$. It follows that $m_l(p) = l - 1$. Since $l - 1 = s_0ec(np + 1)$, we see that $m_l(p^s) = e$ for $s = s_0c(np + 1)$ and that $l - 1$ is divisible by e . Thus this l satisfies the assertion of the lemma. The lemma is proved. \square

2.9 For given integers $s_0 \geq 1, e \geq 2$, we choose a prime number p such that p is not a factor of e, s_0 and that $p \equiv 1 \pmod{4}$, and fix it once and for all. For a multiple s of s_0 , put $F' = F^s a$ and $q = p^s$. Under the setting in 1.6, we shall describe the set \mathcal{Z}_{reg} more precisely. As in [L3, Lemma 2.2], \mathcal{Z}_{reg} can be written as $\mathcal{Z}_{\text{reg}} = \mathcal{Z} - \bigcup_{\beta} \ker(\beta|_{\mathcal{Z}})$, where β runs over all the roots in $\Phi - \Phi_L$. ($\beta|_{\mathcal{Z}}$ gives a non-trivial character of \mathcal{Z} for $\beta \in \Phi - \Phi_L$).

First consider the case (a). Let L'_{der} be the derived subgroup of L' , and S' be the split maximal torus of L'_{der} contained in T . Then $S' \subset \mathcal{Z}$. Put $S'_{\text{reg}} = S' \cap \mathcal{Z}_{\text{reg}}$. Now $W_{L'}$ leaves the set $\Phi - \Phi_L$ invariant. For each $\beta \in \Phi - \Phi_L$, put $H_{\beta} = \bigcap_{x \in \Gamma} \ker(x(\beta)|_{S'})$. Then H_{β} is an F' -stable subgroup of S' , and we see that

$$(2.9.1) \quad S'^{F'}_{\text{reg}} = S'^{F'} - \bigcup_{\beta \in \Phi - \Phi_L} H_{\beta}^{F'}.$$

H_{β} is a closed subgroup of S' , and we put $e_{\beta} = |H_{\beta}/H_{\beta}^0|$ for each $\beta \in \Phi - \Phi_L$.

Let \mathcal{P}' be the set of all prime numbers l satisfying the condition in Lemma 2.8. Thus \mathcal{P}' is an infinite set. We denote by \mathcal{P} the subset of \mathcal{P}' consisting of l such that $l > |\Phi - \Phi_L|$ and that l does not divide e_β ($\beta \in \Phi - \Phi_L$). Thus \mathcal{P} is an infinite set also.

Next we consider the case (b). We may assume that G has a connected center of dimension 1, and that the derived subgroup of G is simply connected, almost simple. Let k be an algebraic closure of \mathbf{F}_q . We see that there exists a subtorus S of \mathcal{Z} such that $S \simeq (k^*)^c$, where c is the number of irreducible components of Φ_L . Since a permutes the factors k^* in S , we see that $S^{F'} \simeq (\mathbf{F}_{q^e}^*)^r \times (\mathbf{F}_q^*)^{r'}$, where r' is equal to 1 or 0 according to the cases where X_0 is non-empty or empty. Since $\Gamma \subset N_W(W_L)$, Γ preserves the set $\Phi - \Phi_L$. For each $\beta \in \Phi - \Phi_L$, put $K_\beta = \bigcap_{x \in \Gamma} \ker(x(\beta)|_S)$. Then K_β is an F' -stable subgroup of S , and we have

$$(2.9.2) \quad S_{\text{reg}}^{F'} = S^{F'} - \bigcup_{\beta \in \Phi - \Phi_L} K_\beta^{F'},$$

where $S_{\text{reg}} = S \cap \mathcal{Z}_{\text{reg}}$. K_β is a closed subgroup of S , and put $e_\beta = |K_\beta/K_\beta^0|$ for each $\beta \in \Phi - \Phi_L$. Under the identification $S^{F'} \simeq (\mathbf{F}_{q^e}^*)^r \times (\mathbf{F}_q^*)^{r'}$, we see that $K_\beta^{0F'} \simeq (\mathbf{F}_{q^e}^*)^{r-1} \times (\mathbf{F}_q^*)^{r'}$ or $K_\beta^{0F'} \simeq \mathbf{F}_{q^{e'}}^* \times (\mathbf{F}_{q^e}^*)^{r-1} \times (\mathbf{F}_q^*)^{r'}$, where e' is a proper divisor of e . (Let S_i be the subtorus of S corresponding to the factor eA_{n_i-1} for $i = 1, \dots, r$. Then the former case occurs if $\beta|_{S_i}, \beta|_{S_j}$ are non-trivial for some $i \neq j$, and the latter case occurs if $\beta|_{S_i}$ is non-trivial for only one i . Note that by our assumption in 1.4, β is non-trivial on $S_1 \times \dots \times S_r$.)

Let \mathcal{P}' be as in the case (a). We define a subset \mathcal{P} of \mathcal{P}' as the set of prime numbers $l \in \mathcal{P}'$ such that $l > |\Phi - \Phi_L|$ and that l does not divide e_β .

The next lemma is a variant of Lemma 3.4 in [L3].

Lemma 2.10. *Assume that $l \in \mathcal{P}$, and let s be a multiple of s_0 such that $m_l(p^s) = e$ (see Lemma 2.8). Put $F' = F^s a$. Then there exists $t \in \mathcal{Z}_{\text{reg}}$ such that $F'(t) = t$ and that $t^l = 1$.*

Proof. First consider the case (a) in 1.6. It is enough to show, for each $l \in \mathcal{P}$, that there exists $t \in S_{\text{reg}}^{F'}$ such that $t^l = 1$. Note that a is a regular element of order e in $W_{L'}$. Put $V = \mathbf{R} \otimes_{\mathbf{Z}} X(S')$. Thus $W_{L'}$ acts on V as a reflection group. Let ζ be a primitive e -th root of unity, and let $a(e)$ be the dimension of the eigenspace $V(a, \zeta) \subset V$ of a with eigenvalue ζ . We show that

$$(2.10.1) \quad \#\{t \in S^{F'} \mid t^l = 1\} = l^{a(e)}.$$

By a general formula, we have $|S^{F'}| = |\det_V(qI - a)| = P_a(q)$, where $P_a(x)$ is the characteristic polynomial of $a \in W_{L'}$. Since a is regular $P_a(x)$ can be written, by [Sp1, 4.2], as

$$P_a(x) = \Phi_e(x)^{a(e)} \Phi'(x),$$

where $\Phi_e(x)$ is the cyclotomic polynomial of degree e , and $\Phi'(x)$ is a product of cyclotomic polynomials $\Phi_{e'}(x)$ with $e' < e$. By our assumption $m_l(q) = e$, $\Phi_e(q)$ is divisible by l , and $\Phi'(q)$ is not divisible by l . This means that each minimal F' -stable

torus M of S' corresponding to the factor $\Phi_e(x)$ contains an element of order l . Since $\{t \in M^{F'} \mid t^l = 1\} \subset \mathbf{F}_{q^e}^*$, $M^{F'}$ contains exactly l elements t such that $t^l = 1$. Thus (2.10.1) is proved.

For $\beta \in \Phi - \Phi_L$, let V_β be the subspace of V which is orthogonal to $x(\beta)$ for all $x \in \Gamma$. Then V_β can be identified with $\mathbf{R} \otimes_{\mathbf{Z}} X(H_\beta^0)$. Γ stabilizes V_β , and let $V_\beta(a, \zeta)$ be the eigenspace of a on V_β with eigenvalue ζ . Since a is L -regular, we have $\dim V_\beta(a, \zeta) < \dim V(a, \zeta) = a(e)$. It follows that the characteristic polynomial $P'_a(x)$ of a on V_β contains the factor $\Phi_e(x)$ with multiplicity less than $a(e)$. By a similar argument as above, minimal F' -stable subtori of H_β^0 corresponding to $\Phi_e(x)$ only contain elements of order l . This implies that

$$\#\{t \in H_\alpha^{F'} \mid t^l = 1\} = \#\{t \in H_\alpha^{0F'} \mid t^l = 1\} \leq l^{a(e)-1}.$$

It follows, by (2.9.1), that

$$\begin{aligned} \#\{t \in S'_{\text{reg}}{}^{F'} \mid t^l = 1\} &= \#\{t \in S'^{F'} \mid t^l = 1, t \notin \bigcup_{\beta \in \Phi - \Phi_L} H_\beta^{F'}\} \\ &\geq l^{a(e)} - Nl^{a(e)-1} = l^{a(e)-1}(l - N), \end{aligned}$$

where $N = |\Phi - \Phi_L|$. Since $l > N$ by our assumption, there exists $t \in S'_{\text{reg}}{}^{F'}$ such that $t^l = 1$. This proves the lemma in the case (a).

Next consider the case (b) in 1.6. It is enough to show, for each $l \in \mathcal{P}$, that there exists $t \in S'_{\text{reg}}{}^{F'}$ such that $t^l = 1$. We note that $q^{e'} - 1$ is not divisible by l for any divisor $e' < e$ of e by the assumption $m_l(q) = e$. Since $S^{F'} \simeq (\mathbf{F}_{q^e}^*)^r \times (\mathbf{F}_q^*)^{r'}$ (cf. 2.9), we have

$$\#\{t \in S^{F'} \mid t^l = 1\} = l^r.$$

We consider K_β given in 2.9. By the discussion in 2.9, we have

$$\#\{t \in K_\beta^{F'} \mid t^l = 1\} = \#\{t \in K_\beta^{0F'} \mid t^l = 1\} = l^{r-1}.$$

It follows, by (2.9.2), that

$$\begin{aligned} \#\{t \in S'_{\text{reg}}{}^{F'} \mid t^l = 1\} &= \#\{t \in S^{F'} \mid t^l = 1, t \notin \bigcup_{\beta \in \Phi - \Phi_L} K_\beta^{F'}\} \\ &\geq l^r - Nl^{r-1} = l^{r-1}(l - N), \end{aligned}$$

where N is as before. Since $l > N$ by our assumption, the lemma holds also for the case (b). \square

We need the following lemma due to Lusztig.

Lemma 2.11 ([L3, Lemma 3.2]). *Let H be a finite group, and ϕ a virtual character of H (over a field of characteristic 0). Assume that ϕ is integral valued. Let $x, y \in H$ be such that $xy = yx$ and $y^l = 1$ for a prime number l . Then $\phi(xy) - \phi(x) \in l\mathbf{Z}$.*

2.12. Let s_0 be as in 2.7, and \mathcal{P} be as in 2.9. Let $F' = F^s \dot{a}$ be as in Lemma 2.10 for a fixed $l \in \mathcal{P}$. Let $R_{w,i} = R_{T_w}(1)$ be the Deligne-Lusztig's virtual character of $G^{F'^i}$ for $i = 1, \dots, e$, where T_w is an F'^i -stable maximal torus of G corresponding to $w \in W \simeq W(T_1)$ (here $W(T_1) = N_G(T_1)/T_1$ for an F' -stable pair $T_1 \subset B_1$). Let us choose $t \in \mathcal{Z}_{\text{reg}}$ as in Lemma 2.10. Then we have

$$(2.12.1) \quad \begin{aligned} \text{Tr}(F'^i w, H^*(\mathcal{B}_u)) &= \text{Tr}(u, R_{w,i}), \\ \text{Tr}(F'^i w, H^*(\mathcal{B}_{tu})) &= \text{Tr}(tu, R_{w,i}). \end{aligned}$$

We remark that (2.12.1) was proved in [L2] under the assumption that p^s is large enough (which is determined only by the data of the Dynkin diagram of G). Thus if we replace s_0 in 2.7 by a suitable large number, the result in [L2] is applicable. One can also apply [Sh3, Theorem 2.2] instead of [L2], where the restriction on p^s is removed.

Since $R_{w,i}$ are integral valued, one can apply Lemma 2.11 for $H = G^{F'^i}$ and $x = u, y = t$. Hence we have

$$\text{Tr}(u, R_{w,i}) = \text{Tr}(tu, R_{w,i}) \pmod{l\mathbf{Z}}.$$

It follows from (2.12.1) that

$$(2.12.2) \quad \text{Tr}(F'^i w, H^*(\mathcal{B}_u)) = \text{Tr}(F'^i w, H^*(\mathcal{B}_{tu})) \pmod{l\mathbf{Z}}.$$

Let ζ_0 be a fixed primitive e -th root of unity in \mathbf{C} , and R the ring of integers of the cyclotomic field $\mathbf{Q}(\zeta_0)$. Let \mathcal{I} be the set of non-zero prime ideals \mathfrak{p} in R such that \mathfrak{p} contains one of the numbers $1 - \zeta_0^i$ for $i = 1, \dots, e - 1$ and ζ_0 . Let $\bar{\mathcal{I}}$ be the set of prime numbers l such that $\mathfrak{p} \cap \mathbf{Z} = l\mathbf{Z}$ for $\mathfrak{p} \in \mathcal{I}$. Since \mathcal{I} is a finite set, $\bar{\mathcal{I}}$ is a finite set. So, $\mathcal{P} - \bar{\mathcal{I}}$ is an infinite set. Let \mathcal{J} be the set of prime ideals \mathfrak{p} of R such that $\mathfrak{p} \cap \mathbf{Z} = l\mathbf{Z}$ with $l \in \mathcal{P} - \bar{\mathcal{I}}$. Then \mathcal{J} is an infinite set. Now R/\mathfrak{p} is a finite extension of \mathbf{F}_l . Let $\bar{\zeta}_0$ be the image of ζ_0 in R/\mathfrak{p} . Since $l \in \mathcal{P}$, $l - 1$ is divisible by e . Hence $\bar{\zeta}_0 \in \mathbf{F}_l^*$, which has order e by our choice of \mathfrak{p} . Since $m_l(p^s) = e$, the image of p^s in $\mathbf{Z}/l\mathbf{Z}$ has order e . Hence there exists j such that

$$(2.12.3) \quad p^s - \zeta_0^j \in \mathfrak{p}.$$

Note that the number j is determined by the choice of \mathfrak{p} , which we denote by $j(\mathfrak{p})$. For $j = 1, \dots, e - 1$, let \mathcal{J}_j be the set of prime ideals \mathfrak{p} in \mathcal{J} such that $j(\mathfrak{p}) = j$. Thus $\mathcal{J} = \bigcup_j \mathcal{J}_j$, and so there exists j_0 such that $\mathcal{J}_0 = \mathcal{J}_{j_0}$ is an infinite set. We put $\zeta = \zeta_0^{j_0}$. By (2.12.3), ζ is a primitive e -th root of unity.

We remark that $H^*(\mathcal{B}_{tu}) = H^*(\mathcal{B}_u \cap \mathcal{B}_{\mathcal{Z}})$ is independent of the choice of $t \in \mathcal{Z}_{\text{reg}}$. Then in view of (2.7.1) \sim (2.7.4), together with (2.12.3), we see that

$$\begin{aligned} \text{Tr}((F^s \dot{a})^i w, H^*(\mathcal{B}_u)) &= \text{Tr}((w, a^i), H^*(\mathcal{B}_u)^{(\zeta)}) \pmod{\mathfrak{p}}, \\ \text{Tr}((F^s \dot{a})^i w, H^*(\mathcal{B}_u \cap \mathcal{B}_{\mathcal{Z}})) &= \text{Tr}((w, a^i), H^*(\mathcal{B}_u \cap \mathcal{B}_{\mathcal{Z}})^{[\zeta]}) \pmod{\mathfrak{p}} \end{aligned}$$

for any $\mathfrak{p} \in \mathcal{J}_0$. Combined with (2.12.2), we have

$$\mathrm{Tr}((w, a^i), H^*(\mathcal{B}_u)^{(\zeta)}) = \mathrm{Tr}((w, a^i), H^*(\mathcal{B}_u \cap \mathcal{B}_Z)^{[\zeta]}) \pmod{\mathfrak{p}}$$

for $\mathfrak{p} \in \mathcal{J}_0$. Since \mathcal{J}_0 is an infinite set, we conclude that

$$\mathrm{Tr}((w, a^i), H^*(\mathcal{B}_u)^{(\zeta)}) = \mathrm{Tr}((w, a^i), H^*(\mathcal{B}_u \cap \mathcal{B}_Z)^{[\zeta]}).$$

Hence Proposition 2.6 is proved, and the theorem follows.

3. APPLICATIONS

3.1. Let W_L be the subgroup of W , and Γ the subgroup of W generated by $a \in N_W(W_L)$ such that Γ and W_L generate the semidirect product group $\widetilde{W}_L = \Gamma \ltimes W_L$. Let $V = V^{(\zeta)}$ be the $\Gamma \times \widetilde{W}_L$ -module as in 1.3. (We write Γ as Γ_0 if it is regarded as a subgroup of \widetilde{W}_L , cf. 1.3.) Then V can be decomposed as $V = \bigoplus_{i \in \mathbf{Z}/e\mathbf{Z}} V^{(i)}$, where $V^{(i)}$ is the eigenspace of $a \in \Gamma$ with eigenvalue ζ^i , which is a \widetilde{W}_L -submodule of V . Then we have

$$\begin{aligned} \mathrm{Ind}_{W_L}^W V &= \bigoplus_i \bigoplus_{w \in W/\widetilde{W}_L} \bigoplus_j wa^j \otimes V^{(i)} \\ &= \bigoplus_i \bigoplus_{w \in W/\widetilde{W}_L} \bigoplus_k wb_k \otimes V^{(i)}, \end{aligned}$$

where $b_k = \sum_j \zeta^{jk} a^j \in \mathbf{C}[\Gamma]$ (the group ring of Γ). For each $i \in \mathbf{Z}$, let $\psi^{(i)}$ the linear character of Γ defined by $\psi^{(i)}(a) = \zeta^i$. Then Γ -module $\mathbf{C}b_k$ is afforded by $\psi^{(-k)}$. Let $V_n^{(i)}$ be the eigenspace of $a \in \Gamma_0$ on the \widetilde{W}_L -module $V^{(i)}$ with eigenvalue ζ^n . Let $(\Gamma\text{-Ind}_{W_L}^W V)^{(k)}$ be the eigenspace of $a \in \Gamma$ with eigenvalue ζ^k . Then we have the following lemma.

Lemma 3.2. (i) *Let the notations be as above. We have*

$$(3.2.1) \quad (\Gamma\text{-Ind}_{W_L}^W V)^{(k)} \simeq \bigoplus_{w \in W/\widetilde{W}_L} \bigoplus_{j \in \mathbf{Z}/e\mathbf{Z}} \bigoplus_{0 \leq n < e} wb_{k-n-j} \otimes V_n^{(j)}$$

as vector spaces. In particular, $\dim(\Gamma\text{-Ind}_{W_L}^W V^{(\zeta)})^{(k)}$ is independent of the choice of $k \in \mathbf{Z}/e\mathbf{Z}$, which is given by

$$(3.2.2) \quad \dim(\Gamma\text{-Ind}_{W_L}^W V)^{(k)} = [W : \widetilde{W}_L] \dim V.$$

(ii) *Assume that Γ commutes with W_L . Then we have*

$$(3.2.3) \quad (\Gamma\text{-Ind}_{W_L}^W V)^{(k)} \simeq \bigoplus_{j \in \mathbf{Z}/e\mathbf{Z}} \mathrm{Ind}_{\Gamma \times W_L}^W (\psi^{(-k+j)} \otimes V^{(j)})$$

as W -modules.

Proof. Under the action of Γ on $\Gamma\text{-Ind}_{W_L}^W V$, $w b_k \otimes V_n^{(i)}$ is contained in an eigenspace of a with eigenvalue ζ^{k+n+j} . Then (i) follows easily from the discussion in 3.1. Now assume that Γ commutes with W_L . Then $b_k \otimes V^{(i)}$ has a structure of $\Gamma \times W_L$ -module given by $\psi^{(-k)} \otimes V^{(i)}$. (ii) follows from the formula (3.2.1) by noticing that $V^{(i)} = V_0^{(i)}$. The lemma is proved. \square

We consider a Levi subgroup $L \subset G$ and a unipotent element $u \in L$, and take $\Gamma = \langle a \rangle \subset N_W(W_L)$ satisfying the condition in 1.6. We apply the preceding argument to the situation $V^{(\zeta)} = H^*(\mathcal{B}_u^L)^{(\zeta)}$. Then as a corollary to Theorem 1.7, we have

Proposition 3.3. *Under the setting in Theorem 1.7, we have, for $0 \leq k \leq e - 1$,*

$$(3.3.1) \quad \bigoplus_{n \equiv k \pmod e} H^{2n}(\mathcal{B}_u) \simeq \bigoplus_{w \in W/\widetilde{W}_L} \bigoplus_{j \in \mathbf{Z}/e\mathbf{Z}} \bigoplus_{0 \leq n < e} w b_{k-n-j} \otimes H^{2j}(\mathcal{B}_u^L)_n$$

as vector spaces, where $b_i \in \mathbf{C}[\Gamma]$ and $H^{2n}(\mathcal{B}_u^L)_n$ is the eigenspace of $a \in \Gamma_0$ with eigenvalue ζ^n . In particular, $\dim\left(\bigoplus_{n \equiv k \pmod e} H^{2n}(\mathcal{B}_u)\right)$ is independent of the choice of k . In the case (a) in 1.6, (3.3.1) can be made more precise as follows;

$$(3.3.2) \quad \bigoplus_{n \equiv k \pmod e} H^{2n}(\mathcal{B}_u) \simeq \text{Ind}_{\Gamma \times W_L}^W \left(\bigoplus_{j \in \mathbf{Z}/e\mathbf{Z}} \psi^{(-k+j)} \otimes H^{2j}(\mathcal{B}_u^L) \right)$$

as W -modules.

Proof. By Theorem 1.7 $\Gamma\text{-Ind}_{W_L}^W H^*(\mathcal{B}_u^L)^{(\zeta)}$ is isomorphic to $H^*(\mathcal{B}_u)^{(\zeta)}$ as $\Gamma \times W$ -modules. Since $(H^*(\mathcal{B}_u)^{(\zeta)})^{(k)} = \bigoplus_{n \equiv k \pmod e} H^{2n}(\mathcal{B}_u)$, the corollary follows from Lemma 3.2. \square

Remarks 3.4. (i) In the case where $u = 1$, the cohomology ring $H^*(\mathcal{B}_u) = H^*(\mathcal{B})$ coincides with the coinvariant algebra R of W . In the special case where G is of type A_{n-1} , i.e., $W \simeq \mathfrak{S}_n$, we consider $W_L \simeq \mathfrak{S}_{n-re}$ for $1 \leq e \leq n$. Then $W_{L'} \simeq \mathfrak{S}_{re}$, and if we choose a regular element $a \in W_{L'}$ as a product of disjoint cycles of length e , Proposition 3.3 can be applied. This recovers the formula obtained by Morita and Nakajima [MN1].

More generally, consider the Weyl group W acting on the real vector space V as the reflection module. For $v \in V$, let W_v be the stabilizer of v in W , and N_v the stabilizer of the line $\mathbf{R}v$ in W . Note that W_v is normal in N_v , and W_v coincides with W_L for a certain Levi subgroup in G . Then for any $\Gamma = \langle a \rangle$ such that $\Gamma \subset N_v$, Bonnafé, Lehrer and Michel [BLM] have proved a similar formula as in Proposition 3.3. So our formula (3.3.2) can be regarded as a special case of theirs. (Note that they treat a more general case, where W is a complex reflection group and Γ is not necessarily cyclic, in a framework of coinvariant algebras.)

(ii) We consider a unipotent element $u \in L$ in the case where $G = GL_n$. $u \in G$ can be written as $u = u_\mu$ by a partition $\mu = (1^{m_1}, 2^{m_2}, \dots, n^{m_n})$ of n . Take a positive integer $e \geq 2$, and let I be a subset of $\{1, \dots, n\}$ such that $e \leq m_i$ for $i \in I$.

We consider a Levi subgroup L of type $X_0 + e \sum_{i \in I} A_{i-1}$, where $X_0 = A_k$ with $k = \sum_{i \notin I} im_i + \sum_{i \in I} i(m_i - e) - 1$. Then we have $W_{L'} = \{1\}$, and one can choose $u \in L$ so that it satisfies the assumption of the case (b) in 1.6. Thus Proposition 3.3 can be applied. This covers the results on the stability of dimensions obtained in [MN2], [MN3], where they considered the case $|I| = 1$ or the case all the m_i are divisible by e .

Returning to the general setup, we consider the case where u is a regular unipotent element in L . Then $H^*(\mathcal{B}_u^L) = H^0(\mathcal{B}_u^L) \simeq \mathbf{C}$ is a trivial W_L -module. Thus Proposition 3.3 implies the following.

Corollary 3.5. *Let G be a simple algebraic group modulo center, and L a Levi subgroup in G . Let u be a regular unipotent element in L . Let $\Gamma = \langle a \rangle$ be a subgroup of $N_W(W_L)$ of order e satisfying the conditions in 1.6. Then for $k = 0, \dots, e-1$, we have*

$$\bigoplus_{n \equiv k \pmod{e}} H^{2n}(\mathcal{B}_u) \simeq \text{Ind}_{\Gamma \rtimes W_L}^W \tilde{\psi}^{(-k)}$$

as W -modules, where $\tilde{\psi}^{(-k)}$ is the character of $\Gamma \rtimes W_L$ obtained as the pull back of $\psi^{(-k)}$ under the projection $\Gamma \rtimes W_L \rightarrow \Gamma$.

Proof. In the case (a), the assertion follows from (3.3.2). So we consider the case (b). In the setup of 3.1, $V^{(i)}$ is a trivial W_L -module \mathbf{C} for $i = 0$ and zero otherwise. Then we see that $V^{(0)} = V_0^{(0)}$, and $b_k \otimes V^{(0)}$ has a structure of \widetilde{W}_L -module $\tilde{\psi}^{(-k)}$. The assertion follows from the formula in 3.1. \square

3.6. Let G be a simple algebraic group defined over \mathbf{F}_q with Frobenius map F . We assume that G^F is of split type. The Green function Q_{T_w} is defined as the restriction of the Deligne-Lusztig's virtual character $R_{T_w}(1)$ to the set of unipotent elements in G^F . We assume that $p = \text{ch } \mathbf{F}_q$ is good, and in the case where G is of type E_8 , we further assume that $q \equiv 1 \pmod{4}$. Then as explained in 2.7, for each unipotent class C of G , there exists a split element $u \in C^F$. As in 2.12, we have

$$(3.6.1) \quad Q_{T_w}(u) = \sum_{n \geq 0} \text{Tr}(w, H^{2n}(\mathcal{B}_u)) q^n.$$

Hence there exists a polynomial $\mathbf{Q}_{w,C}(x) \in \mathbf{Z}[x]$ such that $Q_{T_w}(u) = \mathbf{Q}_{w,C}(q)$. Concerning the values of Green functions at root of unity, we have the following.

Proposition 3.7. *Suppose that G, L and $u \in L$ are as in Corollary 3.5. Then we have*

$$(3.7.1) \quad \mathbf{Q}_{w,C}(\zeta^j) = |W_L|^{-1} \sharp \{x \in W \mid x^{-1}wx \in a^j W_L\}$$

for $j = 0, \dots, e-1$. In particular, the value $\mathbf{Q}_{w,C}(\zeta')$ is independent of the choice of a primitive e -th root of unity ζ' .

Proof. Put $c_i(w) = \#\{x \in W \mid x^{-1}wx \in a^iW_L\}$ for $i = 0, \dots, e-1$. Then

$$\begin{aligned} (\text{Ind}_{\Gamma \times W_L}^W \tilde{\psi}^{(-k)})(w) &= |\Gamma \times W_L|^{-1} \sum_{i=0}^{e-1} \sum_{\substack{x \in W \\ x^{-1}wx \in a^iW_L}} \tilde{\psi}^{(-k)}(x^{-1}wx) \\ &= |\Gamma \times W_L|^{-1} \sum_{i=0}^{e-1} c_i(w) \zeta^{-ki}. \end{aligned}$$

It follows, by (3.6.1) together with Corollary 3.5, that

$$\begin{aligned} \mathbf{Q}_{w,C}(\zeta^j) &= \sum_{k=0}^{e-1} \zeta^{kj} \sum_{n \equiv k \pmod{e}} \text{Tr}(w, H^{2n}(\mathcal{B}_u)) \\ &= |\Gamma \times W_L|^{-1} \sum_{i=0}^{e-1} c_i(w) \sum_{k=0}^{e-1} \zeta^{(j-i)k} \\ &= |W_L|^{-1} c_j(w). \end{aligned}$$

Hence we obtain the formula (3.7.1). Let ζ^j be a primitive e -th root of unity. There exists an element $\tau \in \text{Gal}(\mathbf{Q}(\zeta)/\mathbf{Q})$ such that $\tau(\zeta) = \zeta^j$. By (3.6.1), we see that $\mathbf{Q}_{w,C}(\zeta) \in \mathbf{Q}(\zeta)$ and that $\tau(\mathbf{Q}_{w,C}(\zeta)) = \mathbf{Q}_{w,C}(\zeta^j)$. But since $\mathbf{Q}_{w,C}(\zeta) \in \mathbf{Z}$ by (3.7.1), we conclude that $\mathbf{Q}_{w,C}(\zeta) = \mathbf{Q}_{w,C}(\zeta^j)$. This proves the proposition. \square

Remark 3.8. In the case where $G = GL_n$ and L is of type $A_{m-1} + \dots + A_{m-1}$ (e -times) with $n = em$, take a regular unipotent element u in L . Then $u = u_\mu \in G$ with $\mu = (m^e)$. For $w \in W = \mathfrak{S}_n$, let $\lambda(w) = (1^{l_1}, 2^{l_2}, \dots)$ be the partition of n corresponding to the cycle decomposition of w . Then one can show by a direct computation (cf. [M, (6.2)]) that

$$|W_L|^{-1} \#\{x \in W \mid x^{-1}wx \in aW_L\} = \begin{cases} e^{l(\lambda(w))} & \text{if } e \mid l_i \text{ for all } i, \\ 0 & \text{otherwise,} \end{cases}$$

where $l(\lambda)$ is the number of parts for a partition λ . Thus we recover the formula in [LLT, Theorem 3.2, Theorem 3.4] concerning the values of Green polynomials of GL_n at roots of unity.

3.9. We give some more examples where Proposition 3.3 can be applied.

(i) Assume that G is of type B_n and L is a Levi subgroup of type B_m with $m < n$. Then L' is of type A_{n-m-1} . For any $u \in L$ and a divisor e of $n-m$, the proposition can be applied. Similar results hold also for C_n or D_n .

(ii) Assume that $G = Sp_{2n}$. Then a unipotent element $u \in G$ can be written as $u = u_\mu$ as an element of GL_{2n} , where $\mu = (1^{m_1}, 2^{m_2}, \dots)$ is a partition of $2n$ such that m_i is even for odd i . Take an even integer $e \geq 2$, and let I be a subset of odd integers $\{1, 3, \dots, 2n-1\}$ such that $e \leq m_i$ for $i \in I$. We consider a Levi subgroup L of type $X_0 + e \sum_{i \in I} A_{i-1}$, where X_0 is of type C_k with $2k = \sum_{i \notin I} im_i + \sum_{i \in I} i(m_i - e)$.

Then $W_{L'} = \{1\}$, and as in Remarks 3.4 (ii), one can find $u \in L$ so that the case (b) in 1.6 can be applied. Similar results hold also for type B_n and D_n .

(iii) Assume that G is of type E_7 , and choose L of type A_2 so that L' is of type A_4 . Take any unipotent element $u \in L$. Then the proposition can be applied with $e = 5$.

REFERENCES

- [AL] D. Alvis and G. Lusztig, On Springer's correspondence for simple groups of type E_n ($n = 6, 7, 8$), *Math. Proc. Camb. Phil. Soc.* **92** (1982), 65–72.)
- [BLM] C. Bonnafé, G.I. Lehrer and J. Michel, Twisted invariant theory for reflection groups, preprint.
- [BS] W.M. Beynon and N. Spaltenstein, Green functions of finite Chevalley groups of type E_n ($n = 6, 7, 8$), *J. Algebra* (1984), 584–614.
- [L1] G. Lusztig, Green polynomials and singularities of unipotent classes, *Adv. in Math.* **42** (1981), 169–178.
- [L2] G. Lusztig, Green functions and character sheaves, *Annals of Mathematics*, **131** (1990), 355–408.
- [L3] G. Lusztig, An induction theorem for Springer's representations, *Adv. Studies in Pure Math.*, **40**, In "Representation theory of algebraic groups and quantum groups", pp. 253 – 259, 2004.
- [LLT] A. Lascoux, B. Leclerc and J.-Y. Thibon, Green polynomials and Hall-Littlewood functions at roots of unity, *Euro. J. Combi.* **15** (1994), 173–180.
- [M] H. Morita, Decomposition of Green polynomials of type A and De Concini-Procesi-Tanisaki algebras of certain types, preprint.
- [MN1] H. Morita and T. Nakajima, The coinvariant algebra of the symmetric group as a direct sum of induced modules, *Osaka J. Math.*, **42** (2005), 217–231.
- [MN2] H. Morita and T. Nakajima, A formula of Lascoux-Leclerc-Thibon and representations of symmetric groups, preprint.
- [MN3] H. Morita and T. Nakajima, The Green polynomials at roots of unity and its applications, preprint.
- [Se] J.-P. Serre, "A course in Arithmetic" GTM, Springer-Verlag, 1973.
- [Sh1] T. Shoji, On the Green polynomials of Chevalley groups of type F_4 , *Comm. in Alg.* **10**, (1982), 505–543.
- [Sh2] T. Shoji, On the Green polynomials of classical groups, *Invent. Math.* **74**, (1983), 237–267.
- [Sh3] T. Shoji, Character sheaves and almost characters of reductive groups, *Adv. in Math.* **111** (1995), 244–313.
- [Sp1] T.A. Springer, Regular elements of finite reflection groups, *Invent. Math.* **25**, (1974), 159–198.
- [Sp2] T.A. Springer, Trigonometric sums, Green functions of finite groups and representations of Weyl groups, *Invent. Math.* **36** (1976), 173–207.
- [St] J.R. Stembridge, On the eigenvalues of representations of reflection groups and wreath products, *Pacific J. Math.*, **140** (1989), 353–396.