

RINGS WITH q -TORSIONFREE CANONICAL MODULES

NAOKI ENDO, LAURA GHEZZI, SHIRO GOTO, JOOYOUN HONG, SHIN-ICHIRO IAI,
TOSHINORI KOBAYASHI, NAOYUKI MATSUOKA, AND RYO TAKAHASHI

Dedicated to the memory of Wolmer V. Vasconcelos

ABSTRACT. Let A be a Noetherian local ring with canonical module K_A . We characterize A when K_A is a torsionless, reflexive, or q -torsionfree module for an integer $q \geq 3$. If A is a Cohen-Macaulay ring, H.-B. Foxby proved in 1974 that the A -module K_A is q -torsionfree if and only if the ring A is q -Gorenstein. With mild assumptions, we provide a generalization of Foxby's result to arbitrary Noetherian local rings admitting the canonical module. In particular, since the reflexivity of the canonical module is closely related to the ring being Gorenstein in low codimension, we also explore quasi-normal rings, introduced by W. V. Vasconcelos. We provide several examples as well.

1. INTRODUCTION

This paper investigates the question of the structure of a Noetherian local ring A if its canonical module K_A is a torsionless, reflexive, or more generally, q -torsionfree A -module for an integer $q \geq 3$. The notion of q -torsionfree modules was one of the central contributions of the famous research of M. Auslander and M. Bridger [1]. It turned out to be an important property in H.-B. Foxby's study of q -Gorenstein rings [11]. Among many interesting results, Foxby settled the above question in the case where A is a Cohen-Macaulay ring. More precisely, the A -module K_A is q -torsionfree if and only if the ring A is q -Gorenstein, i.e., $A_{\mathfrak{p}}$ is a Gorenstein ring for every $\mathfrak{p} \in \text{Spec } A$ with $\text{depth } A_{\mathfrak{p}} < q$ (see [10, Proposition 3.2]). It remains unclear what happens if we do not assume the ring A is Cohen-Macaulay. The theory of canonical modules nowadays has been developed mainly over Cohen-Macaulay rings in connection with the Gorenstein property; see e.g., [6, 13, 14, 15, 19]. However, over Noetherian local (not necessarily Cohen-Macaulay) rings, there are also remarkable preceding researches on canonical modules, including the study of their endomorphism algebras; see [2, 3, 5]. Therefore, behaviors of canonical modules, even for non-Cohen-Macaulay rings, are interesting and the q -torsionfree property is well worth studying. The motivation for the present

2020 *Mathematics Subject Classification.* 13H10, 13A02, 13A15.

Key words and phrases. Canonical module, Gorenstein ring, Cohen-Macaulay ring, q -torsionfree module, q -Gorenstein ring, quasi-normal ring.

N. Endo was partially supported by JSPS Grant-in-Aid for Young Scientists 20K14299. L. Ghezzi was partially supported by the Fellowship Leave from the New York City College of Technology-CUNY (Fall 2022-Spring 2023) and by a grant from the City University of New York PSC-CUNY Research Award Program Cycle 53. S. Goto was partially supported by JSPS Grant-in-Aid for Scientific Research (C) 21K03211. J. Hong was partially supported by the Sabbatical Leave Program at Southern Connecticut State University (Spring 2022). T. Kobayashi was partly supported by JSPS Grant-in-Aid for JSPS Fellows 21J00567. N. Matsuoka was partially supported by JSPS Grant-in-Aid for Scientific Research (C) 18K03227. R. Takahashi was partially supported by JSPS Grant-in-Aid for Scientific Research (C) 19K03443.

research started with this question that arose while the second and fourth authors were writing the last paper with Vasconcelos concerning (torsionless) canonical modules [4].

To explain our results more precisely, let us start from definitions which we will use throughout this paper. For a Noetherian local ring A of dimension d with maximal ideal \mathfrak{m} , a *canonical module* K of A is a finitely generated A -module satisfying

$$\widehat{A} \otimes_A K \cong \mathrm{Hom}_{\widehat{A}}(\mathrm{H}_{\widehat{\mathfrak{m}}}^d(\widehat{A}), \widehat{E})$$

where $\mathrm{H}_{\widehat{\mathfrak{m}}}^d(\widehat{A})$ denotes the d^{th} local cohomology module of the $\widehat{\mathfrak{m}}$ -adic completion \widehat{A} of A with respect to its maximal ideal $\widehat{\mathfrak{m}}$ and \widehat{E} is the injective hull of the \widehat{A} -module $\widehat{A}/\widehat{\mathfrak{m}}$ ([14, Definition 5.6]). Equivalently, a finitely generated A -module K is a canonical module of A if $\mathrm{Hom}_A(K, E) \cong \mathrm{H}_{\mathfrak{m}}^d(A)$, where $\mathrm{H}_{\mathfrak{m}}^d(A)$ is the d^{th} local cohomology module of A with respect to \mathfrak{m} and E is the injective hull of A/\mathfrak{m} ([5, Definition 12.1.2, Remarks 12.1.3]). The canonical module K_A is uniquely determined up to isomorphisms ([2, (1.5)], see also [14, Lemma 5.8]) if it exists. Although the existence is not guaranteed even for Cohen-Macaulay local domains, provided A is Cohen-Macaulay, the ring A admits the canonical module if and only if A is a homomorphic image of a Gorenstein ring ([16, 18]). The fundamental theory of canonical modules over Cohen-Macaulay rings was developed in the monumental book [14] of J. Herzog and E. Kunz. We shall in this paper freely refer to [14] for basic results on canonical modules (see [6, Chapter 3] also).

We now continue to state our setup. Let R be a Noetherian (not necessarily local) ring. For an R -module M , we have a canonical homomorphism

$$\varphi : M \rightarrow M^{**}$$

defined by $[\varphi(x)](f) = f(x)$ for each $f \in M^*$ and $x \in M$, where $(-)^* = \mathrm{Hom}_R(-, R)$ denotes the R -dual functor. We say that M is *torsionless* (resp. *reflexive*) if φ is injective (resp. bijective). Torsionless modules are torsionfree, i.e., there is no nonzero torsion elements, and the converse holds if the total ring of fractions $\mathrm{Q}(R)$ of R is Gorenstein ([21, Theorem (A.1)]). Moreover, the R -module M is torsionless (resp. reflexive) if and only if $\mathrm{Ext}_R^i(\mathrm{D}(M), R) = (0)$ for $i = 1$ (resp. $i = 1, 2$), where $\mathrm{D}(M)$ denotes the Auslander transpose of M ([1]). From this point of view, Auslander and Bridger introduced a *q -torsionfree module* M to be $\mathrm{Ext}_R^i(\mathrm{D}(M), R) = (0)$ for all $i = 1, 2, \dots, q$. In addition, for an integer n , we say that

- M satisfies (S_n) if $\mathrm{depth} M_{\mathfrak{p}} \geq \min\{n, \dim R_{\mathfrak{p}}\}$ for every $\mathfrak{p} \in \mathrm{Spec} R$,
- M satisfies (\widetilde{S}_n) if $\mathrm{depth} M_{\mathfrak{p}} \geq \min\{n, \mathrm{depth} R_{\mathfrak{p}}\}$ for every $\mathfrak{p} \in \mathrm{Spec} R$,
- R satisfies (G_n) if $R_{\mathfrak{p}}$ is Gorenstein for every $\mathfrak{p} \in \mathrm{Spec} R$ with $\dim R_{\mathfrak{p}} \leq n$,
- R satisfies (\widetilde{G}_n) if $R_{\mathfrak{p}}$ is Gorenstein for every $\mathfrak{p} \in \mathrm{Spec} R$ with $\mathrm{depth} R_{\mathfrak{p}} \leq n$.

The condition (S_n) is known as Serre's condition. A Noetherian ring satisfying (\widetilde{G}_n) coincides with $(n+1)$ -Gorenstein ring in earlier publications such as [1, 11]. The condition (\widetilde{G}_n) is equivalent to saying that the ring satisfies both (S_{n+1}) and (G_n) .

Let us now state our results, explaining how this paper is organized. In Section 2, after recalling the necessary definitions and preliminaries, we give a criterion for a Noetherian local ring A to have the torsionless canonical module. We show that the A -module K_A is torsionless if and only if $A_{\mathfrak{p}}$ is Gorenstein for every $\mathfrak{p} \in \mathrm{Assh} A$, where $\mathrm{Assh} A = \{\mathfrak{p} \in \mathrm{Spec} A \mid \dim A/\mathfrak{p} = \dim A\} =$

$\text{Ass}_A K_A$ (Proposition 2.3). Section 3 is devoted to the characterizations of local rings A with reflexive canonical modules. When $\dim A = 1$, this is exactly the case where A is a Gorenstein ring (Proposition 3.3). We elaborate on the one-dimensional case in Section 5. For the higher dimensional case, the reflexivity of K_A is characterized by the local ring $A_{\mathfrak{p}}$ being Gorenstein for every $\mathfrak{p} \in \text{Supp}_A K_A$ with $\dim A_{\mathfrak{p}} \leq 1$ and $\text{Ass } A \cap V(U) = \text{Assh } A$, where U denotes the unmixed component of (0) in A and $V(U)$ is the set of all prime ideals in A containing U (Theorem 3.6). This lead us to obtain Corollary 3.8, which claims that K_A is reflexive if and only if A satisfies (G_1) , provided $\text{Assh } A = \text{Ass } A$. This indicates that the reflexivity of canonical modules is deeply related to the ring being Gorenstein in low codimension. Thus Section 4 is dedicated to quasi-normal rings, i.e., rings with (S_2) and (G_1) , which were introduced by Vasconcelos. In Section 6, we generalize Foxby's result on q -torsionfree canonical modules to arbitrary Noetherian local rings A admitting a canonical module. Our results of Sections 2 and 3 provide a complete generalization in case $q = 1, 2$. When $q \geq 3$, Theorem 6.6 states that the A -module K_A is q -torsionfree if and only if the ring A satisfies (G_{q-1}) and (S_{q-1}) on $\text{Supp}_A K_A$, provided that K_A satisfies (S_q) . In the final section we provide concrete examples of Cohen-Macaulay and q -Gorenstein rings in order to illustrate our theorems.

2. TORSIONLESS CANONICAL MODULES

Throughout the section, let (A, \mathfrak{m}) be a Noetherian local ring of dimension d . We begin with some preliminaries. Let $(0) = \bigcap_{\mathfrak{p} \in \text{Ass } A} Q(\mathfrak{p})$ denote a primary decomposition of (0) in A . We set

$$\text{Assh } A = \{\mathfrak{p} \in \text{Spec } A \mid \dim A/\mathfrak{p} = d\} \quad \text{and} \quad U = \bigcap_{\mathfrak{p} \in \text{Assh } A} Q(\mathfrak{p})$$

where U is called the *unmixed component* of (0) in A . Let $V(U)$ denote the set of all prime ideals of A containing U .

Lemma 2.1. *There is an embedding $0 \rightarrow A/U \rightarrow A$ of A -modules.*

Proof. We may assume that $U \neq (0)$. Then $\text{Assh } A \subsetneq \text{Ass } A$. Let

$$L = \bigcap_{\mathfrak{p} \in \text{Ass } A \setminus \text{Assh } A} Q(\mathfrak{p}).$$

We then have $L \not\subseteq \bigcup_{\mathfrak{p} \in \text{Assh } A} \mathfrak{p}$. Choose an element $a \in L$ but $a \notin \bigcup_{\mathfrak{p} \in \text{Assh } A} \mathfrak{p}$. Since a is a non-zero-divisor on A/U and $aU \subseteq L \cap U = (0)$, we obtain $((0) :_A a) = U$. Then U is the kernel of the homomorphism $\varphi : A \rightarrow A$ given by $\varphi(1) = a$. Thus, $A/U \cong \text{Im}(\varphi) \hookrightarrow A$. \square

In the rest of this section, we assume the ring A admits the canonical module K_A . We recall several known facts about K_A which we will use throughout this article; see [2, (1.6), (1.7), (1.8), (1.9), (1.10), Theorem 3.2] and [14, Korollar 6.3] (also [5, Chapter 12]) for the proofs.

Proposition 2.2. *The following assertions hold true.*

(1) *The annihilator of K_A is U . In particular, $\dim_A K_A = d$ and $\text{Ass}_A K_A = \text{Assh } A$.*

- (2) If $a \in \mathfrak{m}$ is A -regular, then a is K_A -regular.
- (3) $V(U) = \text{Supp}_A K_A = \{\mathfrak{p} \in \text{Spec } A \mid \dim A = \dim A/\mathfrak{p} + \text{ht}_A \mathfrak{p}\}$.
- (4) Both K_A and $\text{Hom}_A(K_A, K_A)$ satisfy (S_2) .
- (5) $K_{A_{\mathfrak{p}}} = [K_A]_{\mathfrak{p}}$ for every $\mathfrak{p} \in \text{Supp}_A K_A$.
- (6) $\text{Supp}_A K_A = \text{Spec } A$ if and only if $\text{Min } A = \text{Assh } A$.
- (7) $\text{Ass } A = \text{Assh } A$ if and only if K_A is a faithful A -module.
- (8) Suppose A is Cohen-Macaulay. If $a \in \mathfrak{m}$ is A -regular, then $K_{A/(a)}$ exists and $K_{A/(a)} \cong K_A/aK_A$.
- (9) Suppose that $K_{A/U}$ exists. Then $K_{A/U}$, as an A -module, is the canonical module of A .

By Proposition 2.2-(2), the canonical module K_A is torsionfree as an A -module. In general, torsionless modules are torsionfree, and the converse holds if and only if $A_{\mathfrak{p}}$ is a Gorenstein local ring for every $\mathfrak{p} \in \text{Ass } A$ ([21, Theorem (A.1)]). Therefore, if the total ring of fractions $Q(A)$ of A is Gorenstein, then K_A is torsionless. The following proposition shows the case where K_A is torsionless without assuming $Q(A)$ is Gorenstein. It is also a generalization of [4, Proposition 3.2].

Proposition 2.3. *The following conditions are equivalent:*

- (1) K_A is a torsionless A -module;
- (2) $A_{\mathfrak{p}}$ is a Gorenstein ring for every $\mathfrak{p} \in \text{Assh } A$;
- (3) $K_A \cong I$ for some ideal I of A .

Proof. (1) \Rightarrow (2) Since K_A is torsionless, there exists an exact sequence $0 \rightarrow K_A \rightarrow F$ of A -modules, where F is a finitely generated free A -module. Let $\mathfrak{p} \in \text{Assh } A \subseteq \text{Supp}_A K_A$. Then $A_{\mathfrak{p}}$ is Artinian and $[K_A]_{\mathfrak{p}}$ is the canonical module of $A_{\mathfrak{p}}$. Therefore, we may assume that $[K_A]_{\mathfrak{p}}$ is the injective hull of $A_{\mathfrak{p}}/\mathfrak{p}A_{\mathfrak{p}}$. The splitting monomorphism $0 \rightarrow [K_A]_{\mathfrak{p}} \rightarrow F_{\mathfrak{p}}$ induces that $[K_A]_{\mathfrak{p}}$ is a direct summand of the free $A_{\mathfrak{p}}$ -module $F_{\mathfrak{p}}$. Since $A_{\mathfrak{p}}$ is Artinian, by the Matlis duality, we have $[K_A]_{\mathfrak{p}} \cong A_{\mathfrak{p}}$. Hence $A_{\mathfrak{p}}$ is a Gorenstein ring.

(2) \Rightarrow (3) Let $W = A \setminus \bigcup_{\mathfrak{p} \in \text{Assh } A} \mathfrak{p}$. By assumption, $[K_A]_{\mathfrak{p}} \cong A_{\mathfrak{p}}$ for every $\mathfrak{p} \in \text{Assh } A$. Thus, $W^{-1}K_A \cong W^{-1}A$. Moreover, we have $W^{-1}A \cong W^{-1}(A/U)$ because $W^{-1}U = (0)$ by the proof of Lemma 2.1. Since every element of W is a non-zero-divisor on both K_A and A/U , the isomorphism $W^{-1}K_A \cong W^{-1}(A/U)$ induces the embedding $K_A \hookrightarrow A/U$. By Lemma 2.1, there is an embedding $K_A \hookrightarrow A$.

(3) \Rightarrow (1) is clear. □

If A is *reduced*, which means there are no nonzero nilpotents, then the local ring $A_{\mathfrak{p}}$ is a field for every $\mathfrak{p} \in \text{Ass } A$. Hence we obtain the following.

Corollary 2.4. *If A is a reduced ring, then $K_A \cong I$ for some ideal I of A .*

We recall that if A is Cohen-Macaulay, the canonical module K_A has rank one if and only if the ring A is *generically Gorenstein*, i.e., $A_{\mathfrak{p}}$ is a Gorenstein local ring for every $\mathfrak{p} \in \text{Min } A$. When one of the equivalent conditions of [6, Proposition 3.3.18] is satisfied, the canonical module can be identified with an ideal of A (see also [14, Satz 6.21]). The example below shows that the assumption $\mathfrak{p} \in \text{Assh } A$ is necessary for Proposition 2.3.

Example 2.5. Let $S = k[[X, Y, Z]]$ be the formal power series ring over a field k and set $A = S/[(X) \cap (Y, Z)^2]$. Let x, y, z denote the images of X, Y, Z in A , respectively. Then we have

$$U = (x) \text{ and } K_A \cong A/(x) \cong y^2A.$$

By Proposition 2.3, K_A is torsionless. However, A is not generically Gorenstein. In fact, $A_{\mathfrak{q}}$ is not a Gorenstein ring for $\mathfrak{q} = (y, z) \in \text{Min } A$.

3. REFLEXIVE CANONICAL MODULES

Let (A, \mathfrak{m}) be a Noetherian local ring of dimension d admitting the canonical module K_A . We denote by U the unmixed component of (0) in A . In this section we will show how the reflexivity of the canonical module is related to the Gorensteinness of the ring. We begin with the following simple but effective lemma.

Lemma 3.1. *Suppose that K_A is reflexive. Then $\text{Ass } A \cap V(U) = \text{Assh } A$. In particular, if K_A is reflexive and $\text{depth } A = 0$, then $\dim A = 0$.*

Proof. The assertion follows from

$$V(U) \cap \text{Ass } A = \text{Supp}_A K_A^* \cap \text{Ass } A = \text{Ass}_A \text{Hom}_A(K_A^*, A) = \text{Ass}_A K_A^{**} = \text{Ass}_A K_A = \text{Assh } A. \quad \square$$

The example below shows that the reflexivity of K_A may require a rather strong restriction on A .

Example 3.2. Let $S = k[[X, Y]]$ be the formal power series ring over a field k and set $A = S/[(X) \cap (X^2, Y)]$. Let x, y denote the images of X, Y in A , respectively. Let $\mathfrak{m} = (x, y)$ be the maximal ideal of A . Then we have $\text{Assh } A = \{(x)\}$, $U = (x)$, and $K_A = A/U$.

- (1) Let $\mathfrak{p} = (x)$. Since $A_{\mathfrak{p}}$ is a field, by Proposition 2.3, K_A is torsionless.
- (2) Since $\text{depth } A = 0$ and $\dim A = 1$, by Lemma 3.1, K_A is not reflexive.

Proposition 3.3. *Suppose $d = 1$. Then K_A is a reflexive A -module if and only if A is a Gorenstein ring.*

Proof. Suppose that K_A is reflexive. By Lemma 3.1, A is Cohen-Macaulay. Since K_A is reflexive, there exists an exact sequence $0 \rightarrow K_A \rightarrow F_1 \rightarrow F_0$, where F_0, F_1 are finite free A -modules [11, Proposition 2.1]. Let $a \in \mathfrak{m}$ be an A -regular element. Since A is Cohen-Macaulay, K_A/aK_A is the canonical module of A/aA . Moreover, the embedding $0 \rightarrow K_A/aK_A \rightarrow F_1/aF_1$ proves that $K_{A/aA}$ is torsionless. Therefore, by Proposition 2.3, A/aA is a Gorenstein ring. Thus, A is a Gorenstein ring. The converse is clear. \square

Remark 3.4. There exist non-Cohen-Macaulay local rings with reflexive canonical module. Example 6.2 shows a two-dimensional non-Cohen-Macaulay local ring A with K_A reflexive. The example also shows that, even if K_A is reflexive, the equality $\text{Ass } A = \text{Assh } A$ does not hold true in general.

Recall that a finitely generated A -module M is reflexive, i.e., the canonical map $\varphi : M \rightarrow M^{**}$ is an isomorphism, if and only if there is at least one isomorphism $M \cong M^{**}$ of A -modules.

Lemma 3.5. *Suppose that there is an exact sequence*

$$0 \rightarrow K_A \rightarrow K_A^{**} \rightarrow C \rightarrow 0$$

of A -modules. If $C \neq (0)$, then $A_{\mathfrak{p}}$ is a Cohen-Macaulay ring with $\dim A_{\mathfrak{p}} = 1$ for every $\mathfrak{p} \in \text{Ass}_A C$ with $\text{depth } A_{\mathfrak{p}} \geq 1$. In particular, $\mathfrak{p} \in V(U)$ and $U_{\mathfrak{p}} = (0)$.

Proof. Let $\mathfrak{p} \in \text{Ass}_A C$ such that $\text{depth } A_{\mathfrak{p}} \geq 1$. Then $\mathfrak{p} \in \text{Supp}_A K_A = V(U)$. Since $[K_A]_{\mathfrak{p}} \cong K_{A_{\mathfrak{p}}}$, by passing to the ring $A_{\mathfrak{p}}$, we may assume $\text{depth } A > 0$ and $\text{depth}_A C = 0$. Since $\text{Ass}_A K_A^{**} \subseteq \text{Ass } A$, we have $\text{depth}_A K_A^{**} \geq 1$. From the exact sequence $0 \rightarrow K_A \rightarrow K_A^{**} \rightarrow C \rightarrow 0$, we obtain

$$0 = \text{depth}_A C \geq \min\{\text{depth}_A K_A - 1, \text{depth}_A K_A^{**}\}.$$

Thus, $\text{depth}_A K_A = 1$. Since K_A satisfies (S_2) , we have

$$1 = \text{depth}_A K_A \geq \min\{2, \dim A\}.$$

Therefore, A is a Cohen-Macaulay ring of dimension 1. In particular, $U = (0)$. \square

Now we aim to generalize Proposition 3.3.

Theorem 3.6. *The following conditions are equivalent:*

- (1) K_A is a reflexive A -module;
- (2) $\text{Ass } A \cap V(U) = \text{Assh } A$, and $A_{\mathfrak{p}}$ is Gorenstein for every $\mathfrak{p} \in \text{Supp}_A K_A$ with $\text{ht}_A \mathfrak{p} \leq 1$.

Proof. (1) \Rightarrow (2) By Lemma 3.1, we have $\text{Ass } A \cap V(U) = \text{Assh } A$. Let $\mathfrak{p} \in \text{Supp}_A K_A$ with $\text{ht}_A \mathfrak{p} \leq 1$. If $\mathfrak{p} \in \text{Assh } A$, then $A_{\mathfrak{p}}$ is Gorenstein by Proposition 2.3. Otherwise, we have $\dim A_{\mathfrak{p}} = 1$. Since $K_{A_{\mathfrak{p}}}$ is a reflexive $A_{\mathfrak{p}}$ -module, the ring $A_{\mathfrak{p}}$ is Gorenstein by Proposition 3.3.

(2) \Rightarrow (1) Since $A_{\mathfrak{p}}$ is Gorenstein for every $\mathfrak{p} \in \text{Assh } A$, by Proposition 2.3, K_A is torsionless. Hence we have the exact sequence

$$0 \rightarrow K_A \xrightarrow{\varphi} K_A^{**} \rightarrow C \rightarrow 0$$

of A -modules, where φ is the canonical homomorphism. Suppose that $C \neq (0)$. Let $\mathfrak{p} \in \text{Ass}_A C$. Note that $\mathfrak{p} \in \text{Supp}_A K_A$ and $[K_A]_{\mathfrak{p}} \cong K_{A_{\mathfrak{p}}}$. If $\text{ht}_A \mathfrak{p} \leq 1$, then by assumption $A_{\mathfrak{p}}$ is Gorenstein. Then $C_{\mathfrak{p}} = (0)$, which is a contradiction. Thus, $\text{ht}_A \mathfrak{p} \geq 2$. Since $\text{Ass } A \cap V(U) = \text{Assh } A$, we have $\text{depth } A_{\mathfrak{p}} \geq 1$. This shows, by Lemma 3.5, that $A_{\mathfrak{p}}$ is a Cohen-Macaulay ring with $\dim A_{\mathfrak{p}} = 1$, which is a contradiction. Therefore $C = (0)$ and K_A is a reflexive A -module. \square

We summarize some consequences of Theorem 3.6. Note that A satisfies (S_1) if and only if $\text{Ass } A = \text{Min } A$, and the latter condition implies $\text{Ass } A \cap V(U) = \text{Assh } A$.

Corollary 3.7. *If A satisfies (S_1) and (G_1) , then K_A is a reflexive A -module.*

Recall that if $\text{Assh } A = \text{Ass } A$, then $\text{Spec } A = \text{Supp}_A K_A$. Thus, we obtain the following as another direct consequence of Theorem 3.6.

Corollary 3.8. *Suppose that $\text{Assh } A = \text{Ass } A$. Then K_A is a reflexive A -module if and only if A satisfies (G_1) .*

If A is a Cohen-Macaulay ring, the above corollary recovers [14, Korollar 7.29]. Recall that A is a *generalized Cohen-Macaulay ring*, if the i^{th} local cohomology module $H_{\mathfrak{m}}^i(A)$ is a finitely generated A -module for every $i \neq d$.

Corollary 3.9. *Suppose that A is a generalized Cohen-Macaulay ring and $d > 0$. Then K_A is a reflexive A -module if and only if $\text{depth } A > 0$ and A satisfies (G_1) .*

Proof. By assumption, we have $\text{Ass } A \setminus \{\mathfrak{m}\} \subseteq \text{Assh } A$. If K_A is reflexive, then by Lemma 3.1 we see that $\text{depth } A > 0$. Without loss of generality, we may assume $\text{depth } A > 0$. Hence $\text{Ass } A = \text{Assh } A$. The assertion follows from Corollary 3.8. \square

It seems natural to ask for the relation between the reflexivity of the A -module K_A and that of the A/U -module $K_{A/U}$. As for this question, we have the following.

Theorem 3.10. *The following conditions are equivalent:*

- (1) K_A is a reflexive A -module;
- (2) $K_{A/U}$ is a reflexive A/U -module and $\text{Ass } A \cap V(U) = \text{Assh } A$.

Proof. Let $B = A/U$. Then $K_A = K_B$ ([2, 1.8]). Also note that $\text{Ass } B = \text{Assh } B$.

(1) \Rightarrow (2) By Lemma 3.1, we have $\text{Ass } A \cap V(U) = \text{Assh } A$. By Corollary 3.8, it suffices to show that B satisfies (G_1) . Let $\mathfrak{P} \in \text{Spec } B$ be a prime with $\text{ht}_B \mathfrak{P} \leq 1$. We write $\mathfrak{P} = \mathfrak{p}/U$ for some $\mathfrak{p} \in V(U)$. Then $\text{ht}_A \mathfrak{p} = \text{ht}_B \mathfrak{P} \leq 1$. Moreover, $K_{A_{\mathfrak{p}}}$ is a reflexive $A_{\mathfrak{p}}$ -module. By Propositions 2.3 and 3.3, $A_{\mathfrak{p}}$ is Gorenstein. Since $U_{\mathfrak{p}} = (0) :_{A_{\mathfrak{p}}} K_{A_{\mathfrak{p}}} = (0)$, we obtain $B_{\mathfrak{P}} = A_{\mathfrak{p}}$. Thus, $B_{\mathfrak{P}}$ is a Gorenstein ring.

(2) \Rightarrow (1) Let $\mathfrak{p} \in \text{Supp}_A K_A$ with $\text{ht}_A \mathfrak{p} \leq 1$. By Theorem 3.6, it is enough to show that $A_{\mathfrak{p}}$ is Gorenstein. Let $\mathfrak{P} = \mathfrak{p}/U$. Then by Corollary 3.8, $B_{\mathfrak{P}}$ is a Gorenstein ring. Since $\text{Ass } A \cap V(U) = \text{Assh } A$, the ring $A_{\mathfrak{p}}$ is Cohen-Macaulay. In particular, $U_{\mathfrak{p}} = (0)$ and $A_{\mathfrak{p}} = B_{\mathfrak{P}}$. Therefore $A_{\mathfrak{p}}$ is Gorenstein. \square

Corollary 3.11. *Suppose that A/U is a Gorenstein ring. Then the following assertions hold true.*

- (1) K_A is a reflexive A -module if and only if $\text{Ass } A \cap V(U) = \text{Assh } A$.
- (2) If A satisfies (S_1) , then K_A is reflexive.

Proof. Note that (1) follows directly from Theorem 3.10. To prove (2), it is enough to show that $\text{Ass } A \cap V(U) \subseteq \text{Assh } A$. Let $\mathfrak{p} \in \text{Ass } A \cap V(U)$. Since A satisfies (S_1) , we have $\text{ht}_A \mathfrak{p} = 0$. Since $\mathfrak{p} \in V(U)$, we have $\dim A = \dim A/\mathfrak{p} + \text{ht}_A \mathfrak{p} = \dim A/\mathfrak{p}$. Therefore $\mathfrak{p} \in \text{Assh } A$. \square

Closing this section, we provide the examples of (not necessarily Cohen-Macaulay) local rings admitting reflexive canonical modules.

Example 3.12. Let $S = k[[X, Y_1, Y_2, \dots, Y_n]]$ ($n \geq 2$) be the formal power series ring over a field k and let $A = S/[(X^m) \cap J]$ where $m \geq 1$ and J is a (Y_1, Y_2, \dots, Y_n) -primary ideal of S . Let x denote the image of X in A . Then $U = (x^m)$, $\text{Assh } A = \{(x)\}$, and A/U is a Gorenstein ring. By Corollary 3.11, the A -module K_A is reflexive.

Example 3.13. Let k be a field and $R = k[\Delta]$ be the Stanley-Reisner ring of a simplicial complex Δ . Since R is reduced, the graded canonical module K_R (see [13, 20]) is torsionless. Moreover, if $\#(\text{Assh } R) = 1$, the ring R/U is Gorenstein, so that K_R is reflexive as an R -module, where U stands for the unmixed component of (0) in R and $\text{Assh } R = \{\mathfrak{p} \in \text{Spec } R \mid \dim R/\mathfrak{p} = \dim R\}$.

4. QUASI-NORMAL RINGS

Quasi-normal rings were introduced by Vasconcelos ([21, Definition 1.2]) and they are exactly 2-Gorenstein rings.

Definition 4.1. A Noetherian ring R is said to be *quasi-normal* if R satisfies (S_2) and (G_1) .

The following is a direct consequence of [11, Proposition 2.3] (for a local ring case, see also [8, Theorem 3.8]). Here we include our alternative proof specifically for quasi-normal rings.

Proposition 4.2. *Let R be a quasi-normal ring and let M be a finitely generated R -module. If $\text{depth}_{R_{\mathfrak{p}}} M_{\mathfrak{p}} \geq \min\{2, \dim R_{\mathfrak{p}}\}$ for every $\mathfrak{p} \in \text{Spec } R$, then M is reflexive.*

Proof. Consider the exact sequence of R -modules

$$0 \rightarrow X \rightarrow M \xrightarrow{\varphi} M^{**} \rightarrow C \rightarrow 0,$$

where φ denotes the canonical homomorphism. Suppose $X \neq (0)$ and choose $\mathfrak{p} \in \text{Ass}_R X$. By assumption, we have $\dim R_{\mathfrak{p}} = 0$. Since R satisfies (G_1) , the local ring $R_{\mathfrak{p}}$ is Gorenstein, whence $M_{\mathfrak{p}}$ is reflexive. Hence $X_{\mathfrak{p}} = (0)$, which is a contradiction. So $X = (0)$, and we have the exact sequence

$$0 \rightarrow M \rightarrow M^{**} \rightarrow C \rightarrow 0.$$

Suppose $C \neq (0)$. Let $\mathfrak{p} \in \text{Ass}_R C$. If $\dim R_{\mathfrak{p}} = 0$, then $M_{\mathfrak{p}}$ is reflexive, so $C_{\mathfrak{p}} = (0)$. This is a contradiction. Thus $\dim R_{\mathfrak{p}} \geq 1$. As R satisfies (S_2) , we have $\text{depth}_{R_{\mathfrak{p}}} M_{\mathfrak{p}} \geq \min\{2, \dim R_{\mathfrak{p}}\}$. Hence $\text{depth}_{R_{\mathfrak{p}}} M_{\mathfrak{p}} \geq 1$. Since $\text{depth}_{R_{\mathfrak{p}}} M_{\mathfrak{p}}^{**} \geq \min\{2, \text{depth}_{R_{\mathfrak{p}}} M_{\mathfrak{p}}\}$, we then have $\text{depth}_{R_{\mathfrak{p}}} M_{\mathfrak{p}}^{**} \geq 1$. The exact sequence

$$0 \rightarrow M_{\mathfrak{p}} \rightarrow M_{\mathfrak{p}}^{**} \rightarrow C_{\mathfrak{p}} \rightarrow 0$$

gives that

$$0 = \text{depth}_{R_{\mathfrak{p}}} C_{\mathfrak{p}} \geq \min\{\text{depth}_{R_{\mathfrak{p}}} M_{\mathfrak{p}} - 1, \text{depth}_{R_{\mathfrak{p}}} M_{\mathfrak{p}}^{**}\}.$$

Hence $\text{depth}_{R_{\mathfrak{p}}} M_{\mathfrak{p}} \leq 1$. By assumption, we have

$$1 \geq \text{depth}_{R_{\mathfrak{p}}} M_{\mathfrak{p}} \geq \min\{2, \dim R_{\mathfrak{p}}\}.$$

Therefore $\dim R_{\mathfrak{p}} = 1$ and $\text{depth}_{R_{\mathfrak{p}}} M_{\mathfrak{p}} = 1$. Since R satisfies (G_1) , the ring $R_{\mathfrak{p}}$ is Gorenstein. By [21, Corollary 2.3], we see that $M_{\mathfrak{p}}$ is reflexive. Hence $C_{\mathfrak{p}} = (0)$, which is a contradiction. \square

A finitely generated R -module ω_R is a *canonical module* of R , if $(\omega_R)_{\mathfrak{m}}$ is the canonical module of $R_{\mathfrak{m}}$ for all maximal ideals \mathfrak{m} of R . In contrast to the local case, the canonical module is in general not unique up to isomorphisms; see e.g., [6, Remark 3.3.17].

Corollary 4.3. *Let R be a Noetherian ring with $d = \dim R > 0$. Suppose that there exists a canonical module ω_R and $\text{Ass } R_{\mathfrak{m}} = \text{Assh } R_{\mathfrak{m}}$ for every maximal ideal \mathfrak{m} . Then R is quasi-normal if and only if R satisfies (S_2) and ω_R is reflexive.*

Proof. Suppose R is quasi-normal. Since ω_R satisfies (S_2) and $\dim_{R_{\mathfrak{p}}}[\omega_R]_{\mathfrak{p}} = \dim R_{\mathfrak{p}}$ for every $\mathfrak{p} \in \text{Spec } R$, by Proposition 4.2, we conclude that ω_R is reflexive. For the converse, it remains to show that R satisfies (G_1) . Let $A = R_{\mathfrak{m}}$, where \mathfrak{m} is a maximal ideal of R . Then $K_A = (\omega_R)_{\mathfrak{m}}$ is reflexive. Therefore we have $\text{Ass } A = \text{Assh } A$. By Corollary 3.8, A satisfies (G_1) . Thus R satisfies (G_1) . \square

We summarize some examples. First, we note examples of quasi-normal rings which are not normal. The simplest ones are non-normal Gorenstein rings.

Example 4.4. Suppose that R is a Cohen-Macaulay ring with canonical module ω_R . We set $T = R \ltimes \omega_R$ to be the idealization of ω_R over R . Then T is a Gorenstein ring ([16]), but not normal because it is never a reduced ring.

For a commutative ring R , we denote by \overline{R} the integral closure of R in $\mathbb{Q}(R)$. We refer to [6, p. 178] for background on numerical semigroups.

Example 4.5. Let $H = \langle a_1, a_2, \dots, a_\ell \rangle$ be a symmetric numerical semigroup. We consider $R = k[s, t^{a_1}, t^{a_2}, \dots, t^{a_\ell}]$, where s, t are indeterminates and k is a field. Then R is a two-dimensional Gorenstein ring with $\overline{R} = k[s, t]$, so that R is not normal if $1 \notin H$. As a special case, the ring $R = k[s, t^2, t^3]$ is quasi-normal, but not normal.

Next, we note examples of quasi-normal but non-normal Cohen-Macaulay rings which are moreover not Gorenstein.

Example 4.6. Let k be a field and $k[X, Y]$ the polynomial ring over k . Let $H = \langle a_1, a_2, \dots, a_\ell \rangle$ be a symmetric numerical semigroup such that $1 \notin H$ and let $k[H] = k[t^{a_1}, t^{a_2}, \dots, t^{a_\ell}]$ denote the semigroup ring of H over k , where t is an indeterminate. Let $T = k[X^n, X^{n-1}Y, \dots, XY^{n-1}, Y^n]$, where $n \geq 3$ is an integer. We set $R = T \otimes_k k[H]$. Then R is a quasi-normal Cohen-Macaulay ring with $\dim R = 3$, which is neither Gorenstein nor normal. Indeed, because $\overline{R} = T \otimes_k k[t]$ and $k[H] \neq k[t]$, the ring R is not normal. As T is normal, we see that R is a quasi-normal ring (see Proposition 7.3 (2)). Moreover, R is not a Gorenstein ring because T is not Gorenstein. The simplest example in this class is $R = k[X^3, X^2Y, XY^2, Y^3] \otimes_k k[t^2, t^3]$.

Example 4.7. Let $T = k[X, Y, Z, V]$ be the polynomial ring over a field k . We denote by $\mathbf{I}_2(\mathbb{N})$ the ideal of T generated by all the 2×2 minors of a matrix \mathbb{N} . Let $I = \mathbf{I}_2(\mathbb{M})$ where $\mathbb{M} = \begin{pmatrix} X^a & Y^b + V & Z^c \\ Y^{b'} & Z^{c'} & X^{a'} \end{pmatrix}$ for some integers $a, b, c, a', b', c' \geq 1$. We set $R = T/I$. Then R is a Cohen-Macaulay ring of dimension 2. Let x, y, z, v denote the images of X, Y, Z, V in R , respectively. We first check the isomorphism $\omega_R \cong (x^a, y^{b'})R$. In fact, by setting

$$f = Z^{c+c'} - X^{a'}(Y^b + V), \quad g = X^{a+a'} - Y^{b'}Z^c, \quad \text{and} \quad h = -X^aZ^{c'} + Y^{b'}(Y^b + V),$$

we can consider the exact sequence

$$0 \longrightarrow T^2 \xrightarrow{t_{\mathbb{M}}} T^3 \xrightarrow{\begin{pmatrix} f & g & h \end{pmatrix}} T \longrightarrow R \longrightarrow 0$$

of T -modules. By taking the T -dual, we get the presentation of ω_R of the form $T^3 \xrightarrow{\mathbb{M}} T^2 \rightarrow \omega_R \rightarrow 0$. Therefore, the complex of R -modules

$$R^3 \xrightarrow{\mathbb{M}} R^2 \xrightarrow{\begin{pmatrix} Y^{b'} & -X^a \end{pmatrix}} (x^a, y^{b'})R \longrightarrow 0$$

induces a natural epimorphism

$$\varphi : \omega_R \twoheadrightarrow (x^a, y^{b'})R$$

of R -modules. Moreover, φ is an isomorphism because ω_R is a torsionfree R -module of rank one and x^a is a non-zero-divisor on R . Hence $\omega_R \cong (x^a, y^{b'})R$, as claimed. We similarly have

$$\omega_R \cong (y^b + v, z^{c'})R \cong (z^c, x^{a'})R.$$

We also note that the isomorphisms can be obtained by using the procedure of [22, Section 6.1.2]. In particular, R is not a Gorenstein ring, since the type of R is two.

Next, we show that R is a quasi-normal ring. Let $\mathfrak{p} \in \text{Spec } R$ with $\text{ht}_R \mathfrak{p} \leq 1$. If $x \notin \mathfrak{p}$, then $[\omega_R]_{\mathfrak{p}} \cong (x^a, y^{b'})R_{\mathfrak{p}} = R_{\mathfrak{p}}$, so that $R_{\mathfrak{p}}$ is a Gorenstein ring. Assume that $x \in \mathfrak{p}$. Similarly, we may assume that $y, z \in \mathfrak{p}$. Then, $v \notin \mathfrak{p}$, since $\text{ht}_R \mathfrak{p} \leq 1$. Therefore, $[\omega_R]_{\mathfrak{p}} \cong (y^b + v, z^{c'})R_{\mathfrak{p}} = R_{\mathfrak{p}}$, so that $R_{\mathfrak{p}}$ is a Gorenstein ring. Hence R is a quasi-normal ring.

Finally, we prove that R is a normal ring if and only if $a' = b' = c = 1$. Assume $a' = b' = c = 1$ and consider the ideal

$$J = \mathbf{I}_2 \begin{pmatrix} Z^{c'} & (a+1)X^a & Y^b + V \\ -(b+1)Y^b + V & -Z & bXY^{b-1} \\ c'X^aZ^{c'-1} & -Y & -(c'+1)Z^{c'} \\ -Y & 0 & X \end{pmatrix}.$$

Then $J + I/I$ is the Jacobian ideal of R over k . A direct computation shows that $\sqrt{J+I} = (X, Y, Z, V)$ and hence, by the Jacobian criterion, the local ring $R_{\mathfrak{p}}$ is regular for every $\mathfrak{p} \in \text{Spec } R \setminus \{(x, y, z, v)\}$. Hence R is a normal ring. Conversely, we assume $a' \geq 2$, or $b' \geq 2$, or $c \geq 2$. By taking

$$P = \begin{cases} (X, Y^b + V, Z) & (\text{if } c \geq 2) \\ (X, Y, Z) & (\text{if } c = 1), \end{cases}$$

we then have $J \subseteq P$. We set $\mathfrak{p} = PR$. Then $\text{ht}_R \mathfrak{p} = 1$, but $R_{\mathfrak{p}}$ is not a DVR. Indeed, because $\varepsilon = Y$ or $\varepsilon' = Y^b + V$ is invertible in T_P , we see that

$$JT_P = \begin{cases} \left(Z^{c+c'} - X^{a'}(Y^b + V), \frac{X^{a+a'}}{\varepsilon^{b'}} - Z^c, -\frac{X^a Z^{c'}}{\varepsilon} + (Y^b + V) \right) \subseteq (Y^b + V) + (X, Z)^2 & (\text{if } c \geq 2) \\ \left(\frac{Z^{c+c'}}{\varepsilon'} - X^{a'}, X^{a+a'} - Y^{b'}Z^c, -\frac{X^a Z^{c'}}{\varepsilon'} + Y^{b'} \right) \subseteq (Y) + (X, Z)^2 & (\text{if } c = 1) \end{cases}$$

in T_P . Thus $R_{\mathfrak{p}} = T_P/JT_P$ cannot be a DVR. Hence R is not a normal ring. As a special case, $R = k[X, Y, Z, V]/\mathbf{I}_2 \begin{pmatrix} X & Y+V & Z \\ Y & Z & X^2 \end{pmatrix}$ is a quasi-normal ring, but not normal.

5. REFLEXIVE CANONICAL MODULES IN DIMENSION ONE

Let (A, \mathfrak{m}) be a Cohen-Macaulay local ring with $\dim A = 1$ admitting the canonical module K_A . In this section, we explore the question of when A has a reflexive canonical module. We denote by $\mathbb{Q}(A)$ the total ring of fractions of A . Throughout this section, we assume that there exists an A -submodule K of $\mathbb{Q}(A)$ such that $A \subseteq K \subseteq \overline{A}$ and $K \cong K_A$ as an A -module, where \overline{A} denotes the integral closure of A in $\mathbb{Q}(A)$. Note that the assumption is automatically satisfied if $\mathbb{Q}(A)$ is Gorenstein and the residue class field A/\mathfrak{m} is infinite; see [12, Corollaries 2.8, 2.9]. For A -submodules X and Y of $\mathbb{Q}(A)$, let $X : Y = \{a \in \mathbb{Q}(A) \mid aY \subseteq X\}$. If we consider ideals I, J of A , we set $I :_A J = \{a \in A \mid aJ \subseteq I\}$; hence $I :_A J = (I : J) \cap A$.

Proposition 5.1. *The following conditions are equivalent:*

- (1) A is a Gorenstein ring;
- (2) $K^2 : K = K$;
- (3) K_A is a reflexive A -module.

Proof. (1) \Leftrightarrow (3) See Proposition 3.3.

(3) \Leftrightarrow (2) Since $A : K = [K : K] : K = K : K^2$ ([14, Bemerkung 2.5]), we have

$$A : (A : K) = (K : K) : (K : K^2) = [K : (K : K^2)] : K = K^2 : K.$$

Therefore, $K^2 : K = K$ if and only if $A : (A : K) = K$, that is K_A is a reflexive A -module. \square

Recall that an ideal I of A is called a *canonical ideal* of A , if $I \neq A$ and $I \cong K_A$ as an A -module. By [12, Corollary 2.8], there exists a canonical ideal I of A . We then have the following.

Theorem 5.2. *Let I be a canonical ideal of A . Then the following conditions are equivalent:*

- (1) A is a Gorenstein ring;
- (2) $I^2 :_A I = I$;
- (3) I/I^2 is a free A/I -module;
- (4) I is a reflexive A -module.

Proof. By Proposition 5.1, it suffices to show (2) \Rightarrow (1). Enlarging the residue class field A/\mathfrak{m} of A if necessary, we may assume that A/\mathfrak{m} is infinite. Let $I = (x_1, x_2, \dots, x_n)$ ($n > 0$) so that each (x_i) is a reduction of I . We set $K_i = x_i^{-1}I$ and choose a non-zero-divisor b of A so that $bK_i^2 \subseteq A$ for all $1 \leq i \leq n$. Let $J = bI$ and $y_i = bx_i$ for $1 \leq i \leq n$. Then (y_i) is a reduction of J . Notice that A/I and A/J are both Gorenstein rings, since $I, J \cong K_A$ as A -modules.

Claim 1. $J^2 :_A J = J$.

Proof of Claim 1. Suppose that $J^2 :_A J \supsetneq J$. Then, $J :_A \mathfrak{m} \subseteq J^2 :_A J$. Since A/J is a Gorenstein ring, we have $J :_A \mathfrak{m} = J + A\varphi$ for some $\varphi \in (J :_A \mathfrak{m}) \setminus J$. Hence, $\frac{\varphi}{b} \in \mathbb{Q}(A)$ and $\mathfrak{m} \cdot \frac{\varphi}{b} \subseteq I$, so that $\frac{\varphi}{b} \in I : \mathfrak{m}$. Because $I \subsetneq I :_A \mathfrak{m} \subseteq I : \mathfrak{m}$ and $\ell_A((I : \mathfrak{m})/I) = 1$ (since A/I is a Gorenstein ring), we get $\frac{\varphi}{b} \in I :_A \mathfrak{m}$, so that $\frac{\varphi}{b} \in A$. On the other hand, $\frac{\varphi}{b} \cdot I \subseteq I^2$, since $\varphi \cdot bI = \varphi J \subseteq J^2 = b^2 I^2$. Consequently, $\frac{\varphi}{b} \in I^2 :_A I = I$, whence $\varphi \in bI = J$, which is impossible. Thus $J^2 :_A J = J$. \square

Let \bar{y}_i denote the image of y_i in J/J^2 . We then have $J/J^2 = \sum_{i=1}^n (A/J) \cdot \bar{y}_i$, and therefore, $(0) :_{A/J} \bar{y}_i = (0)$ for some i , since A/J is a Gorenstein ring and $(0) :_{A/J} J/J^2 = (0)$ by Claim 1. Without loss of generality, assume $i = 1$. Then, $J^2 :_A y_1 = J$. On the other hand, since $bK_1^2 \subseteq A$ and $K_1 = y_1^{-1}J$, we get $b \cdot (y_1^{-1}J)^2 \subseteq A$, whence $bJ^2 \subseteq (bx_1)^2$. Therefore, $J^2 \subseteq (bx_1^2) \subseteq (bx_1) = (y_1)$. Hence, $J^2 = y_1 \cdot (J^2 :_A y_1) = y_1 J$. Thus A is a Gorenstein ring (see [12, Theorem 3.7]). \square

6. q -TORSIONFREE CANONICAL MODULES

The purpose of this section is to give a generalization of Proposition 2.3 and Theorem 3.6, which characterize local rings with q -torsionfree canonical modules for $q = 1, 2$.

Let R be a Noetherian (not necessarily local) ring and q an integer. Let M be a finitely generated R -module with a finite projective presentation $P_1 \xrightarrow{\sigma} P_0 \rightarrow M \rightarrow 0$. By applying the R -dual functor $(-)^* = \text{Hom}_R(-, R)$, we obtain the exact sequence

$$0 \longrightarrow M^* \longrightarrow P_0^* \xrightarrow{\sigma^*} P_1^* \longrightarrow D_\sigma(M) \longrightarrow 0$$

of R -modules. We set $D(M) = D_\sigma(M)$ and call it the *Auslander transpose* of M . Note that $D(M)$ is uniquely determined up to projective equivalence.

Definition 6.1 ([1, Definition 2.15]). A finitely generated R -module M is said to be *q -torsionfree* if $\text{Ext}_R^i(D(M), R) = 0$ for all $i = 1, 2, \dots, q$.

By [1, Proposition 2.6], there exists an exact sequence

$$0 \longrightarrow \text{Ext}_R^1(D(M), R) \longrightarrow M \xrightarrow{\varphi} M^{**} \longrightarrow \text{Ext}_R^2(D(M), R) \longrightarrow 0$$

of R -modules, where φ is the canonical homomorphism, and furthermore, we have

$$\text{Ext}_R^{i+2}(D(M), R) \cong \text{Ext}_R^i(M^*, R) \quad \text{for all } i > 0.$$

This shows M is torsionless (resp. reflexive) if and only if M is 1-torsionfree (2-torsionfree). When $q \geq 3$, the R -module M is q -torsionfree if and only if M is reflexive and $\text{Ext}_R^i(M^*, R) = (0)$ for all $i = 1, 2, \dots, q - 2$.

Example 6.2. Let $S = k[[X, Y, Z]]$ be the formal power series ring over a field k and set $A = S/[(X) \cap (Y, Z)]$. Let x, y, z denote the images of X, Y, Z in A , respectively. Then we have $\text{Assh } A = \{(x)\}$, $U = (x)$, and $K_A = A/U$, where U denotes the unmixed component of (0) in A . By dualizing the exact sequence

$$A \xrightarrow{\cdot x} A \rightarrow A/(x) = K_A \rightarrow 0,$$

we obtain

$$0 \rightarrow K_A^* \rightarrow A \xrightarrow{\cdot x} A \rightarrow A/(x) \rightarrow 0.$$

Thus, $D(K_A) = K_A$. Consider the free resolution of $D(K_A) = D$:

$$A^5 \xrightarrow{\tau_4} A^3 \xrightarrow{\tau_3} A^2 \xrightarrow{\tau_2} A \xrightarrow{\cdot x} A \longrightarrow D \longrightarrow 0,$$

where $\tau_2 = [y \ z]$, $\tau_3 = \begin{bmatrix} x & 0 & z \\ 0 & x & -y \end{bmatrix}$, and $\tau_4 = \begin{bmatrix} y & z & 0 & 0 & 0 \\ 0 & 0 & y & z & 0 \\ 0 & 0 & 0 & 0 & x \end{bmatrix}$. Dualize this free resolution to obtain

$$0 \longrightarrow D^* \longrightarrow A \xrightarrow{\cdot x} A \xrightarrow{\sigma_2} A^2 \xrightarrow{\sigma_3} A^3 \xrightarrow{\sigma_4} A^5,$$

where $\sigma_2, \sigma_3, \sigma_4$ are the transposes of τ_2, τ_3, τ_4 , respectively. Let $a \in \text{Ker } \sigma_2$. Then

$$a \in [(0) :_A y] \cap [(0) :_A z] = (x) \cap (x) = (x) = \text{Im}(\cdot x).$$

Thus, $\text{Ext}_A^1(D, A) = (0)$. Let $\begin{pmatrix} a_1 \\ a_2 \end{pmatrix} \in \text{Ker } \sigma_3$. Then $a_1, a_2 \in (0) :_A x = (y, z)$ and $a_1 z - a_2 y = 0$. Hence $\begin{pmatrix} -a_2 \\ a_1 \end{pmatrix} \in \text{Ker } \tau_2 = \text{Im}(\tau_3)$. Let $-a_2 = c_1 x - c_3 z$, and $a_1 = c_2 x + c_3 y$ for some $c_1, c_2, c_3 \in A$. Then $c_1 x = -a_2 + c_3 z \in (y, z)$. Thus, $c_1 = 0$ and $a_2 = c_3 z$. Similarly, $a_1 = c_3 y$. We obtain $\begin{pmatrix} a_1 \\ a_2 \end{pmatrix} \in \text{Im } \sigma_2$. Then $\text{Ext}_A^2(D, A) = (0)$. Hence K_A is 2-torsionfree. Note that $\text{Ker } \sigma_4$ is generated by $\begin{pmatrix} x \\ 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ x \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ 0 \\ y \end{pmatrix}, \begin{pmatrix} 0 \\ 0 \\ z \end{pmatrix}$. Then $\text{Ext}_A^3(D, A) \neq (0)$. Thus, K_A is not 3-torsionfree.

Definition 6.3 ([1, Definition 2.15]). A finitely generated R -module M is called q -syzygy, if there exist finite free R -modules F_1, F_2, \dots, F_q and an exact sequence $0 \rightarrow M \rightarrow F_1 \rightarrow F_2 \rightarrow \dots \rightarrow F_q$ of R -modules.

Note that (a) M is torsionless if and only if M is 1-syzygy, (b) every q -torsionfree R -module is q -syzygy, and (c) if M is q -syzygy and x is an R -regular element, then M/xM is $(q-1)$ -syzygy as an R/xR -module.

Although the following theorem has been proved by Foxby in a more general setting involving Gorenstein modules, we restate it and give its proof in our context for the sake of completeness. Recall that R is q -Gorenstein if $R_{\mathfrak{p}}$ is Gorenstein for every prime \mathfrak{p} with $\text{depth } R_{\mathfrak{p}} < q$.

Theorem 6.4 ([10, Proposition 3.2]). *Let A be a Cohen-Macaulay local ring admitting the canonical module K_A . Then the following conditions are equivalent:*

- (1) A is q -Gorenstein;
- (2) K_A is q -torsionfree;
- (3) K_A is q -syzygy.

Proof. Since A is Cohen-Macaulay, we have $\text{Spec } A = \text{Supp}_A K_A$ and $[K_A]_{\mathfrak{p}} = K_{A_{\mathfrak{p}}}$ for every $\mathfrak{p} \in \text{Spec } A$. Notice that $K_{A_{\mathfrak{p}}}$ is maximal Cohen-Macaulay as an $A_{\mathfrak{p}}$ -module.

(1) \Rightarrow (2) Since every A -regular sequence is K_A -regular, the A -module K_A is q -torsionfree by [11, Proposition 2.3].

(2) \Rightarrow (3) This follows from [11, Proposition 2.1].

(3) \Rightarrow (1) Let $\mathfrak{p} \in \text{Spec } A$ with $\text{depth } A_{\mathfrak{p}} < q$. Set $n = \text{dim } A_{\mathfrak{p}}$. When $n = 0$, the ring $A_{\mathfrak{p}}$ is Gorenstein. Assume $n > 0$ and choose a system f_1, f_2, \dots, f_n of parameters of $A_{\mathfrak{p}}$. Then it is an $A_{\mathfrak{p}}$ -regular sequence, so that $K_{A_{\mathfrak{p}}}/(f_1, f_2, \dots, f_n)K_{A_{\mathfrak{p}}}$ is 1-syzygy because $n < q$. Since $K_{A_{\mathfrak{p}}}/(f_1, f_2, \dots, f_n)K_{A_{\mathfrak{p}}} \cong K_{A_{\mathfrak{p}}}/(f_1, f_2, \dots, f_n)A_{\mathfrak{p}}$, we conclude that $A_{\mathfrak{p}}/(f_1, f_2, \dots, f_n)A_{\mathfrak{p}}$ is Gorenstein by Proposition 2.3, whence so is the ring $A_{\mathfrak{p}}$. This completes the proof. \square

As a direct consequence of Theorem 6.4, we have the following.

Corollary 6.5. *Let A be a Cohen-Macaulay local ring with $d = \dim A$ admitting the canonical module K_A . Then A is a Gorenstein ring if and only if K_A is $(d + 1)$ -torsionfree.*

Theorem 6.4 and Corollary 6.5 lead us to the question of the structure of a local ring A with q -torsionfree canonical module without the assumption that A is Cohen-Macaulay.

For a subset Φ of prime ideals in a Noetherian ring R , we say that

- R satisfies (S_n) on Φ if $\text{depth } R_{\mathfrak{p}} \geq \min\{n, \dim R_{\mathfrak{p}}\}$ for every $\mathfrak{p} \in \Phi$,
- R satisfies (G_n) on Φ if $R_{\mathfrak{p}}$ is Gorenstein for every $\mathfrak{p} \in \Phi$ with $\dim R_{\mathfrak{p}} \leq n$.

The main result of this section gives an answer to the above question.

Theorem 6.6. *Let A be a Noetherian local ring admitting the canonical module K_A . Suppose that K_A satisfies (S_q) . Then the following conditions are equivalent:*

- (1) A satisfies both (G_{q-1}) and (S_{q-1}) on $\text{Supp}_A K_A$;
- (2) K_A is q -torsionfree;
- (3) K_A is q -syzygy.

To show this, we need some auxiliaries. The following plays an important role in our argument.

Lemma 6.7 ([7, Lemma 4.9]). *Let A be a Noetherian local ring and M a nonzero finitely generated A -module. Assume that $q \geq \text{depth } A + 2$ and M is q -syzygy. Then $\text{depth}_A M = \text{depth } A$.*

We apply Lemma 6.7 to get the following.

Theorem 6.8. *Let R be a Noetherian ring and M a finitely generated R -module. If M is $(q + 1)$ -syzygy, then one has*

$$\text{depth } R_{\mathfrak{p}} \geq \min\{q, \text{depth}_{R_{\mathfrak{p}}} M_{\mathfrak{p}}\} \quad \text{for all } \mathfrak{p} \in \text{Supp}_R M.$$

In particular, if M satisfies (S_q) , then R satisfies (S_q) on $\text{Supp}_R M$.

Proof. By localizing at $\mathfrak{p} \in \text{Supp}_R M$, it suffices to show $\text{depth } R \geq \min\{q, \text{depth}_R M\}$. If $\text{depth } R \geq q$, the assertion is obvious. Otherwise, if $\text{depth } R < q$, the assertion follows from Lemma 6.7. \square

As consequences of Theorems 6.4, 6.8, we get the following.

Corollary 6.9. *Let A be a Noetherian local ring with $d = \dim A$ admitting the canonical module K_A . Then the following conditions are equivalent:*

- (1) A is Gorenstein;
- (2) K_A is a $(d + 1)$ -torsionfree maximal Cohen-Macaulay A -module;
- (3) K_A is a $(d + 1)$ -syzygy maximal Cohen-Macaulay A -module.

Proof. We only need to show (3) \Rightarrow (1). By Theorem 6.8 we have that $\text{depth } A = d$, so that A is Cohen-Macaulay. Hence the assertion follows from Theorem 6.4. \square

Corollary 6.10. *Let (A, \mathfrak{m}) be a Noetherian local ring with $d = \dim A$ admitting the canonical module K_A . Furthermore, we assume one of the following conditions (i) and (ii).*

- (i) $H_{\mathfrak{m}}^i(A) = (0)$ for every integer $i \neq 0, 1, d$.

(ii) $d \leq 2$.

Then the following conditions are equivalent:

- (1) A is Gorenstein;
- (2) K_A is $(d+1)$ -torsionfree;
- (3) K_A is $(d+1)$ -syzygy.

Proof. (i) By passing to the \mathfrak{m} -adic completion, we may assume A is \mathfrak{m} -adically complete. In view of [17, (2.3) Satz], it follows that K_A is maximal Cohen-Macaulay. Therefore the assertion follows from Corollary 6.9.

(ii) Since K_A satisfies (S_2) , the assertion follows from Corollary 6.9. \square

Remark 6.11. Let A be a Noetherian local ring admitting the canonical module K_A . We say that A is *quasi-Gorenstein* if $K_A \cong A$ as an A -module. When $d \geq 3$, there exist non-Gorenstein quasi-Gorenstein local rings of dimension d (see e.g., [2, Theorem 2.11]). Notice that, in such a ring A , K_A is q -torsionfree for all $q \geq 1$. So, Corollary 6.10 fails without the condition (i) or (ii).

Based on the above observation, it is natural to raise the following question.

Question 6.12. Let A be a Noetherian local ring with $d = \dim A \geq 3$ admitting the canonical module K_A . When are the following conditions equivalent?

- (i) A is a quasi-Gorenstein ring, i.e., $K_A \cong A$.
- (ii) K_A is a $(d+1)$ -torsionfree A -module.

In what follows, let R be a Noetherian ring and M a finitely generated R -module. The equivalence of (1) and (2) in the next theorem was essentially proved by Auslander and Bridger [1]. Notice that this is a q^{th} version of [6, Proposition 1.4.1].

Theorem 6.13. *The following conditions are equivalent:*

- (1) M is q -torsionfree;
- (2) M satisfies the conditions below:
 - (i) $M_{\mathfrak{p}}$ is q -torsionfree for every $\mathfrak{p} \in \text{Supp}_R M$ with $\text{depth } R_{\mathfrak{p}} < q$;
 - (ii) M satisfies (\widetilde{S}_q) ;
- (3) M satisfies the conditions below:
 - (i) $M_{\mathfrak{p}}$ is q -torsionfree for every $\mathfrak{p} \in \text{Supp}_R M$ with $\text{depth}_{R_{\mathfrak{p}}} M_{\mathfrak{p}} < q$;
 - (ii) $\text{depth}_{R_{\mathfrak{p}}} M_{\mathfrak{p}} = \text{depth } R_{\mathfrak{p}}$ for every $\mathfrak{p} \in \text{Supp}_R M$ with $\text{depth } R_{\mathfrak{p}} < q - 1$.

Proof. Without loss of generality, we may assume $q \geq 1$.

(1) \Rightarrow (2) This follows from [11, Proposition 2.1].

(2) \Rightarrow (3) (i) Let $\mathfrak{p} \in \text{Supp}_R M$ with $\text{depth}_{R_{\mathfrak{p}}} M_{\mathfrak{p}} < q$. Since M satisfies (\widetilde{S}_q) , we have $\text{depth}_{R_{\mathfrak{p}}} M_{\mathfrak{p}} \geq \text{depth } R_{\mathfrak{p}}$. This implies $\text{depth } R_{\mathfrak{p}} < q$, and hence $M_{\mathfrak{p}}$ is q -torsionfree. (ii) Let $\mathfrak{p} \in \text{Supp}_R M$ with $\text{depth } R_{\mathfrak{p}} < q - 1$. Then $M_{\mathfrak{p}}$ is q -torsionfree, so that $\text{depth}_{R_{\mathfrak{p}}} M_{\mathfrak{p}} = \text{depth } R_{\mathfrak{p}}$ by Lemma 6.7.

(3) \Rightarrow (1) For each $i \in \{1, 2, \dots, q\}$, we set $E^i = \text{Ext}_R^i(\text{D}(M), R)$. Suppose $E^q \neq 0$ and seek a contradiction. Take $\mathfrak{p} \in \text{Ass}_R E^q$. Since $\mathfrak{p} \in \text{Supp}_R M$, by (i) we have $\text{depth}_{R_{\mathfrak{p}}} M_{\mathfrak{p}} \geq q$. Then by

(ii), $\text{depth } R_{\mathfrak{p}} \geq q - 1$. By passing to the localization $R_{\mathfrak{p}}$ at \mathfrak{p} , we may assume R is a local ring, $\text{depth } R \geq q - 1$, $\text{depth}_R M \geq q$, and $\text{depth}_R E^q = 0$.

We proceed by induction on q . First, assume that $q = 1$. Since E^1 is isomorphic to a submodule of M , it follows that $\text{depth}_R M = 0$, a contradiction. Thus $E^1 = (0)$. Next, we assume $q = 2$. Applying the depth lemma to the exact sequence $0 \rightarrow M \rightarrow M^{**} \rightarrow E^2 \rightarrow 0$ of R -modules, we get $\text{depth}_R M = 1$, as $\text{depth}_R M^{**} \geq 1$. This is impossible, whence $E^2 = (0)$. Suppose $q \geq 3$ and the assertion holds for $q - 1$, i.e., M is $(q - 1)$ -torsionfree. Hence $E^1 = \dots = E^{q-1} = (0)$. Consider a free resolution (F_i, ∂_i) of M^* . Applying the R -dual functor $(-)^*$, we get the exact sequence

$$0 \rightarrow M^{**} \rightarrow F_0^* \xrightarrow{\partial_1^*} F_1^* \rightarrow \dots \rightarrow F_{q-3}^* \xrightarrow{\partial_{q-2}^*} F_{q-2}^*$$

of R -modules because $E^3 = \dots = E^{q-1} = (0)$. Let C be the cokernel of ∂_{q-2}^* . Since M is reflexive as an R -module, we obtain the exact sequence of the form

$$0 \rightarrow M \rightarrow F_0^* \rightarrow \dots \rightarrow F_{q-2}^* \rightarrow C \rightarrow 0.$$

Since $E^q = \text{Ext}_R^{q-2}(M^*, R)$ may be regarded as a submodule of C , we see that $\text{depth}_R C = 0$. Hence $\text{depth}_R M = q - 1$. This gives a contradiction. Hence we conclude that $E^q = (0)$, which shows M is q -torsionfree. \square

Corollary 6.14. *Suppose that the following conditions are satisfied:*

- (a) $M_{\mathfrak{p}}$ is q -torsionfree for every $\mathfrak{p} \in \text{Supp}_R M$ with $\dim R_{\mathfrak{p}} < q$;
- (b) M satisfies (S_q) ;
- (c) $\text{depth } R_{\mathfrak{p}} \geq \min\{q - 1, \dim R_{\mathfrak{p}} - 1\}$ for every $\mathfrak{p} \in \text{Supp}_R M$.

Then M is q -torsionfree.

Proof. We will check condition (3) in Theorem 6.13. Let $\mathfrak{p} \in \text{Supp}_R M$. (i) Assume that $\text{depth}_{R_{\mathfrak{p}}} M_{\mathfrak{p}} < q$. By (b), $\text{depth}_{R_{\mathfrak{p}}} M_{\mathfrak{p}} = \dim R_{\mathfrak{p}}$, so that $\dim R_{\mathfrak{p}} < q$. Therefore $M_{\mathfrak{p}}$ is q -torsionfree by (a). (ii) Assume that $\text{depth } R_{\mathfrak{p}} < q - 1$. By (c), $\text{depth } R_{\mathfrak{p}} \geq \dim R_{\mathfrak{p}} - 1$, so that $\dim R_{\mathfrak{p}} < q$. Therefore $M_{\mathfrak{p}}$ is q -torsionfree by (a), whence $\text{depth}_{R_{\mathfrak{p}}} M_{\mathfrak{p}} = \text{depth } R_{\mathfrak{p}}$ by Lemma 6.7. \square

Corollary 6.15. *Suppose that the following conditions are satisfied:*

- (a) M satisfies (S_q) ;
- (b) R satisfies both (G_{q-1}) and (S_{q-1}) on $\text{Supp}_R M$.

Then M is q -torsionfree.

Proof. By Corollary 6.14, it suffices to show that $M_{\mathfrak{p}}$ is q -torsionfree on $\{\mathfrak{p} \in \text{Supp}_R M \mid \dim R_{\mathfrak{p}} < q\}$. Let $\mathfrak{p} \in \text{Supp}_R M$ with $\dim R_{\mathfrak{p}} < q$. Then by (b), $R_{\mathfrak{p}}$ is Gorenstein. As M satisfies (S_q) , $\text{depth}_{R_{\mathfrak{p}}} M_{\mathfrak{p}} \geq \dim R_{\mathfrak{p}}$. Hence $M_{\mathfrak{p}}$ is maximal Cohen-Macaulay as an $R_{\mathfrak{p}}$ -module. In particular, $M_{\mathfrak{p}}$ is q -torsionfree. \square

We are now ready to prove Theorem 6.6.

Proof of Theorem 6.6. (1) \Rightarrow (2) This follows from Corollary 6.15.

(2) \Rightarrow (3) This follows from [11, Proposition 2.1].

(3) \Rightarrow (1) By Theorem 6.8, the ring A satisfies (S_{q-1}) on $\text{Supp}_A K_A$. Let $\mathfrak{p} \in \text{Supp}_A K_A$ with $\dim A_{\mathfrak{p}} \leq q - 1$. Then $A_{\mathfrak{p}}$ is a Cohen-Macaulay ring of dimension at most $q - 1$. Hence Theorem 6.4 implies that $A_{\mathfrak{p}}$ is a Gorenstein ring. \square

As consequences of Theorem 6.6, we get the following corollaries.

Corollary 6.16. *Let A be a Noetherian local ring admitting the canonical module K_A . Suppose that $q \geq 2$ and K_A satisfies (S_q) . Consider the following conditions:*

- (1) K_A is $(q + 1)$ -torsionfree;
- (2) K_A is $(q + 1)$ -syzygy;
- (3) A satisfies both (S_q) and (G_{q-1}) on $\text{Supp}_A K_A$;
- (4) A satisfies both (S_q) and (G_{q-1}) , that is, A is q -Gorenstein;
- (5) A satisfies (S_q) and K_A is q -torsionfree;
- (6) A satisfies (S_q) and K_A is q -syzygy.

Then the implications $(1) \Leftrightarrow (2) \Rightarrow (3) \Leftrightarrow (4) \Leftrightarrow (5) \Leftrightarrow (6)$ hold true.

Proof. The implications $(1) \Rightarrow (2)$ and $(4) \Rightarrow (3)$ are clear. The equivalence of (3), (5), and (6) immediately follows from Theorem 6.6. Thus it suffices to check the implications $(2) \Rightarrow (3)$, $(3) \Rightarrow (4)$, and $(2) \Rightarrow (1)$.

$(2) \Rightarrow (3)$ Theorem 6.6 shows the ring A satisfies (G_{q-1}) on $\text{Supp}_A K_A$. On the other hand, by Theorem 6.8, we deduce that A satisfies (S_q) on $\text{Supp}_A K_A$.

$(3) \Rightarrow (4)$ Since $q \geq 2$, by [3, Lemma 1.1] we have $\text{Supp}_A K_A = \text{Spec } A$.

$(2) \Rightarrow (1)$ The implication $(2) \Rightarrow (4)$ guarantees that A is q -Gorenstein. Hence K_A is q -torsionfree by [1, Proposition 4.21]. \square

Since K_A satisfies (S_2) , from Corollary 6.16 we have the following.

Corollary 6.17. *Let A be a Noetherian local ring admitting the canonical module K_A . Consider the following conditions:*

- (1) K_A is 3-torsionfree;
- (2) K_A is 3-syzygy;
- (3) A satisfies both (S_2) and (G_1) , that is, A is quasi-normal;
- (4) A satisfies (S_2) and K_A is 2-torsionfree;
- (5) A satisfies (S_2) and K_A is 2-syzygy.

Then the implications $(1) \Leftrightarrow (2) \Rightarrow (3) \Leftrightarrow (4) \Leftrightarrow (5)$ hold true.

Corollary 6.18. *Let A be a Noetherian local ring admitting the canonical module K_A . Suppose that $q \geq 2$ and K_A satisfies (S_{q+1}) . Then the following conditions are equivalent:*

- (1) K_A is $(q + 1)$ -torsionfree;
- (2) K_A is $(q + 1)$ -syzygy;
- (3) A satisfies both (S_q) and (G_q) on $\text{Supp}_A K_A$;
- (4) A satisfies both (S_q) and (G_q) .

Proof. This follows from Theorem 6.6 and the fact that $\text{Supp}_A K_A = \text{Spec } A$ ([3, Lemma 1.1]). \square

Corollary 6.19. *Let A be a Noetherian local ring with $d = \dim A$ which is a homomorphic image of a Gorenstein ring. Suppose that $q \geq \frac{d}{2} + 1$ and K_A is $(q + 1)$ -syzygy satisfying (S_q) . Then A is a Cohen-Macaulay ring.*

Proof. By Theorem 6.8, we see that A satisfies (S_q) . We may assume $d > 0$. Then $q \geq 2$, so that A is equidimensional by [3, Lemma 1.1]. Furthermore, either A is Cohen-Macaulay or $\text{depth } A \geq q$. We assume $\text{depth } A \geq q$. Since K_A satisfies (S_q) , every A -regular sequence of length at most q is K_A -regular ([11, Proposition 2.1]). The assertion follows from [9, Corollary (2.6)] (see also [10, Proposition 4.2]). \square

7. EXAMPLES OF q -GORENSTEIN RINGS

Closing this paper, in order to illustrate our theorems, we provide additional examples of Cohen-Macaulay and q -Gorenstein rings, i.e., rings with (S_q) and (G_{q-1}) conditions, or equivalently, rings with (\tilde{G}_{q-1}) condition.

Theorem 7.1. *Let A be a Gorenstein local ring with $d = \dim A \geq 3$ and let a_1, a_2, \dots, a_d be a system of parameters of A . Let $\mathbf{a} = (a_1, a_2, \dots, a_\ell)$ ($3 \leq \ell \leq d$) and let*

$$\mathcal{R} = A[a_1t, a_2t, \dots, a_\ell t] \subseteq A[t]$$

be the Rees algebra of \mathbf{a} , where t denotes an indeterminate. Then, \mathcal{R} is not a Gorenstein ring, but it is a Cohen-Macaulay $(\ell + 1)$ -Gorenstein ring of dimension $d + 1$.

Proof. Recall that \mathcal{R} is a Cohen-Macaulay ring of dimension $d + 1$. Let $S = A[X_1, X_2, \dots, X_\ell]$ be the polynomial ring over A and let $\varphi : S \rightarrow \mathcal{R}$ denote the surjective homomorphism of A -algebras defined by $\varphi(X_i) = a_it$ for each $1 \leq i \leq \ell$. The homomorphism φ preserves the grading and $\text{Ker}(\varphi) = \mathbf{I}_2 \begin{pmatrix} X_1 & X_2 & \dots & X_\ell \\ a_1 & a_2 & \dots & a_\ell \end{pmatrix}$ is the perfect ideal of S of grade $\ell - 1$ generated by the 2×2 minors of the matrix $\begin{pmatrix} X_1 & X_2 & \dots & X_\ell \\ a_1 & a_2 & \dots & a_\ell \end{pmatrix}$. We set $I = \text{Ker}(\varphi)$. We then have the following.

Claim 2. Let $P \in \text{Spec } S$ such that $I \subseteq P$ but $(X_1, X_2, \dots, X_\ell) + (a_1, a_2, \dots, a_\ell) \not\subseteq P$. Then, S_P/IS_P is a Gorenstein ring.

Proof of Claim 2. We may assume that $X_1 \notin P$. Let $\tilde{S} = S[\frac{1}{X_1}]$, $\tilde{A} = A[X_1, \frac{1}{X_1}]$, and $Y_i = \frac{X_i}{X_1}$ for $2 \leq i \leq \ell$. Then, $\tilde{S} = \tilde{A}[Y_2, Y_3, \dots, Y_\ell]$ and $I\tilde{S} = (a_i - a_1Y_i \mid 2 \leq i \leq \ell)\tilde{S}$. Because $a_1\tilde{S} + (a_i - a_1Y_i \mid 2 \leq i \leq \ell)\tilde{S} = (a_i \mid 1 \leq i \leq \ell)\tilde{S}$ and a_1, a_2, \dots, a_ℓ is an \tilde{S} -regular sequence, the sequence $a_2 - a_1Y_2, a_3 - a_1Y_3, \dots, a_\ell - a_1Y_\ell$ is \tilde{S}_P -regular, so that S_P/IS_P is a Gorenstein ring. \square

Let $P \in \text{Spec } S$ and suppose that $I \subseteq P$. We set $\mathfrak{p} = \varphi(P) \in \text{Spec } \mathcal{R}$. Then, $(X_1, X_2, \dots, X_\ell) + (a_1, a_2, \dots, a_\ell) \not\subseteq P$ if $\text{ht}_S P < 2\ell$, while

$$\text{ht}_{\mathcal{R}} \mathfrak{p} = \text{ht}_{S/I} P/I = \text{ht}_S P - (\ell - 1).$$

Therefore, if $\text{ht}_{\mathcal{R}} \mathfrak{p} < \ell + 1$, then $\text{ht}_S P - (\ell - 1) < \ell + 1$, that is $\text{ht}_S P < 2\ell$, so that $(X_1, X_2, \dots, X_\ell) + (a_1, a_2, \dots, a_\ell) \not\subseteq P$, whence $\mathcal{R}_{\mathfrak{p}} = S_P/IS_P$ is a Gorenstein ring by Claim 2. Thus, \mathcal{R} is an $(\ell + 1)$ -Gorenstein ring. \square

Since the proofs of the following assertions are standard, we left them to the interested readers.

Lemma 7.2. *Let $\varphi : A \rightarrow B$ be a flat local homomorphism of Noetherian local rings and $q \geq 1$ be an integer. Then the following conditions are equivalent:*

- (1) B is a q -Gorenstein ring;
- (2) A is a q -Gorenstein ring and $B_P/\mathfrak{p}B_P$ is a Gorenstein ring for every $P \in \text{Spec } B$ with $\text{depth } B_P < q$, where $\mathfrak{p} = \varphi^{-1}(P)$.

Proposition 7.3. *Let R be a Noetherian ring. Then the following assertions hold true.*

- (1) *Let $q \geq 1$ be an integer. Then $R[t]$ is a q -Gorenstein ring if and only if R is a q -Gorenstein ring, where t is an indeterminate.*
- (2) *Let H be a symmetric numerical semigroup. If R is a q -Gorenstein ring, then the semigroup ring $R[H]$ of H over R is a q -Gorenstein ring.*
- (3) *Let $X = \{X_{ij}\}_{1 \leq i \leq \ell, 1 \leq j \leq m}$ be indeterminates where $\ell, m \geq 2$, and set $T = R[X]$. Let t be an integer such that $2 \leq t \leq \min \{\ell, m\}$ and let $I = \mathbf{I}_t(X)$ denote the ideal of S generated by the $t \times t$ minors of the matrix X . We set $S = T/I$.*
 - (a) *Let $\ell = m$. If R is a q -Gorenstein ring, then S is a q -Gorenstein ring.*
 - (b) *Suppose that R is a field and let $t = 2$. Then S is a d -Gorenstein ring, where $d = \ell + m - 1$.*

REFERENCES

- [1] M. Auslander and M. Bridger, Stable module theory, Amer. Math. Soc., Memoirs, **94**, 1969. [1](#), [2](#), [12](#), [13](#), [15](#), [17](#)
- [2] Y. Aoyama, Some basic results on canonical modules, J. Math. Kyoto Univ, **23** (1983), no. 1, 85–94. [1](#), [2](#), [3](#), [7](#), [15](#)
- [3] Y. Aoyama and S. Goto, On the endomorphism ring of the canonical module. J. Math. Kyoto Univ., **25** (1985), no. 1, 21–30. [1](#), [17](#), [18](#)
- [4] J. Brennan, L. Ghezzi, J. Hong and W. V. Vasconcelos, Generalization of bi-canonical degrees, São Paulo J. Math. Sci. (2022), <https://doi.org/10.1007/s40863-022-00333-9>. [2](#), [4](#)
- [5] M. P. Brodmann and R. Y. Sharp, Local cohomology, An algebraic introduction with geometric applications, Cambridge Studies in Advanced Mathematics **136**, 2nd edition, Cambridge University Press, 1993. [1](#), [2](#), [3](#)
- [6] W. Bruns and J. Herzog, Cohen-Macaulay Rings, Cambridge University Press, 2013. [1](#), [2](#), [4](#), [8](#), [9](#), [15](#)
- [7] S. Dey and R. Takahashi, On the subcategories of n -torsionfree modules and related modules, Collect. Math., **74** (2023), no. 1, 113–132. [14](#)
- [8] E. G. Evans and P. Griffith, Syzygies, London Mathematical Society Lecture Note Series, **106**, Cambridge University Press, Cambridge, 1985. [8](#)
- [9] R. Fossum, H.-B. Foxby, P. Griffith, I. Reiten, Minimal injective resolutions with applications to dualizing modules and Gorenstein modules, Inst. Hautes Études Sci. Publ. Math., **45** (1975), 193–215. [18](#)
- [10] H.-B. Foxby, Gorenstein modules and related modules, Math. Scand., **31** (1972), 267–284. [1](#), [13](#), [18](#)
- [11] H.-B. Foxby, n -Gorenstein rings, Proc. Amer. Math. Soc., **42** (1974), 67–72. [1](#), [2](#), [5](#), [8](#), [13](#), [15](#), [16](#), [18](#)
- [12] S. Goto, N. Matsuoka, and T. T. Phuong, Almost Gorenstein rings, J. Algebra, **379** (2013), 355–381. [11](#), [12](#)
- [13] S. Goto and K.-i. Watanabe, On graded rings I, J. Math. Soc. Japan, **30** (1978), no. 2, 179–213. [1](#), [8](#)
- [14] J. Herzog and E. Kunz, Der kanonische Modul eines Cohen-Macaulay-Rings, Lecture Notes in Mathematics, 238, Springer-Verlag, Berlin-New York, 1971. [1](#), [2](#), [3](#), [4](#), [7](#), [11](#)
- [15] E. Kunz, The value-semigroup of a one-dimensional Gorenstein ring, Proc. Amer. Math. Soc., **25** (1970), 748–751. [1](#)
- [16] I. Reiten, The converse to a theorem of Sharp on Gorenstein modules, Proc. Amer. Math. Soc., **32** (1972), 417–420. [2](#), [9](#)
- [17] P. Schenzel, Zur lokalen Kohomologie des kanonischen Moduls. *Math. Z.* **165** (1979), no. 3, 223–230. [15](#)
- [18] R. Y. Sharp, On Gorenstein modules over a complete Cohen-Macaulay ring, Quart. J. Math., **22**, no. 3 (1971), 425–434. [2](#)

- [19] R. Stanley, Hilbert functions of graded algebras, *Adv. Math.*, **28** (1978), no. 1, 57–83. [1](#)
- [20] R. Stanley, *Combinatorics and commutative algebra*, Second Edition, Birkhäuser, Boston, 1996. [8](#)
- [21] W. V. Vasconcelos, Reflexive modules over Gorenstein rings, *Proc. Amer. Math. Soc.*, **19** (1968), 1349–1355. [2](#), [4](#), [8](#)
- [22] W. V. Vasconcelos, *Integral Closure. Rees algebras, multiplicities, algorithms*. Springer Monographs in Mathematics. Berlin, Springer-Verlag, 2005. [10](#)

SCHOOL OF POLITICAL SCIENCE AND ECONOMICS, MEIJI UNIVERSITY, 1-9-1 EIFUKU, SUGINAMI-KU, TOKYO 168-8555, JAPAN

E-mail address: endo@meiji.ac.jp

URL: <https://www.isc.meiji.ac.jp/~endo/>

DEPARTMENT OF MATHEMATICS, NEW YORK CITY COLLEGE OF TECHNOLOGY AND THE GRADUATE CENTER, THE CITY UNIVERSITY OF NEW YORK, 300 JAY STREET, BROOKLYN, NY 11201, U.S.A.; 365 FIFTH AVENUE, NEW YORK, NY 10016, U.S.A.

E-mail address: lghezzi@citytech.cuny.edu

DEPARTMENT OF MATHEMATICS, SCHOOL OF SCIENCE AND TECHNOLOGY, MEIJI UNIVERSITY, 1-1-1 HIGASHIMITA, TAMA-KU, KAWASAKI 214-8571, JAPAN

E-mail address: shirogoto@gmail.com

DEPARTMENT OF MATHEMATICS, SOUTHERN CONNECTICUT STATE UNIVERSITY, 501 CRESCENT STREET, NEW HAVEN, CT 06515-1533, U.S.A.

E-mail address: hongj2@southernct.edu

MATHEMATICS LABORATORY, SAPPORO COLLEGE, HOKKAIDO UNIVERSITY OF EDUCATION, 1-3 AINOSATO 5-3, KITA-KU, SAPPORO 002-8502, JAPAN

E-mail address: iai.shinichiro@s.hokkyodai.ac.jp

DEPARTMENT OF MATHEMATICS, SCHOOL OF SCIENCE AND TECHNOLOGY, MEIJI UNIVERSITY, 1-1-1 HIGASHIMITA, TAMA-KU, KAWASAKI 214-8571, JAPAN

E-mail address: toshinorikobayashi@icloud.com

DEPARTMENT OF MATHEMATICS, SCHOOL OF SCIENCE AND TECHNOLOGY, MEIJI UNIVERSITY, 1-1-1 HIGASHIMITA, TAMA-KU, KAWASAKI 214-8571, JAPAN

E-mail address: naomatsu@meiji.ac.jp

GRADUATE SCHOOL OF MATHEMATICS, NAGOYA UNIVERSITY, FUROCHO, CHIKUSAKU, NAGOYA, AICHI 464-8602, JAPAN

E-mail address: takahashi@math.nagoya-u.ac.jp