

JOINT VALUE-DISTRIBUTION THEOREMS ON LERCH ZETA-FUNCTIONS. II

A. Laurinćikas

Faculty of Mathematics and Informatics, Vilnius University, Naugarduko 24, LT-03225 Vilnius
(e-mail: antanas.laurincikas@maf.vu.lt)

Kohji Matsumoto

Graduate School of Mathematics, Nagoya University, Chikusa-ku, Nagoya, 464-8602, Japan
(e-mail: kohjimat@math.nagoya-u.ac.jp)

Abstract. We give corrected statements of some theorems from [5] and [6] on joint value distribution of Lerch zeta-functions (limit theorems, universality, functional independence). We also present a new direct proof of a joint limit theorem in the space of analytic functions and an extension of a joint universality theorem.

Key words: Lerch zeta-function, limit theorem, probability measure, space of analytic functions, support, universality, weak convergence.

1. INTRODUCTION

Let, as usual, $s = \sigma + it$, denote a complex variable, and let \mathbb{N} , \mathbb{N}_0 , \mathbb{Z} , \mathbb{R} and \mathbb{C} be the set of all positive integers, non-negative integers, integers, real and complex numbers, respectively. The Lerch zeta-function $L(\lambda, \alpha, s)$ with fixed parameters $\alpha, \lambda \in \mathbb{R}$, $0 < \alpha \leq 1$, for $\sigma > 1$, is defined by

$$L(\lambda, \alpha, s) = \sum_{m=0}^{\infty} \frac{e^{2\pi i \lambda m}}{(m + \alpha)^s}.$$

If $\lambda \in \mathbb{Z}$, then the Lerch zeta-function becomes the Hurwitz zeta-function

$$\zeta(s, \alpha) = \sum_{m=0}^{\infty} \frac{1}{(m + \alpha)^s}, \quad \sigma > 1,$$

and $\zeta(s, \alpha)$, for $\alpha = 1$, reduces to the Riemann zeta-function.

The paper is conditioned by [5] and [6], where the joint value distribution of Lerch zeta-functions was considered and some inaccuracies in the statements of some theorems in these papers were remained. The aim of this paper is to correct and comment the results of [5] and [6], and to give for some of them new proofs

as well as certain their extensions. For this, first we recall the results of [5]. Denote by $\mathcal{B}(S)$ the class of Borel sets of the space S , and let, for $T > 0$,

$$\nu_T^t(\dots) = \frac{1}{T} \text{meas}\{t \in [0, T] : \dots\},$$

where $\text{meas}\{A\}$ is the Lebesgue measure of a measurable set $A \subset \mathbb{R}$, in place of dots a condition satisfied by t is to be written, and the sign t in ν_T^t means that the measure is taken over $t \in [0, T]$.

Let $r \in \mathbb{N} \setminus \{1\}$, and let $L(\lambda_1, \alpha_1, \sigma_1 + it), \dots, L(\lambda_r, \alpha_r, \sigma_r + it)$ be a collection of Lerch zeta-functions. Throughout the paper, as in [5], we suppose that $\lambda_j \notin \mathbb{Z}$, $j = 1, \dots, r$.

Theorem 1 of [5] remains without any changes. It asserts that, for $\min_{1 \leq j \leq r} \sigma_j > \frac{1}{2}$, on $(\mathbb{C}^r, \mathcal{B}(\mathbb{C}^r))$, there exists a probability measure P such that the probability measure

$$\nu_T^t((L(\lambda_1, \alpha_1, \sigma_1 + it), \dots, L(\lambda_r, \alpha_r, \sigma_r + it)) \in A), \quad A \in \mathcal{B}(\mathbb{C}^r),$$

converges weakly to P as $T \rightarrow \infty$. Now we will present the corrected statement of Theorem 2 from [5]. This theorem deals with joint distribution of Lerch zeta-functions in the space of analytic functions, and differently from Theorem 1, gives the explicit form of the limit measure. For its statement, we need some additional notation. For $D = \{s \in \mathbb{C} : \sigma > \frac{1}{2}\}$, denote by $H(D)$ the space of analytic on D functions equipped with the topology of uniform convergence on compacta, and let $H^r(D) = \underbrace{H(D) \times \dots \times H(D)}_r$. Let $\gamma = \{s \in \mathbb{C} : |s| = 1\}$, and define

$$\Omega = \prod_{m=0}^{\infty} \gamma_m,$$

where $\gamma_m = \gamma$ for all $m \in \mathbb{N}_0$. By the Tikhonov theorem the infinite-dimensional torus Ω is a compact topological Abelian group, therefore, the probability Haar measure m_H on $(\Omega, \mathcal{B}(\Omega))$ can be defined. This gives a probability space $(\Omega, \mathcal{B}(\Omega), m_H)$. Denote by $\omega(m)$ the projection of $\omega \in \Omega$ to the coordinate space γ_m . Moreover, define $\Omega^r = \Omega_1 \times \dots \times \Omega_r$, where $\Omega_j = \Omega$ for $j = 1, \dots, r$. Then Ω^r is also a compact topological group. Denote by m_{H^r} the probability Haar measure on $(\Omega^r, \mathcal{B}(\Omega^r))$, and on the probability space $(\Omega^r, \mathcal{B}(\Omega^r), m_{H^r})$ define an $H^r(D)$ -valued random element $\underline{L}(s, \underline{\omega})$ by

$$\underline{L}(s, \underline{\omega}) = (L(\lambda_1, \alpha_1, s, \omega_1), \dots, L(\lambda_r, \alpha_r, s, \omega_r)),$$

where

$$L(\lambda_j, \alpha_j, s, \omega_j) = \sum_{m=0}^{\infty} \frac{e^{2\pi i \lambda_j m} \omega_j(m)}{(m + \alpha_j)^s},$$

$\omega_j \in \Omega$, $j = 1, \dots, r$, and $\underline{\omega} = (\omega_1, \dots, \omega_r)$. Denote by $P_{\underline{L}}$ the distribution of the random element $\underline{L}(s, \underline{\omega})$, that is

$$P_{\underline{L}}(A) = m_{H^r}(\underline{\omega} \in \Omega^r : \underline{L}(s, \underline{\omega}) \in A), \quad A \in \mathcal{B}(H^r(D)),$$

and define the probability measure P_T by

$$P_T(A) = \nu_T^\tau((L(\lambda_1, \alpha_1, s + i\tau), \dots, L(\lambda_r, \alpha_r, s + i\tau)) \in A), \quad A \in \mathcal{B}(H^r(D)).$$

We recall that the numbers a_1, \dots, a_r are algebraically independent over the field of rational numbers \mathbb{Q} if the coefficients of every polynomial p with rational coefficients satisfying $p(a_1, \dots, a_r) = 0$ are equal to zero.

THEOREM 1. *Suppose that $\alpha_1, \dots, \alpha_r$ are algebraically independent over \mathbb{Q} and that $\lambda_j \notin \mathbb{Z}$, $j = 1, \dots, r$. Then the probability measure P_T converges weakly to $P_{\underline{L}}$ as $T \rightarrow \infty$.*

Theorem 1 is the corrected statement of Theorem 2 from [5]. In Theorem 2 of [5] it is required that the numbers $\alpha_1, \dots, \alpha_r$ should be transcendental, however, this is not sufficient for the proof of Lemma 9 of [5]. Since, for $\sigma > 1$, the shifting parameters of the functions $L(\lambda_1, \alpha_1, s), \dots, L(\lambda_r, \alpha_r, s)$ are different, for the proof of Lemma 9 of [5] we need a limit theorem on the torus Ω^r (see Lemma 4 below). Moreover, in this theorem, the limit measure must be the Haar measure. Therefore, we have to use a condition that the set

$$\bigcup_{j=1}^r \bigcup_{m=0}^{\infty} \{\log(m + \alpha_j)\}$$

should be linearly independent over \mathbb{Q} . The latter requirement is satisfied if the numbers $\alpha_1, \dots, \alpha_r$ are algebraically independent over \mathbb{Q} . This is also used in application of elements of ergodic theory for the identification of the limit measure. So, except for the above change in the proof of Lemma 9 of [5], the proof of Theorem 2 from [5] remains the same, and only in its statement the condition of the transcendence of $\alpha_1, \dots, \alpha_r$ is changed by the algebraic independence over \mathbb{Q} . Below we will give a new direct proof of Theorem 1 which does not use the modified Cramér-Wald criterion (a statement of the type of Lemma 9 in [5]).

We note that Theorem 3 of [5] is true. In the case of rational λ , the Dirichlet series for $L(\lambda, \alpha, s)$ is reduced to ordinary Dirichlet series (with exponents $\log m$), therefore, even in the multidimensional case, we can use a limit theorem on the torus

$$\hat{\Omega} = \prod_p \gamma_p,$$

where $\gamma_p = \gamma$ for all primes p , and the linear independence over \mathbb{Q} of logarithms of prime numbers. The further proof runs in a standard way.

The same changes must be done also in Theorems 1 and 2 from [6], on the joint universality and functional independence of Lerch zeta-functions, since their

proof is based on Theorem 2 of [5]. Thus, in Theorems 1 and 2 of [6] the condition that $\alpha_1, \dots, \alpha_r$ are transcendental must be changed by a more strong requirement that the numbers $\alpha_1, \dots, \alpha_r$ are algebraically independent over \mathbb{Q} .

We also note that the same changes must be done in Theorems 5.3.1, 5.3.2, 6.3.1 and 7.2.1 of [4].

For example, the transcendental numbers $\alpha_1 = e^{-1}$ and $\alpha_2 = e^{-2}$ are not algebraically independent over \mathbb{Q} , therefore they do not satisfy the hypotheses of Theorem 1. On the other hand, it is known that the numbers e^π and π are algebraically independent over \mathbb{Q} . So, we can take, for example, $\alpha_1 = e^\pi/12$ and $\alpha_2 = \pi/4$ in Theorem 1.

Now we state one generalization of Theorems 1 and 2 (after the above correction) from [6]. Let $\lambda_1, \dots, \lambda_r$ be arbitrary rational numbers with denominators q_1, \dots, q_r , respectively. Denote by $k = [q_1, \dots, q_r]$ the least common multiple, and define

$$A = \begin{pmatrix} e^{2\pi i \lambda_1} & e^{2\pi i \lambda_2} & \dots & e^{2\pi i \lambda_r} \\ e^{4\pi i \lambda_1} & e^{4\pi i \lambda_2} & \dots & e^{4\pi i \lambda_r} \\ \dots & \dots & \dots & \dots \\ e^{2\pi k i \lambda_1} & e^{2\pi k i \lambda_2} & \dots & e^{2\pi k i \lambda_r} \end{pmatrix}.$$

THEOREM 2. *Suppose that $\alpha_1, \dots, \alpha_r$ are algebraically independent over \mathbb{Q} and that $\text{rank}(A) = r$. Let K_j be a compact subset of the strip $D_0 = \{s \in \mathbb{C} : \frac{1}{2} < \sigma < 1\}$ with connected complement, and let $f_j(s)$ be a continuous on K_j function which is analytic in the interior of K_j , $j = 1, \dots, r$. Then, for every $\varepsilon > 0$*

$$\liminf_{T \rightarrow \infty} \nu_T^r \left(\sup_{1 \leq j \leq r} \sup_{s \in K_j} |L(\lambda_j, \alpha_j, s + i\tau) - f_j(s)| < \varepsilon \right) > 0.$$

THEOREM 3. *Suppose that $\alpha_1, \dots, \alpha_r$ are algebraically independent over \mathbb{Q} and that $\text{rank}(A) = r$. Let F_j be a continuous on \mathbb{C}^{N^r} function, $j = 0, \dots, l$, and*

$$\sum_{j=0}^l s^j F_j(L(\lambda_1, \alpha_1, s), \dots, L(\lambda_r, \alpha_r, s), L'(\lambda_1, \alpha_1, s), \dots, L'(\lambda_r, \alpha_r, s), \dots, L^{(N-1)}(\lambda_1, \alpha_1, s), \dots, L^{(N-1)}(\lambda_r, \alpha_r, s)) = 0$$

identically for $s \in \mathbb{C}$. Then $F_j \equiv 0$, $j = 0, \dots, l$.

Theorem 3 is deduced from Theorem 2 in the same way as in [6] where Theorem 2 of [6] is obtained from Theorem 1 of [6].

2. PROOF OF THEOREM 1

2.1. Joint limit theorems for Dirichlet polynomials. First we will prove a joint limit theorem on the torus Ω^r for the probability measure

$$Q_{T,r}(A) = \nu_T^r \left(\left((m + \alpha_1)^{i\tau} : m \in \mathbb{N}_0 \right), \dots, \left((m + \alpha_r)^{i\tau} : m \in \mathbb{N}_0 \right) \in A \right), \quad A \in \mathcal{B}(\Omega^r).$$

LEMMA 4. *The probability measure $Q_{T,r}$ converges weakly to the Haar measure m_{H^r} on $(\Omega^r, \mathcal{B}(\Omega^r))$ as $T \rightarrow \infty$.*

Proof. The dual group of Ω^r is

$$\bigoplus_{j=1}^r \bigoplus_{m=0}^{\infty} \mathbb{Z}_{m_j},$$

where $\mathbb{Z}_{m_j} = \mathbb{Z}$ for all $m \in \mathbb{N}_0$ and $j = 1, \dots, r$. The element $(\underline{k}_1, \dots, \underline{k}_r) = (k_{01}, k_{11}, \dots, k_{0r}, k_{1r}, \dots) \in \bigoplus_{j=1}^r \bigoplus_{m=0}^{\infty} \mathbb{Z}_{m_j}$, where only a finite number of integers k_{mj} , $m \in \mathbb{N}_0$, $j = 1, \dots, r$, are distinct from zero, acts on Ω^r by

$$(\underline{x}_1, \dots, \underline{x}_r) \rightarrow (\underline{x}_1^{k_1}, \dots, \underline{x}_r^{k_r}) = \prod_{j=1}^r \prod_{m=0}^{\infty} x_{m_j}^{k_{mj}}, \quad x_j = (x_{1j}, x_{2j}, \dots), \quad x_{mj} \in \gamma, \quad m \in \mathbb{N}_0, \quad j = 1, \dots, r.$$

Therefore, the Fourier transform $g_{T,r}(\underline{k}_1, \dots, \underline{k}_r)$ of the measure $Q_{T,r}$ is

$$\begin{aligned} g_{T,r}(\underline{k}_1, \dots, \underline{k}_r) &= \int_{\Omega^r} \prod_{j=1}^r \prod_{m=0}^{\infty} x_{m_j}^{k_{mj}} dQ_{T,r} = \\ &= \frac{1}{T} \int_0^T \prod_{j=1}^r \prod_{m=0}^{\infty} e^{i\tau k_{mj} \log(m+\alpha_j)} d\tau = \\ &= \frac{1}{T} \int_0^T \exp \left\{ i\tau \sum_{j=1}^r \sum_{m=1}^{\infty} k_{mj} \log(m+\alpha_j) \right\} d\tau. \end{aligned}$$

Since $\alpha_1, \dots, \alpha_r$ are algebraically independent over \mathbb{Q} , we find that

$$g_{T,r}(\underline{k}_1, \dots, \underline{k}_r) = \begin{cases} 1 & \text{if } (\underline{k}_1, \dots, \underline{k}_r) = (\underline{0}, \dots, \underline{0}), \\ \frac{\exp\{iT \sum_{j=1}^r \sum_{m=0}^{\infty} k_{mj} \log(m+\alpha_j)\} - 1}{iT \sum_{j=1}^r \sum_{m=0}^{\infty} k_{mj} \log(m+\alpha_j)} & \text{if } (\underline{k}_1, \dots, \underline{k}_r) \neq (\underline{0}, \dots, \underline{0}). \end{cases}$$

Consequently, we have

$$\lim_{T \rightarrow \infty} g_{T,r}(\underline{k}_1, \dots, \underline{k}_r) = \begin{cases} 1 & \text{if } (\underline{k}_1, \dots, \underline{k}_r) = (\underline{0}, \dots, \underline{0}), \\ 0 & \text{if } (\underline{k}_1, \dots, \underline{k}_r) \neq (\underline{0}, \dots, \underline{0}). \end{cases}$$

Hence by continuity theorems for probability measures on locally compact groups ([2], Theorem 1.4.2), we obtain that the measure $Q_{T,r}$ converges weakly to m_{H^r} as $T \rightarrow \infty$.

Now let $\sigma_{1j} > \frac{1}{2}$, and, for $m, n \in \mathbb{N}_0$,

$$v_j(m, n) = \exp \left\{ - \left(\frac{m + \alpha_j}{n + \alpha_j} \right)^{\sigma_{1j}} \right\}, \quad j = 1, \dots, r.$$

Define, for $N_j \in \mathbb{N}_0$, $\hat{\omega}_j \in \Omega$, $s \in D$,

$$L_{N_j, j, n}(\lambda_j, \alpha_j, s) = \sum_{m=0}^{N_j} \frac{e^{2\pi i \lambda_j m} v_j(m, n)}{(m + \alpha_j)^s},$$

$$L_{N_j, j, n}(\lambda_j, \alpha_j, s, \hat{\omega}_j) = \sum_{m=0}^{N_j} \frac{e^{2\pi i \lambda_j m} v_j(m, n) \hat{\omega}_j(m)}{(m + \alpha_j)^s}, \quad j = 1, \dots, r,$$

and consider the weak convergence of the probability measures

$$P_{T, N_1, \dots, N_r, n}(A) = \nu_T^r((L_{N_1, 1, n}(\lambda_1, \alpha_1, s + i\tau), \dots, L_{N_r, r, n}(\lambda_r, \alpha_r, s + i\tau)) \in A)$$

and

$$\hat{P}_{T, N_1, \dots, N_r, n}(A) = \nu_T^r((L_{N_1, 1, n}(\lambda_1, \alpha_1, s + i\tau, \hat{\omega}_1), \dots, L_{N_r, r, n}(\lambda_r, \alpha_r, s + i\tau, \hat{\omega}_r)) \in A),$$

where $A \in \mathcal{B}(H^r(D))$.

THEOREM 5. *The probability measures $P_{T, N_1, \dots, N_r, n}$ and $\hat{P}_{T, N_1, \dots, N_r, n}$ both converge weakly to the same probability measure on $(H^r(D), \mathcal{B}(H^r(D)))$ as $T \rightarrow \infty$.*

Proof. Let a function $h : \Omega^r \rightarrow H^r(D)$ be defined by

$$h(\omega_1, \dots, \omega_r) = \left(\sum_{m=0}^{N_1} \frac{e^{2\pi i \lambda_1 m} v_1(m, n) \omega_1^{-1}(m)}{(m + \alpha_1)^s}, \dots, \sum_{m=0}^{N_r} \frac{e^{2\pi i \lambda_r m} v_r(m, n) \omega_r^{-1}(m)}{(m + \alpha_r)^s} \right),$$

$(\omega_1, \dots, \omega_r) \in \Omega^r$. Then the function h is continuous, and

$$h(((m + \alpha_1)^{i\tau} : m \in \mathbb{N}_0), \dots, ((m + \alpha_r)^{i\tau} : m \in \mathbb{N}_0))$$

$$= ((L_{N_1, 1, n}(\lambda_1, \alpha_1, s + i\tau), \dots, L_{N_r, r, n}(\lambda_r, \alpha_r, s + i\tau)).$$

Hence $P_{T, N_1, \dots, N_r, n} = Q_{T, r} h^{-1}$, therefore by Lemma 4 and Theorem 5.1 of [1] we obtain that the measure $P_{T, N_1, \dots, N_r, n}$ converges weakly to $m_{H^r} h^{-1}$ as $T \rightarrow \infty$.

Now let $h_1 : \Omega^r \rightarrow \Omega^r$ be given by

$$h_1(\omega_1, \dots, \omega_r) = (\omega_1 \hat{\omega}_1^{-1}, \dots, \omega_r \hat{\omega}_r^{-1}).$$

Then

$$(L_{N_1, 1, n}(\lambda_1, \alpha_1, s + i\tau, \hat{\omega}_1), \dots, L_{N_r, r, n}(\lambda_r, \alpha_r, s + i\tau, \hat{\omega}_r)) =$$

$$= h \left(h_1 \left((m + \alpha_1)^{i\tau} : m \in \mathbb{N}_0 \right), \dots, (m + \alpha_r)^{i\tau} : m \in \mathbb{N}_0 \right).$$

Similarly as above we find that $\hat{P}_{T,N_1,\dots,N_r,n}$ converges weakly to the measure $m_{H^r}(hh_1)^{-1}$ as $T \rightarrow \infty$. However, the invariance of the measure m_{H^r} yields

$$m_{H^r}(hh_1)^{-1} = (m_{H^r}h_1^{-1})h^{-1} = m_{H^r}h^{-1},$$

and the theorem follows.

2.2. Limit theorems for absolutely convergent series. For $j = 1, \dots, r$, let $\omega_j \in \Omega$, and

$$L_{n,j}(\lambda_j, \alpha_j, s) = \sum_{m=0}^{\infty} \frac{e^{2\pi i \lambda_j m} v_j(m, n)}{(m + \alpha_j)^s}$$

and

$$L_{n,j}(\lambda_j, \alpha_j, s, \omega_j) = \sum_{m=0}^{\infty} \frac{e^{2\pi i \lambda_j m} \omega_j(m) v_j(m, n)}{(m + \alpha_j)^s}.$$

Note that these two series both converge absolutely for $\sigma > \frac{1}{2}$. Define the probability measures

$$P_{T,n}(A) = \nu_T^\tau \left((L_{n,1}(\lambda_1, \alpha_1, s + i\tau), \dots, L_{n,r}(\lambda_r, \alpha_r, s + i\tau)) \in A \right)$$

and

$$\hat{P}_{T,n}(A) = \nu_T^\tau \left((L_{n,1}(\lambda_1, \alpha_1, s + i\tau, \omega_1), \dots, L_{n,r}(\lambda_r, \alpha_r, s + i\tau, \omega_r)) \in A \right),$$

where $A \in \mathcal{B}(H^r(D))$.

THEOREM 6. *On $(H^r(D), \mathcal{B}(H^r(D)))$ there exists a probability measure P_n such that the measures $P_{T,n}$ and $\hat{P}_{T,n}$ both converge weakly to P_n as $T \rightarrow \infty$.*

Proof. We apply Theorem 5 with $N_1 = \dots = N_r \stackrel{\text{def}}{=} N$. Then by Theorem 5 the measures $P_{T,N_1,\dots,N_r,n} \stackrel{\text{def}}{=} P_{T,N,n}$ and $\hat{P}_{T,N_1,\dots,N_r,n} \stackrel{\text{def}}{=} \hat{P}_{T,N,n}$ both converge to the same measure $P_{N,n}$ as $T \rightarrow \infty$.

First we observe that, for any fixed n , the family $\{P_{N,n} : N \in \mathbb{N}_0\}$ is tight. Let η be a random variable defined on a certain probability space $(\hat{\Omega}, \mathcal{B}(\hat{\Omega}), \mathbb{P})$ and uniformly distributed on $[0, 1]$. Define, for $j = 1, \dots, r$,

$$X_{T,N,j,n} = X_{T,N,j,n}(s) = L_{N,j,n}(\lambda_j, \alpha_j, s + iT\eta),$$

which is an $H(D)$ -valued random element defined on $(\hat{\Omega}, \mathcal{B}(\hat{\Omega}), \mathbb{P})$. Then by Theorem 5

$$\underline{X}_{T,N,n} \stackrel{\text{def}}{=} (X_{T,N,1,n}, \dots, X_{T,N,r,n}) \xrightarrow[T \rightarrow \infty]{\mathcal{D}} \underline{X}_{N,n}, \quad (1)$$

where $\underline{X}_{N,n} = (X_{N,1,n}, \dots, X_{N,r,n})$ is an $H^r(D)$ -valued random element with the distribution $P_{N,n}$, and $\xrightarrow{\mathcal{D}}$ means the convergence in distribution.

For the investigation of the weak convergence of the measures $P_{T,n}$ and $\hat{P}_{T,n}$ we need a metric on $H^r(D)$ which induces its topology. Let $\{K_n\}$ be a sequence of compact subsets of D such that $\bigcup_{n=1}^{\infty} K_n = D$, $K_n \subset K_{n+1}$, and if K is a compact of D , then $K \subseteq K_n$ for some n . Then

$$\rho(f, g) = \sum_{n=1}^{\infty} 2^{-n} \frac{\rho_n(f, g)}{1 + \rho_n(f, g)}, \quad f, g \in H(D),$$

$$\rho_n(f, g) = \sup_{s \in K_n} |f(s) - g(s)|,$$

is a metric on $H(D)$ which induces its topology. Then

$$\rho_r(\underline{f}, \underline{g}) = \max_{1 \leq j \leq r} \rho(f_j, g_j),$$

where $\underline{f} = (f_1, \dots, f_r) \in H^r(D)$, $\underline{g} = (g_1, \dots, g_r) \in H^r(D)$, is a desired metric on $H^r(D)$.

The series for $L_{n,j}(\lambda_j, \alpha_j, s)$, $j = 1, \dots, r$, converge absolutely for $\sigma > \frac{1}{2}$. Therefore, for $M_{lj} > 0$, $j = 1, \dots, r$, $l \in \mathbb{N}$, and some $\sigma_l > 1/2$

$$\begin{aligned} & \limsup_{T \rightarrow \infty} \mathbb{P} \left(\sup_{s \in K_l} |X_{T,N,j,n}(s)| > M_{lj} \quad \text{for at least one } j = 1, \dots, r \right) \leq \\ & \leq \sum_{j=1}^r \limsup_{T \rightarrow \infty} \mathbb{P} \left(\sup_{s \in K_l} |X_{T,N,j,n}(s)| > M_{lj} \right) \leq \\ & \leq \sum_{j=1}^r \frac{1}{M_{lj}} \sup_{N \geq 0} \limsup_{T \rightarrow \infty} \frac{1}{T} \int_0^T \sup_{s \in K_l} |L_{N,j,n}(\lambda_j, \alpha_j, s + i\tau)| d\tau \ll_l \\ & \ll_l \sum_{j=1}^r \frac{1}{M_{lj}} \sup_{N \geq 0} \limsup_{T \rightarrow \infty} \frac{1}{T} \int_0^{2T} |L_{N,j,n}(\lambda_j, \alpha_j, \sigma_l + it)| dt \ll_l \\ & \ll_l \sum_{j=1}^r \frac{1}{M_{lj}} \sup_{N \geq 0} \limsup_{T \rightarrow \infty} \left(\frac{1}{2T} \int_0^{2T} |L_{N,j,n}(\lambda_j, \alpha_j, \sigma_l + it)|^2 dt \right)^{1/2} \\ & \ll_l \sum_{j=1}^r \frac{1}{M_{lj}} \sup_{N \geq 0} \left(\sum_{m=0}^N \frac{v_j^2(m, n)}{(m + \alpha_j)^{2\sigma_l}} \right)^{1/2} \leq \\ & \leq C_l \sum_{j=1}^r \frac{1}{M_{lj}} \left(\sum_{m=0}^{\infty} \frac{1}{(m + \alpha_j)^{2\sigma_l}} \right)^{1/2} \stackrel{def}{=} C_l \sum_{j=1}^r \frac{R_{lj}}{M_{lj}} \end{aligned}$$

with a certain $C_l > 0$ and

$$R_{lj} = \left(\sum_{m=0}^{\infty} \frac{1}{(m + \alpha_j)^{2\sigma_l}} \right)^{1/2} < \infty.$$

Taking $M_{lj} = C_l R_{lj} 2^l r / \varepsilon$, hence we find that

$$\limsup_{T \rightarrow \infty} \mathbb{P} \left(\sup_{s \in K_l} |X_{T,N,j,n}(s)| > M_{lj} \quad \text{for at least one } j = 1, \dots, r \right) \leq \frac{\varepsilon}{2^l}. \quad (2)$$

Now (1) and (2) show that, for all $l \in \mathbb{N}$,

$$\mathbb{P}(\sup_{s \in K_l} |X_{N,j,n}(s)| > M_{lj} \quad \text{for at least one } j = 1, \dots, r) \leq \frac{\varepsilon}{2^l}. \quad (3)$$

Define

$$H_\varepsilon^r = \{(f_1, \dots, f_r) \in H^r(D) : \sup_{s \in K_l} |f_j(s)| \leq M_{lj}, j = 1, \dots, r, l \in \mathbb{N}\}.$$

Then the set H_ε^r is compact, and by (3)

$$\mathbb{P}(\underline{X}_{N,n}(s) \in H_\varepsilon^r) \geq 1 - \varepsilon,$$

or, by the definition of $\underline{X}_{N,n}$,

$$P_{N,n}(H_\varepsilon^r) \geq 1 - \varepsilon$$

for all $N \in \mathbb{N}_0$. This proves the tightness of the family $\{P_{N,n} : N \in \mathbb{N}_0\}$.

By the definition, for $j = 1, \dots, r$,

$$\lim_{N \rightarrow \infty} L_{N,j,n}(\lambda_j, \alpha_j, s) = L_{n,j}(\lambda_j, \alpha_j, s),$$

the convergence being uniform on compact subsets of D . Therefore, for every $\varepsilon > 0$,

$$\begin{aligned} & \lim_{N \rightarrow \infty} \limsup_{T \rightarrow \infty} \nu_T^r(\rho_r(\underline{L}_{N,n}(s + i\tau), \underline{L}_n(s + i\tau)) \geq \varepsilon) \leq \\ & \leq \lim_{N \rightarrow \infty} \limsup_{T \rightarrow \infty} \frac{1}{T\varepsilon} \int_0^T \rho_r(\underline{L}_{N,n}(s + i\tau), \underline{L}_n(s + i\tau)) d\tau \leq \\ & \leq \lim_{N \rightarrow \infty} \limsup_{T \rightarrow \infty} \frac{1}{T\varepsilon} \sum_{j=1}^r \int_0^T \rho(L_{N,j,n}(s + i\tau), L_{n,j}(s + i\tau)) d\tau = 0. \end{aligned} \quad (4)$$

Here

$$\underline{L}_{N,n}(s) = (L_{N,1,n}(\lambda_1, \alpha_1, s), \dots, L_{N,r,n}(\lambda_r, \alpha_r, s))$$

and

$$\underline{L}_n(s) = (L_{n,1}(\lambda_1, \alpha_1, s), \dots, L_{n,r}(\lambda_r, \alpha_r, s)).$$

Now define, for $j = 1, \dots, r$,

$$X_{T,j,n} = X_{T,j,n}(s) = L_{n,j}(\lambda_j, \alpha_j, s + iT\eta),$$

and put

$$\underline{X}_{T,n} = (X_{T,1,n}, \dots, X_{T,r,n}).$$

Then in view of (4)

$$\lim_{N \rightarrow \infty} \limsup_{T \rightarrow \infty} \mathbb{P}(\rho_r(\underline{X}_{T,N,n}, \underline{X}_{T,n}) \geq \varepsilon) = 0. \quad (5)$$

Since the family $\{P_{N,n} : N \in \mathbb{N}_0\}$ is tight, by the Prokhorov theorem it is relatively compact. Let $\{P_{N_1,n}\} \subset \{P_{N,n}\}$ be such that $P_{N_1,n}$ converges weakly to some measure P_n as $N_1 \rightarrow \infty$. Then we have that

$$\underline{X}_{N_1,n} \xrightarrow[N_1 \rightarrow \infty]{\mathcal{D}} P_n. \quad (6)$$

The space $H^r(D)$ is separable, and (1), (5) and (6) show that the hypothesis of Theorem 4.2 from [1] are satisfied. Consequently,

$$\underline{X}_{T,n} \xrightarrow[T \rightarrow \infty]{\mathcal{D}} P_n, \quad (7)$$

and this is equivalent to the weak convergence of the measure $P_{T,n}$ to P_n as $T \rightarrow \infty$.

Formula (7) shows that the measure P_n is independent of the sequence N_1 . Therefore, we have

$$\underline{X}_{N,n} \xrightarrow[N \rightarrow \infty]{\mathcal{D}} P_n. \quad (8)$$

Now, repeating the above arguments for the random elements

$$\widehat{\underline{X}}_{T,N,n} = (L_{N,1,n}(\lambda_1, \alpha_1, s + iT\eta, \omega), \dots, L_{N,r,n}(\lambda_r, \alpha_r, s + iT\eta, \omega))$$

and

$$\widehat{\underline{X}}_{T,n} = (L_{n,1}(\lambda_1, \alpha_1, s + iT\eta, \omega), \dots, L_{n,r}(\lambda_r, \alpha_r, s + iT\eta, \omega)),$$

and using (8) we similarly obtain that the measure $\widehat{P}_{T,n}$ also converges weakly to P_n as $T \rightarrow \infty$. The theorem is proved.

2.3. Proof of Theorem 1. We start with approximation in mean of the vectors $(L(\lambda_1, \alpha_1, s), \dots, L(\alpha_1, \lambda_1, s))$ and $(L(\lambda_1, \alpha_1, s, \omega_1), \dots, L(\lambda_r, \alpha_r, s, \omega_r))$ by the vectors $(L_{n,1}(\lambda_1, \alpha_1, s), \dots, L_{n,r}(\lambda_r, \alpha_r, s))$ and $(L_{n,1}(\lambda_1, \alpha_1, s, \omega_1), \dots, L_{n,r}(\lambda_r, \alpha_r, s, \omega_r))$, respectively. Let

$$\begin{aligned} \underline{L}(s) &= (L(\lambda_1, \alpha_1, s), \dots, L(\lambda_r, \alpha_r, s)), \\ \underline{L}(s, \underline{\omega}) &= (L(\lambda_1, \alpha_1, s, \omega_1), \dots, L(\lambda_r, \alpha_r, s, \omega_r)) \end{aligned}$$

and

$$\underline{L}_n(s, \underline{\omega}) = (L_{n,1}(\lambda_1, \alpha_1, s, \omega_1), \dots, L_{n,r}(\lambda_r, \alpha_r, s, \omega_r)).$$

LEMMA 7. *We have*

$$\lim_{n \rightarrow \infty} \limsup_{T \rightarrow \infty} \frac{1}{T} \int_0^T \rho_r(\underline{L}(s + i\tau), \underline{L}_n(s + i\tau)) d\tau = 0$$

and, for almost all $\underline{\omega} \in \Omega^r$,

$$\lim_{n \rightarrow \infty} \limsup_{T \rightarrow \infty} \frac{1}{T} \int_0^T \rho_r(\underline{L}(s + i\tau, \underline{\omega}), \underline{L}_n(s + i\tau, \underline{\omega})) d\tau = 0.$$

Proof. Since $\alpha_1, \dots, \alpha_r$ are algebraically independent over \mathbb{Q} , they are transcendental. Therefore, for each $j = 1, \dots, r$, Lemmas 5.2.11 and 5.2.13 of [4] imply

$$\lim_{n \rightarrow \infty} \limsup_{T \rightarrow \infty} \frac{1}{T} \int_0^T \rho(L(\lambda_j, \alpha_j, s + i\tau), L_{n,j}(\lambda_j, \alpha_j, s + i\tau)) d\tau = 0$$

and, for almost all $\omega_j \in \Omega$,

$$\lim_{n \rightarrow \infty} \limsup_{T \rightarrow \infty} \frac{1}{T} \int_0^T \rho(L(\lambda_j, \alpha_j, s + i\tau, \omega_j), L_{n,j}(\lambda_j, \alpha_j, s + i\tau, \omega_j)) d\tau = 0.$$

From this and the definition of ρ_r the lemma follows.

For $A \in \mathcal{B}(H^r(D))$, define

$$\widehat{P}_T(A) = \nu_T^r((L(\lambda_1, \alpha_1, s + i\tau, \omega_1), \dots, L(\lambda_r, \alpha_r, s + i\tau, \omega_r)) \in A).$$

THEOREM 8. *On $(H^r(D), \mathcal{B}(H^r(D)))$ there exists a probability measure P such that the measures P_T and \widehat{P}_T (for almost all $\underline{\omega}$) both converge weakly to P as $T \rightarrow \infty$.*

Proof. We use the same way as that in the proof of Theorem 6. First we will show that the family of probability measures $\{P_n : n \in \mathbb{N}_0\}$ is tight.

By Theorem 6

$$\underline{X}_{T,n} \xrightarrow[T \rightarrow \infty]{\mathcal{D}} \underline{X}_n, \tag{9}$$

where $\underline{X}_n = (X_{n,1}, \dots, X_{n,r})$ is an $H^r(D)$ -valued random element with the distribution P_n .

Since $L_{n,j}(\lambda_j, \alpha_j, s)$ is convergent absolutely for $\sigma > 1/2, j = 1, \dots, r$, we have

$$\lim_{T \rightarrow \infty} \frac{1}{T} \int_0^T |L_{n,j}(\lambda_j, \alpha_j, \sigma + it)|^2 dt = \sum_{m=0}^{\infty} \frac{v_j^2(m, n)}{(m + \alpha_j)^{2\sigma}} \leq \sum_{m=0}^{\infty} \frac{1}{(m + \alpha_j)^{2\sigma}}$$

uniformly in n . Hence it is not difficult to see that, for some $\sigma_l > 1/2$,

$$\begin{aligned} \limsup_{T \rightarrow \infty} \frac{1}{T} \int_0^T \sup_{s \in K_l} |L_{n,j}(\lambda_j, \alpha_j, s+i\tau)| d\tau &\leq C_l \limsup_{T \rightarrow \infty} \left(\frac{1}{2T} \int_0^{2T} |L_{n,j}(\lambda_j, \alpha_j, \sigma_l+it)|^2 dt \right)^{1/2} \\ &\leq C_l R_{lj} < \infty \end{aligned}$$

with a certain constant $C_l > 0$, where

$$R_{lj} = \left(\sum_{m=0}^{\infty} \frac{1}{(m + \alpha_j)^{2\sigma_l}} \right)^{1/2}.$$

Therefore, for $M_{lj} = C_l R_{lj} 2^l r / \varepsilon$, $j = 1, \dots, r$, $l \in \mathbb{N}$,

$$\limsup_{T \rightarrow \infty} \mathbb{P}(\sup_{s \in K_l} |X_{T,j,n}(s)| > M_{lj} \text{ for at least one } j = 1, \dots, r) \leq \frac{\varepsilon}{2^l}.$$

This and (9) imply

$$\mathbb{P}(\sup_{s \in K_l} |X_{n,j}(s)| > M_{lj} \text{ for at least one } j = 1, \dots, r) \leq \frac{\varepsilon}{2^l}.$$

Hence we find that

$$P_n(H_\varepsilon^r) \geq 1 - \varepsilon$$

for all $n \in \mathbb{N}_0$, that is the family $\{P_n : n \in \mathbb{N}_0\}$ is tight.

Now let, for $j = 1, \dots, r$,

$$X_{T,j} = X_{T,j}(s) = L(\lambda_j, \alpha_j, s + iT\eta),$$

and let

$$\underline{X}_T = (X_{T,1}, \dots, X_{T,r}).$$

Then in view of Lemma 7, for every $\varepsilon > 0$,

$$\begin{aligned} \lim_{n \rightarrow \infty} \limsup_{T \rightarrow \infty} \nu_T^r(\rho_r(\underline{L}(s+i\tau), \underline{L}_n(s+i\tau)) \geq \varepsilon) &\leq \\ &\leq \frac{1}{\varepsilon T} \int_0^T \rho_r(\underline{L}(s+i\tau), \underline{L}_n(s+i\tau)) d\tau = 0. \end{aligned}$$

Hence

$$\lim_{n \rightarrow \infty} \limsup_{T \rightarrow \infty} \mathbb{P}(\rho_r(\underline{X}_T(s), \underline{X}_{T,n}(s)) \geq \varepsilon) = 0. \quad (10)$$

The family $\{P_n : n \in \mathbb{N}_0\}$ is relatively compact. Let $\{P_{n_1}\} \subset \{P_n\}$ be such that P_{n_1} converges weakly to some measure P on $(H^r(D), \mathcal{B}(H^r(D)))$ as $n_1 \rightarrow \infty$. Then

$$\underline{X}_{n_1} \xrightarrow[n_1 \rightarrow \infty]{\mathcal{D}} P.$$

This, (10), (9) and Theorem 4.2 of [1] show that

$$\underline{X}_T \xrightarrow[T \rightarrow \infty]{\mathcal{D}} P, \quad (11)$$

that is P_T converges weakly to P as $T \rightarrow \infty$.

By (11) the measure P is independent on the sequence n_1 . Therefore,

$$\underline{X}_n \xrightarrow[n \rightarrow \infty]{\mathcal{D}} P. \quad (12)$$

By the same way as above, using (12), we obtain that the measure \widehat{P}_T for almost all $\underline{\omega}$ also converges weakly to P as $T \rightarrow \infty$.

To identify the limit measure in Theorem 8, we will apply some facts from ergodic theory.

Let $a_{\tau,j} = \{(m + \alpha_j)^{-i\tau} : m \in \mathbb{N}_0\}$, $\tau \in \mathbb{R}$, $j = 1, \dots, r$. Then $\{a_{\tau,j} : \tau \in \mathbb{R}\}$, for each $j = 1, \dots, r$, is a one - parameter group. Define the one - parameter family $\{\Phi_\tau : \tau \in \mathbb{R}\} = \{\varphi_{\tau,1}, \dots, \varphi_{\tau,r} : \tau \in \mathbb{R}\}$ of transformations on Ω^r by $\varphi_{\tau,j}(\omega_j) = a_{\tau,j}\omega_j$, $\omega_j \in \Omega_j$, $j = 1, \dots, r$. Then we have a one - parameter group of measurable transformations on Ω^r .

LEMMA 9. *The one - parameter group $\{\Phi_\tau : \tau \in \mathbb{R}\}$ is ergodic.*

Proof. Let $\chi : \Omega^r \rightarrow \gamma$ be a character. Then

$$\chi(\underline{\omega}) = \prod_{j=1}^r \prod_{m=0}^{\infty} \omega_j^{k_{mj}}(m), \quad \underline{\omega} \in \Omega^r,$$

where only a finite number of integers k_{mj} are distinct from zero. Suppose that χ is a non - principal character. Then

$$\chi(a_{\tau,1}, \dots, a_{\tau,r}) = \prod_{j=1}^r \prod_{m=0}^{\infty} (m + \alpha_j)^{-i\tau k_{mj}} = \exp\{-i\tau \sum_{j=1}^r \sum_{m=0}^{\infty} k_{mj} \log(m + \alpha_j)\},$$

where only a finite number of integers $k_{mj} \neq 0$. Since $\alpha_1, \dots, \alpha_r$ are algebraically independent, we have that there exists a $\tau_0 \neq 0$ such that

$$\chi(a_{\tau_0,1}, \dots, a_{\tau_0,r}) \neq 1.$$

The further proof runs in the same way as than in [3], Theorem 5.3.6.

Proof of Theorem 1. Let $A \in \mathcal{B}(H^r(D))$ be a continuity set of the measure P in Theorem 8. Then by Theorem 2.1 of [1] and Theorem 8

$$\lim_{T \rightarrow \infty} \nu_T^\tau((L(\lambda_1, \alpha_1, s + i\tau, \omega_1), \dots, L(\lambda_r, \alpha_r, s + i\tau, \omega_r)) \in A) = P(A) \quad (13)$$

for almost all $\underline{\omega} \in \Omega^r$. Now we fix the set A and define a random variable θ on the space $(\Omega^r, \mathcal{B}(\Omega^r), m_{H^r})$ by the formula

$$\theta(\underline{\omega}) = \begin{cases} 1 & \text{if } L(s, \underline{\omega}) \in A, \\ 0 & \text{if } L(s, \underline{\omega}) \notin A. \end{cases}$$

Denote by $\mathbb{E}(\theta)$ the expectation of θ . Then we have that

$$\mathbb{E}(\theta) = \int_{\Omega^r} \theta dm_{H^r} = m_{H^r}(\underline{\omega} \in \Omega^r : L(s, \underline{\omega}) \in A) = P_L(A). \quad (14)$$

In virtue of Lemma 9 the random process $\theta(\Phi_\tau(\underline{\omega}))$ is ergodic. Therefore, by the Birkhoff-Khinchine theorem

$$\lim_{T \rightarrow \infty} \frac{1}{T} \int_0^T \theta(\Phi_\tau(\underline{\omega})) d\tau = \mathbb{E}(\theta) \quad (15)$$

for almost all $\underline{\omega} \in \Omega^r$. On the other hand, by the definition of θ and Φ_τ we find

$$\frac{1}{T} \int_0^T \theta(\Phi_\tau(\underline{\omega})) d\tau = \nu_T^\tau((L(\lambda_1, \alpha_1, s + i\tau, \omega_1), \dots, L(\lambda_r, \alpha_r, s + i\tau, \omega_r)) \in A).$$

This and (14), (15) show that

$$\lim_{T \rightarrow \infty} \nu_T^\tau((L(\lambda_1, \alpha_1, s + i\tau, \omega_1), \dots, L(\lambda_r, \alpha_r, s + i\tau, \omega_r)) \in A) = P_L(A)$$

for almost all $\underline{\omega} \in \Omega^r$. Hence, by (13), $P(A) = P_L(A)$ for any continuity set A of the measure P , and this implies the equality $P(A) = P_L(A)$ for all $A \in \mathcal{B}(H^r(D))$. The theorem is proved.

3. PROOF OF THEOREM 2.

3.1. A limit theorem. The proof of Theorem 2 is based on a corollary of Theorem 1. Denote by $P_{\underline{L}_0}$ the restriction of the distribution $P_{\underline{L}}$ of the random element $\underline{L}(s, \underline{\omega})$ to $H^r(D_0)$.

COROLLARY 10. *Suppose that $\alpha_1, \dots, \alpha_r$ are algebraically independent over \mathbb{Q} . Then the probability measure*

$$\nu_T^\tau((L(\lambda_1, \alpha_1, s + i\tau), \dots, L(\lambda_r, \alpha_r, s + i\tau)) \in A), \quad A \in \mathcal{B}(H^r(D_0)),$$

converges weakly to $P_{\underline{L}_0}$ as $T \rightarrow \infty$.

Proof. The function $h : H(D) \rightarrow H^r(D_0)$ defined by $h(\underline{g}) = \underline{g}|_{s \in D_0}$, $\underline{g} \in H^r(D)$, obviously, is continuous. Therefore, the corollary follows from Theorem 1 and Theorem 5.1 of [1].

3.2. The support of the measure $P_{\underline{L}_0}$. We recall that the support of the measure $P_{\underline{L}_0}$ is a minimal closed set $S_{P_{\underline{L}_0}} \subseteq H^r(D_0)$ such that $P_{S_{P_{\underline{L}_0}}} = 1$. The set $S_{P_{\underline{L}_0}}$ consists of all $\underline{g} \in H^r(D)$ such that for every neighborhood G of \underline{g} the inequality $P_{\underline{L}_0}(G) > 0$ holds.

THEOREM 10. *Suppose that $\alpha_1, \dots, \alpha_r$ are algebraically independent over \mathbb{Q} and $\text{rank}(A) = r$. Then the support of $P_{\underline{L}_0}$ is the whole of $H^r(D_0)$.*

Let

$$\widehat{L}_0(s, \omega) = (L(\lambda_1, \alpha_1, s, \omega), \dots, L(\lambda_r, \alpha_r, s, \omega)), \quad s \in D_0,$$

where

$$L(\lambda_j, \alpha_j, s, \omega) = \sum_{m=0}^{\infty} \frac{e^{2\pi i m \lambda_j \omega(m)}}{(m + \alpha_j)^s},$$

$\omega \in \Omega, j = 1, \dots, r$. Then, clearly, $S_{P_{\underline{L}_0}} \supseteq S_{P_{\widehat{\underline{L}_0}}}$. Therefore, it suffices to prove that $S_{P_{\widehat{\underline{L}_0}}} = H^r(D_0)$.

The support of each $\omega(m)$ is the unit circle γ . Thus, the support of

$$\frac{e^{2\pi i m \lambda_j \omega(m)}}{(m + \alpha_j)^s}$$

is the set

$$\{g \in H(D_0) : g(s) = \frac{e^{2\pi i m \lambda_j a}}{(m + \alpha_j)^s}, a \in \gamma\}, \quad m \in \mathbb{N}_0, j = 1, \dots, r.$$

Since $\{\omega(m) : m \in \mathbb{N}_0\}$ is a sequence of independent random variables defined on the probability space $(\Omega, \mathcal{B}(\Omega), m_H)$,

$$\left\{ \frac{e^{2\pi i m \lambda_j \omega(m)}}{(m + \alpha_j)^s} : m \in \mathbb{N}_0 \right\}, \quad j = 1, \dots, r,$$

is a sequence of independent $H(D_0)$ -valued random elements defined on the above space. Therefore, by Lemma 5 of [7] the support $S_{P_{\widehat{\underline{L}_0}}}$ is the closure of the set of all convergent series

$$\sum_{m=0}^{\infty} \left(\frac{e^{2\pi i m \lambda_1 a_m}}{(m + \alpha_1)^s}, \dots, \frac{e^{2\pi i m \lambda_r a_m}}{(m + \alpha_r)^s} \right), \quad (16)$$

where $a_m \in \gamma$. Thus, we have to show that the latter set is dense in $H^r(D_0)$. For this we will use Lemma 6 of [7].

Let $\{b_m : b_m \in \gamma, m \in \mathbb{N}_0\}$ be a sequence such that the series

$$\sum_{m=0}^{\infty} \left(\frac{e^{2\pi i m \lambda_1 b_m}}{(m + \alpha_1)^s}, \dots, \frac{e^{2\pi i m \lambda_r b_m}}{(m + \alpha_r)^s} \right)$$

converges in $H^r(D_0)$. Such a sequence exists, since

$$\sum_{m=0}^{\infty} \left(\frac{e^{2\pi i m \lambda_1 \omega(m)}}{(m + \alpha_1)^s}, \dots, \frac{e^{2\pi i m \lambda_r \omega(m)}}{(m + \alpha_r)^s} \right)$$

is an $H^r(D_0)$ - valued random element. By the definition of D_0 , for every compact subset K of D_0 ,

$$\sum_{m=0}^{\infty} \sum_{j=1}^r \sup_{s \in K} \frac{1}{(m + \alpha_j)^{2\sigma}} < \infty.$$

Therefore, it remains to verify only the hypothesis a) of Lemma 6 from [7]. Let μ_1, \dots, μ_r be complex measures on $(\mathbb{C}, \mathcal{B}(\mathbb{C}))$ with compact supports contained in D_0 and such that

$$\sum_{m=0}^{\infty} \left| \sum_{j=1}^r \int_{\mathbb{C}} \frac{e^{2\pi i m \lambda_j}}{(m + \alpha_j)^s} d\mu_j(s) \right| < \infty. \quad (17)$$

Since (see [6])

$$(m + \alpha)^{-s} = m^{-s} + O(m^{-1-\sigma} |s| e^{O(|s|)}),$$

(17) and the properties of the measures μ_1, \dots, μ_r show that

$$\sum_{m=0}^{\infty} \left| \sum_{j=1}^r \int_{\mathbb{C}} \frac{e^{2\pi i m \lambda_j}}{m^s} d\mu_j(s) \right| < \infty.$$

Hence, by the periodicity of $e^{2\pi i m \lambda_j}$, for every $l = 1, \dots, k$,

$$\sum_{\substack{m=0 \\ m \equiv l \pmod{k}}}^{\infty} \left| \sum_{j=1}^r \int_{\mathbb{C}} \frac{e^{2\pi i l \lambda_j}}{m^s} d\mu_j(s) \right| < \infty. \quad (18)$$

Define

$$\nu_l(A) = \sum_{j=1}^r e^{2\pi i l \lambda_j} \mu_j(A), \quad A \in \mathcal{B}(\mathbb{C}), l = 1, \dots, k.$$

Then, clearly, ν_1, \dots, ν_l are complex measures on $(\mathbb{C}, \mathcal{B}(\mathbb{C}))$ with compact supports obtained in D_0 , and in view of (18)

$$\sum_{\substack{m=0 \\ m \equiv l \pmod{k}}}^{\infty} \left| \int_{\mathbb{C}} m^{-s} d\nu_l(s) \right| < \infty, \quad l = 1, \dots, k. \quad (19)$$

Now we put

$$\rho_l(z) = \int_{\mathbb{C}} e^{-sz} d\nu_l(s), \quad z \in \mathbb{C}, \quad l = 1, \dots, k.$$

Then by (19)

$$\sum_{\substack{m=0 \\ m \equiv l \pmod{k}}}^{\infty} \left| \rho_l(\log m) \right| < \infty, \quad l = 1, \dots, k. \quad (20)$$

Clearly, $\rho_l(z)$ is an entire function of exponential type, $l = 1, \dots, k$. Thus, by Lemma 6.4.10 of [3], either $\rho_l \equiv 0$, or

$$\limsup_{x \rightarrow \infty} \frac{\log |\rho_l(x)|}{x} > -1, \quad l = 1, \dots, k. \quad (21)$$

Suppose that (21) is true. Then Lemma 5 of [6] (which is a version of the "positive density method") shows that

$$\sum_{\substack{m=0 \\ m \equiv l \pmod{k}}}^{\infty} \left| \rho_l(\log m) \right| = \infty, \quad l = 1, \dots, k,$$

and this contradicts (20). Therefore, we have that, for $l = 1, \dots, k$, $\rho_l(z) \equiv 0$. Hence, by the definition of the measures ν_1, \dots, ν_l ,

$$\sum_{j=1}^r e^{2\pi i l \lambda_j} \int_{\mathbb{C}} e^{-sz} d\mu_j(s) \equiv 0, \quad l = 1, \dots, k.$$

Since $\text{rank}(A) = r$, this implies

$$\int_{\mathbb{C}} e^{-sz} d\mu_j(s) \equiv 0, \quad j = 1, \dots, r,$$

and we easily deduce that

$$\int_{\mathbb{C}} s^l d\mu_j(s) \equiv 0$$

for all $j = 1, \dots, r$ and $l \in \mathbb{N}_0$. Hence we obtain that all hypotheses of Lemma 6 from [7] are satisfied, and we have that the set of all convergent series

$$\sum_{m=0}^{\infty} \left(\frac{e^{2\pi i m \lambda_1} b_m a_m}{(m + \alpha_1)^s}, \dots, \frac{e^{2\pi i m \lambda_r} b_m a_m}{(m + \alpha_r)^s} \right)$$

with $a_m \in \gamma$ is dense in $H^r(D_0)$. Clearly, the set of all convergent series (16) also has the same property. The theorem is proved.

Proof of Theorem 2. The proof uses Theorem 10 and is the same that of Theorem 1 from [6].

REFERENCES

1. P. Billingsley, *Convergence of Probability measures*, John Wiley, New York (1968).
2. H. Heyer, *Probability Measures on Locally Compact Groups*, Springer - Verlag, Berlin (1977).
3. A. Laurinćikas, *Limit Theorems for the Riemann Zeta-Function*, Kluwer, Dordrecht (1996).
4. A. Laurinćikas, R. Garunkštis, *The Lerch Zeta - Function*, Kluwer, Dordrecht (2002).

5. A. Laurinčikas, K. Matsumoto, Joint value - distribution theorems on Lerch zeta - functions, *Lith. Math. J.*, **38**(3), 238–249 (1998).
6. A. Laurinčikas, K. Matsumoto, The joint universality and the functional independence for Lerch zeta - functions, *Nagoya Math. J.*, **157**, 211–227 (2000).
7. A. Laurinčikas, K. Matsumoto, The joint universality of zeta - functions attached to certain cusp forms, *Fiz. - mat. fak. moksl. seminaro darbai*, **5**, 58–75 (2002).

Jungtinės reikšmių pasiskirstymo teoremos Lercho dzeta funkcijoms. II

A. Laurinčikas, K. Matsumoto.

Pateikti ištaisyti kai kurių teoremų iš [5] ir [6] formulavimai apie jungtinį Lercho dzeta funkcijų reikšmių pasiskirstymą (ribinės teoremos, universalumas, funkcinis nepriklausomumas). Be to, pateiktas naujas tiesioginis jungtinės ribinės teoremos analizinių funkcijų erdvėje įrodymas bei praplėsta jungtinė universalumo teorema.

*Rankraštis gautas
2006 04 00*