

---

**Homework 12**

---

**Exercise 1** Use Green's theorem to compute the integral  $\int_c f$  with  $f(x, y) = (y^2, x)$  when  $c$  corresponds to the following curves, taken counterclockwise:

- (i) The square of vertices  $(0, 0)$ ,  $(2, 0)$ ,  $(2, 2)$ ,  $(0, 2)$ ,
- (ii) The square of vertices  $(\pm 1, \pm 1)$ ,
- (iii) The circle of radius 1 and centered at  $(0, 0)$ ,
- (iv) The ellipse of equation  $(x/a)^2 + (y/b)^2 = 1$  for some  $a, b > 0$ .

**Exercise 2** Check the validity of Green's theorem for the function  $f : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  defined for  $(x, y) \in \mathbb{R}^2$  by  $f(x, y) = (2xy, x^2)$  on the domain  $\Omega = [-1, 2] \times [-1, 3] \subset \mathbb{R}^2$ .

**Exercise 3** Consider the function  $f : \mathbb{R}^2 \setminus \{0, 0\} \rightarrow \mathbb{R}^2$  defined for  $(x, y) \in \mathbb{R}^2$  by

$$f(x, y) = \left( \frac{-y}{x^2 + y^2}, \frac{x}{x^2 + y^2} \right).$$

Let  $c : [a, b] \rightarrow \mathbb{R}^2 \setminus \{(0, 0)\}$  be a parametric curve of class  $C^1$  and non-intersecting such that its interior  $\Omega$  is located on the left of the curve. We also assume that  $(0, 0) \in \Omega$ . Compute  $\int_c f$ , and explain your result.

**Exercise 4** Let  $f : \mathbb{R}^2 \rightarrow \mathbb{R}$  be sufficiently many times differentiable and satisfying the Laplace equation

$$\frac{\partial^2 f}{\partial x^2} + \frac{\partial^2 f}{\partial y^2} = 0.$$

- (i) Let  $c$  be a closed parametric curve oriented counterclockwise and non-intersecting. Show that

$$\int_c \begin{pmatrix} \partial_y f \\ -\partial_x f \end{pmatrix} = 0,$$

- (ii) Show that  $f(0, 0) = \frac{1}{2\pi} \int_0^{2\pi} f(r \cos(\theta), r \sin(\theta)) d\theta$  for any  $r > 0$ .

**Exercise 5** Compute the area of the domain defined in  $\mathbb{R}_+ \times \mathbb{R}_+$  by the four curves of equation

$$y = ax, \quad y = x/a, \quad y = b/x, \quad y = 1/bx \quad \text{for } a > 1, b > 1.$$