
Homework 3

Exercise 1 1) Compute the partial derivatives of the following functions on their respective domain:

$$a) f_1(x, y) = xy, \quad b) f_2(x, y) = (x + 1)(y + 3) \quad c) f_3(x, y) = \frac{xy}{x^2 + y^2} \quad d) f_4(x, y) = \frac{x + y}{x - y}.$$

2) Compute also the partial derivatives of the following functions:

(i) $f : \mathbb{R}_+ \times \mathbb{R} \ni (x, y) \mapsto f(x, y) = x^y \in \mathbb{R}$,

(ii) $g : (\mathbb{R}_+)^3 \rightarrow \mathbb{R}$ given by $g(x, y, z) = x^3y^2 + \sin(xz) - \ln(xyz)$.

Let us recall the polar coordinates $(r, \theta) \in [0, \infty) \times [0, 2\pi)$ which allow one to give an alternative description of all the points of \mathbb{R}^2 . The relations between the usual coordinates $(x, y) \in \mathbb{R} \times \mathbb{R}$ are

$$x = r \cos(\theta) \quad \text{and} \quad y = r \sin(\theta).$$

Exercise 2 Consider the function $f : \mathbb{R}^2 \rightarrow \mathbb{R}$ defined by

$$f(x, y) := \begin{cases} \frac{x^3}{x^2+y^2} & \text{if } (x, y) \neq (0, 0) \\ 0 & \text{if } (x, y) = (0, 0) \end{cases}$$

1. By using the polar coordinates, study the continuity of f at $(0, 0)$,
2. Show that f is differentiable with continuous partial derivatives on $\mathbb{R}^2 \setminus \{(0, 0)\}$,
3. Show that f admits in $(0, 0)$ derivatives in all directions,
4. Show that f is not differentiable at $(0, 0)$.

Exercise 3 Consider the function $f : \mathbb{R}^2 \rightarrow \mathbb{R}$ defined by

$$f(x, y) := \begin{cases} \frac{xy(x^2-y^2)}{x^2+y^2} & \text{if } (x, y) \neq (0, 0) \\ 0 & \text{if } (x, y) = (0, 0) \end{cases}$$

and compute its second derivatives $\partial_j \partial_k f$ for any $j, k \in \{x, y\}$. Are these functions continuous ?

Exercise 4 Consider the functions $\varphi : \mathbb{R} \rightarrow \mathbb{R}^2$ defined by $\varphi(t) := \begin{pmatrix} \cos(t) \\ t^2 \end{pmatrix}$, and $f : \mathbb{R}^2 \rightarrow \mathbb{R}$ defined by $f(x, y) := e^{3x+2y}$. We consider the composition of these two functions, namely $F : \mathbb{R} \rightarrow \mathbb{R}$ given by $F = f \circ \varphi$. Compute the derivative of this function by two different methods: once by a direct computation, and once as the derivative of a composed function (chain rule).